

2021

## Estimation of Stress-Strength Parameter for Rayleigh Distribution based on Progressive Type-II Censoring

M. H. Abu-Moussa

*Department of Mathematics, Faculty of Science, Cairo University, Giza, Egypt,*  
eslmhussam1986@gmail.com

A. M. Abd-Elfattah

*Department of Mathematical Statistics, Institute of Statistical Studies-Research Cairo University, Egypt,*  
eslmhussam1986@gmail.com

E. H. Hafez

*Department of Mathematics, Faculty of Science Helwan University, Cairo, Egypt,*  
eslmhussam1986@gmail.com

Follow this and additional works at: <https://digitalcommons.aaru.edu.jo/isl>

---

### Recommended Citation

H. Abu-Moussa, M.; M. Abd-Elfattah, A.; and H. Hafez, E. (2021) "Estimation of Stress-Strength Parameter for Rayleigh Distribution based on Progressive Type-II Censoring," *Information Sciences Letters*: Vol. 10 : Iss. 1 , Article 12.

Available at: <https://digitalcommons.aaru.edu.jo/isl/vol10/iss1/12>

This Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in *Information Sciences Letters* by an authorized editor. The journal is hosted on [Digital Commons](#), an Elsevier platform. For more information, please contact [rakan@aarj.edu.jo](mailto:rakan@aarj.edu.jo), [marah@aarj.edu.jo](mailto:marah@aarj.edu.jo), [dr\\_ahmad@aarj.edu.jo](mailto:dr_ahmad@aarj.edu.jo).

# Estimation of Stress-Strength Parameter for Rayleigh Distribution based on Progressive Type-II Censoring

M. H. Abu-Moussa<sup>1</sup>, A. M. Abd-Elfattah<sup>2</sup> and E. H. Hafez<sup>3\*</sup>

<sup>1</sup>Department of Mathematics, Faculty of Science, Cairo University, Giza, Egypt

<sup>2</sup>Department of Mathematical Statistics, Institute of Statistical Studies-Research Cairo University, Egypt

<sup>3</sup>Department of Mathematics, Faculty of Science Helwan University, Cairo, Egypt

Received: 7 July. 2020, Revised: 23 Nov. 2020, Accepted: 29 Nov. 2020

Published online: 1 Jan. 2021

**Abstract:** In this paper, the estimation of stress-strength parameter  $R = P(Y < X)$  is considered when  $X, Y$  the strength and stress respectively are two independent random variables of Rayleigh distribution. The samples taken for  $X$  and  $Y$  are progressively censoring of type II. Maximum likelihood estimator (MLE), uniformly minimum variance unbiased estimator (UMVUE) and Bayes estimator of  $R = P(Y < X)$  are obtained. The exact confidence interval of  $R$  based on MLE is obtained. The performance of the proposed estimators is compared using computer simulation.

**Keywords:** Rayleigh distribution; progressive type-II censoring; stress-strength model; unbiased estimator; maximum-likelihood estimator.

## 1 Introduction

Rayleigh distribution was first considered by Lord Rayleigh [1]. It used in modeling lifetime data and reliability. The Rayleigh distribution has the following distribution function for  $X > 0$ :

$$F(x) = 1 - e^{-x^2/2\sigma^2}; \quad \sigma > 0, x \geq 0 \quad (1.1)$$

where, the density function of Rayleigh for  $X > 0$  denoted by Ray ( $\sigma$ ) is

$$f(x; \sigma) = \frac{x}{\sigma^2} e^{-x^2/2\sigma^2}; \quad \sigma > 0 \quad (1.2)$$

Rayleigh distribution is related to other distributions such as Weibull distribution where Rayleigh ( $\sigma$ ) = Weibull ( $2, \sigma\sqrt{2}$ ). In life-testing experiments, one often encounters situations where it takes a substantial amount of time to obtain a reasonable number of failures necessary to carry out reliable inference, so censored samples are used for analyzing lifetime data. Among various censoring schemes, the Type-II progressive censoring scheme has become very popular one in the last decade. It can be described as follows: let  $n$  units be placed on test at time zero with  $m$  failures to be observed. At the first failure a number  $r_1$  of the surviving units ( $n - 1$ ) are randomly selected and removed from the experiment. At the second observed failure,  $r_2$  of the surviving units ( $n - r_1 - 2$ ) are randomly selected and removed from the experiment, and so on until the  $m$ -th failure is observed in which all remaining surviving units  $r_m = n - m - r_1 - r_2 - \dots - r_{m-1}$  are removed. We denote to progressively Type-II censoring with scheme  $(n, m, r_1, r_2, \dots, r_m)$ . Traditional Type-II censoring scheme is included when  $(r_1 = r_2 = \dots = r_{m-1} = 0)$  and  $(r_m = n - m)$  and complete sampling scheme when  $(n = m)$  and  $(r_1 = \dots = r_{m-1} = r_m = 0)$ . Balakrishnan and Aggarwala [2] and Balakrishnan [3] presented a study on different features of progressive censoring schemes.

In stress-strength model, the stress ( $Y$ ) and the strength ( $X$ ) are treated as random variables and the reliability of a component during a given period is taken to be the probability that its strength exceeds the stress during the entire interval, i.e. the reliability  $R$  of a component is  $R = P(Y < X)$ . For a particular situation, if we consider  $Y$  as the pressure of a chamber generated by ignition of a solid propellant and  $X$  as the strength of the chamber. Then  $R$  represents the probability of successful firing of the engine. Stress-strength model can be used as a general measure of the difference between two populations and has applications in many areas. For example, comparing two treatments  $X$  and  $Y$ , then  $R = P(Y < X)$  is the measure of the response of treatment  $X$ . For other applications see Kotz et al. [4]. Many authors considered the problem

\*Corresponding author e-mail: [eslmhussam1986@gmail.com](mailto:eslmhussam1986@gmail.com)

of estimating the stress-strength parameter based on complete samples, it first considered by Birnbaum [5]. Johnson [6] present a good review on stress-strength model in reliability. Awad and Charraf [7] studied the case when  $X$  and  $Y$  are independent Burr random variables of type XII, they obtained maximum likelihood, uniformly minimum variance unbiased estimator (UMVUE) and Bayesian estimates of  $R$ . Fathipour et al. [8] obtained the estimation of  $R$  in the generalized Rayleigh distribution with different scale parameters. Ahmed et al. [9] consider this problem when  $X$  and  $Y$  are two independent random variables have Burr Type X distribution. Based on censored samples, Saraçoğlu et al. [10] obtained the estimation for  $R$  based on exponential distribution with type II progressive censoring. Abd-Elfattah et al. [11] get the estimation of  $R$  based on Burr type XII distribution with type II progressive censoring, they discussed two cases the first when  $X$  and  $Y$  have common shape parameter and different scale parameters while the second case when  $X$  and  $Y$  have common scale parameter and different shape parameters. For some of the recent references, the readers may refer to [12-14].

In the present paper, the study the estimation of  $R = P(Y < X)$  when  $X$  and  $Y$  are two independents but not identically random variables belonging to Rayleigh distribution with parameters  $\sigma_1$  and  $\sigma_2$  respectively. In Section (2), maximum likelihood estimator of  $R$  and exact confidence interval are obtained. Also, UMVUE of  $R$  is obtained in section (3). In section (4) Bayes estimator for  $R$  is obtained based on conjugate priors. Numerical results and simulations are presented in section (5). Finally, some concluding remarks are given in section (6).

## 2 MLE of $R$

In this section the MLE of  $R$  is obtained. Let  $X, Y$  are two independent random variables such that  $X \sim Ray(\sigma_1)$  and  $Y \sim Ray(\sigma_2)$  then  $R$  is:

$$\begin{aligned} R &= P(Y < X) = \int_0^{\infty} \int_0^x f(x)f(y)dydx \\ &= \int_0^{\infty} \int_0^x \frac{x}{\sigma_1^2} e^{-x^2/2\sigma_1^2} \cdot \frac{y}{\sigma_2^2} e^{-y^2/2\sigma_2^2} dydx \\ &= \frac{\sigma_1^2}{\sigma_1^2 + \sigma_2^2} \end{aligned} \quad (2.1)$$

Now to get MLE of  $R$  we first get the MLE of parameters  $\sigma_1$  and  $\sigma_2$ .

Let  $X_{1:m_1:n_1}, \dots, X_{m_1:m_1:n_1}$  be a progressive censored sample from  $Ray(\sigma_1)$  with progressive censoring scheme  $(n_1, m_1, r_1, \dots, r_{m_1})$ , and let  $Y_{1:m_2:n_2}, \dots, Y_{m_2:m_2:n_2}$  be a progressive censored sample from  $Ray(\sigma_2)$  with progressive censoring scheme  $(n_2, m_2, s_1, \dots, s_{m_2})$ , then the jointly likelihood function  $L(\sigma_1, \sigma_2)$  is

$$\begin{aligned} L(\sigma_1, \sigma_2) &= [k_1 \prod_{i=1}^{m_1} f(x_i) [1 - F(x_i)]^{r_i}] \cdot [k_2 \prod_{j=1}^{m_2} f(y_j) [1 - F(y_j)]^{s_j}] \\ &= \frac{k_1 k_2}{\sigma_1^{2m_1} \sigma_2^{2m_2}} (\prod_{i=1}^{m_1} x_i) (\prod_{j=1}^{m_2} y_j) \times \exp\left(\frac{-1}{2\sigma_1^2} \sum_{i=1}^{m_1} (1 + r_i) x_i^2\right) \exp\left(\frac{-1}{2\sigma_2^2} \sum_{j=1}^{m_2} (1 + s_j) y_j^2\right) \end{aligned} \quad (2.2)$$

Where  $k_1$  and  $k_2$  are:

$$\begin{aligned} k_1 &= n_1(n_1 - 1 - r_1)(n_1 - 2 - r_1 - r_2) \dots (n_1 - m_1 + 1 - r_1 - \dots - r_{m_1-1}) \\ k_2 &= n_2(n_2 - 1 - s_1)(n_2 - 2 - s_1 - s_2) \dots (n_2 - m_2 + 1 - s_1 - \dots - s_{m_2-1}) \end{aligned} \quad (2.3)$$

Now the jointly log-likelihood function  $\ell$  is:

$$\ell = \ln(k_1 k_2) - 2m_1 \ln \sigma_1 - 2m_2 \ln \sigma_2 + \sum_{i=1}^{m_1} \ln x_i + \sum_{j=1}^{m_2} \ln y_j - \frac{1}{2\sigma_1^2} \sum_{i=1}^{m_1} (1 + r_i) x_i^2 - \frac{1}{2\sigma_2^2} \sum_{j=1}^{m_2} (1 + s_j) y_j^2 \quad (2.4)$$

By differentiation on equation (2.4) with respect to  $\sigma_1$  and  $\sigma_2$ . Then we get:

$$\frac{\partial \ell}{\partial \sigma_1} = -\frac{2m_1}{\sigma_1} + \frac{1}{\sigma_1^3} \sum_{i=1}^{m_1} (1 + r_i) x_i^2 \quad (2.5)$$

$$\frac{\partial \ell}{\partial \sigma_2} = -\frac{2m_2}{\sigma_2} + \frac{1}{\sigma_2^3} \sum_{j=1}^{m_2} (1 + s_j) y_j^2 \quad (2.6)$$

By putting equations (2.5) and (2.6) equal to zero then we get:

$$\hat{\sigma}_1 = \left[ \frac{1}{2m_1} \sum_{i=1}^{m_1} (1 + r_i) x_i^2 \right]^{0.5} \quad (2.7)$$

$$\hat{\sigma}_2 = \left[ \frac{1}{2m_2} \sum_{j=1}^{m_2} (1 + s_j) y_j^2 \right]^{0.5} \tag{2.8}$$

Then MLE of R is

$$\hat{R} = \frac{1}{1 + \frac{m_1 E}{m_2 T}} = \frac{1}{1 + \frac{\sigma_2^2 \lambda}{\sigma_1^2}} \tag{2.9}$$

where  $T = \sum_{i=1}^{m_1} (1 + r_i) x_i^2$ ,  $E = \sum_{j=1}^{m_2} (1 + s_j) y_j^2$  and  $\lambda = \frac{\sigma_1^2 m_1 E}{\sigma_2^2 m_2 T}$ .

consider the following transformations:

$$\begin{aligned} Z_1 &= n_1 X_1^2 \\ Z_2 &= (n_1 - r_1 - 1) [X_2^2 - X_1^2] \\ &\vdots \\ &\vdots \\ Z_{m_1} &= (n_1 - r_1 - r_2 - \dots - r_{m_1-1} - m_1 + 1) [X_{m_1}^2 - X_{m_1-1}^2] \end{aligned} \tag{2.10}$$

Balakrishnan and Aggarwala [2] show that  $Z_i$ 's are independent and identically distributed exponential random variables with mean  $\sigma_1$ , moreover

$$T = \sum_{i=1}^{m_1} Z_i = \sum_{i=1}^{m_1} (1 + r_i) T_i = \sum_{i=1}^{m_1} (1 + r_i) X_i^2 \sim \text{Gamma}(m_1, \sigma_1). \tag{2.11}$$

Similarly,  $E$  has a gamma distribution with the shape parameter  $m_1$  and scale parameter  $\sigma_1$ . Hence,  $\lambda$  follow F-distribution with degrees of freedom  $2m_2$  and  $2m_1$ . Using the fact That:

$$\frac{R}{1-R} \times \frac{1-\hat{R}}{\hat{R}} \sim F_{2m_2, 2m_1} \tag{2.12}$$

Then the  $100(1 - \alpha)\%$  exact confidence interval of R is:

$$P\left(\frac{1}{1 + F_{2m_1, 2m_2, \alpha/2}(\frac{1}{\hat{R}} - 1)} < R < \frac{1}{1 + F_{2m_1, 2m_2, 1-\alpha/2}(\frac{1}{\hat{R}} - 1)}\right) = 1 - \alpha$$

Where  $\alpha$  is the level of significance and  $2m_1, 2m_2$  are the degree of freedom of F.

### 3 UMVUE of R

In this section the uniformly minimum variance unbiased estimator (UMVUE) is obtained for stress-strength parameter R. Let  $X_{1:m_1:n_1}, \dots, X_{m_1:m_1:n_1}$  be a progressive censored sample from  $Ray(\sigma_1)$  with progressive censoring scheme  $(n_1, m_1, r_1, \dots, r_{m_1})$ , and the progressive censored sample  $Y_{1:m_2:n_2}, \dots, Y_{m_2:m_2:n_2}$  from  $Ray(\sigma_2)$  with progressive censoring scheme  $(n_2, m_2, s_1, \dots, s_{m_2})$ , then from the jointly log-likelihood function of X, Y that given in equation (2.4), we obtained that  $T = \sum_{i=1}^{m_1} (1 + r_i) x_i^2$  is a sufficient statistics for  $\sigma_1$  and  $E = \sum_{j=1}^{m_2} (1 + s_j) y_j^2$  is a sufficient statistics for  $\sigma_2$ . From equation (2.11)

$$f_T(t) = \frac{1}{\sigma_1^{m_1} \Gamma(m_1)} t^{m_1-1} \exp\left(-\frac{t}{\sigma_1}\right), \quad 0 < t < \infty \tag{3.1}$$

**Lemma 3.1** The conditional pdf. of  $T_1 = X_1^2$  given T is:

$$f_{T_1|T}(x) = \frac{f_{T_1, T}(x)}{f_T(t)} = n_1 (m_1 - 1) \frac{(T - n_1 T_1)^{m_1-2}}{T^{m_1-1}}, \quad 0 < T_1 < T/n_1 \tag{3.2}$$

**Proof.** Let  $W = \sum_{i=2}^{m_1} Z_i$  then clearly  $W$  &  $Z_1$  are independent. Then the joint pdf. of  $T_1$  &  $T$ ,  $f_{T_1, T}(x)$  can be easily obtained from the jointly distribution of  $W$  &  $Z_1$  using the transformations  $Z_1 = n_1 T_1$  &  $W = T - Z_1 = T - T_1$  then

$$f_{W, Z_1} = f_W \cdot f_{Z_1} = \frac{1}{\sigma_1^{m_1} \Gamma(m_1-1)} W^{m_1-2} \exp\left(-\frac{W+Z_1}{\sigma_1}\right) \tag{3.3}$$

And

$$f_{T_1, T} = \frac{n_1}{\sigma_1^{m_1} \Gamma(m_1-1)} (T - n_1 T_1)^{m_1-2} \exp\left(-\frac{T}{\sigma_1}\right) \tag{3.4}$$

From equations (3.4), (3.1), we get the result.

Similarly, if  $E = \sum_{j=1}^{m_2} (1 + s_j) E_j$  where  $E_j = Y_j^2$  Then

$$f_{E_1|E}(y) = n_2(m_2 - 1) \frac{(E - n_2 E_1)^{m_2 - 2}}{E^{m_2 - 1}}, \quad 0 < E_1 < E/n_2 \quad (3.5)$$

**Lemma 3.2** The unbiased estimator of  $R$  is:

$$\varphi(T_1, E_1) = \begin{cases} 1 & \text{if } n_2 E_1 < n_1 T_1 \\ 0 & \text{if } n_2 E_1 \geq n_1 T_1 \end{cases} \quad (3.6)$$

Where  $E_1 = Y_1^2$  and  $T_1 = X_1^2$ .

**Proof.**

$$\begin{aligned} E(\varphi) &= 1. P(n_2 E_1 < n_1 T_1) = P(Y_1 < \left[\frac{n_1 X_1^2}{n_2}\right]^{1/2}) = P(Y_1 < a) \\ &= \int_0^\infty \int_0^a f_{X_1}(x) f_{Y_1}(y) dy dx \end{aligned} \quad (3.7)$$

Where the distributions of order statistics  $X_1$  and  $Y_1$  are

$$\begin{aligned} f_{X_1}(x) &= n_1 \frac{x}{\sigma_1^2} e^{-n_1 x^2 / 2\sigma_1^2} \\ f_{Y_1}(y) &= n_2 \frac{y}{\sigma_2^2} e^{-n_2 y^2 / 2\sigma_2^2} \end{aligned} \quad (3.8)$$

Then by using equations (3.8) we get

$$E(\varphi) = \frac{\sigma_1^2}{\sigma_1^2 + \sigma_2^2} = R \quad (3.9)$$

**Theorem 3.3** Based on the sufficient statistics  $T$  and  $E$ , as defined before for  $\sigma_1$  and  $\sigma_2$  respectively and the unbiased statistics  $\varphi$ , the UMVUE of  $R$ , say  $\tilde{R}$ , for  $m_1 \geq 2$  and  $m_2 \geq 2$  can be expressed as follows:

$$\tilde{R} = \begin{cases} 1 - \sum_{k=0}^{m_2-1} (-1)^k \left(\frac{T}{E}\right)^k \frac{\binom{m_2-1}{k}}{\binom{m_1+k-1}{k}} & \text{if } T < E \\ \sum_{k=0}^{m_1-1} (-1)^k \left(\frac{E}{T}\right)^k \frac{\binom{m_1-1}{k}}{\binom{m_2+k-1}{k}} & \text{if } T \geq E \end{cases} \quad (3.10)$$

**Proof.** For  $T < E$  using the Rao-Blackwell theorem

$$\tilde{R} = E(\varphi(T_1, E_1) | T, E) = \int \int_A f_{(T_1|T)} f_{(E_1|E)} dE_1 dT_1, \quad (3.11)$$

Where  $A = \{(E_1, T_1) : 0 < T_1 < \frac{T}{n_1}, 0 < E_1 < \frac{E}{n_2} \text{ and } n_2 E_1 < n_1 T_1\}$  and  $f_{(T_1|T)}$  &  $f_{(E_1|E)}$  are defined in equations (3.2), (3.5) respectively. Then  $\tilde{R}$  becomes:

$$\begin{aligned} \tilde{R} &= \int_0^{T/n_1} \int_0^{n_1 T_1/n_2} n_1 (m_1 - 1) \frac{(T - n_1 T_1)^{m_1 - 2}}{T^{m_1 - 1}} \cdot n_2 (m_2 - 1) \frac{(E - n_2 E_1)^{m_2 - 2}}{E^{m_2 - 1}} dE_1 dT_1 \\ &= 1 - \int_0^{T/n_1} n_1 (m_1 - 1) \frac{(T - n_1 T_1)^{m_1 - 2}}{T^{m_1 - 1}} \frac{(E - n_1 T_1)^{m_2 - 1}}{E^{m_2 - 1}} dT_1 \end{aligned} \quad (3.12)$$

let  $c = \frac{n_1 T_1}{T}$ , then  $\tilde{R}$  becomes:

$$\tilde{R} = 1 - \int_0^1 (m_1 - 1) (1 - c)^{m_1 - 2} \left(1 - c \frac{T}{E}\right)^{m_2 - 1} dc, \quad (3.13)$$

since the binomial expansion of  $(1 - c \frac{T}{E})^{m_2 - 1} = \sum_{k=0}^{m_2 - 1} (-1)^k \binom{m_2 - 1}{k} \left(\frac{cT}{E}\right)^k$ , then  $\tilde{R}$  is obtained as following:

$$\begin{aligned} \tilde{R} &= 1 - \sum_{k=0}^{m_2 - 1} (-1)^k \binom{m_2 - 1}{k} \left(\frac{T}{E}\right)^k \int_0^1 c^k (1 - c)^{m_1 - 2} dc \\ &= 1 - \sum_{k=0}^{m_2 - 1} (-1)^k \left(\frac{T}{E}\right)^k \frac{\binom{m_2 - 1}{k}}{\binom{m_1 + k - 1}{k}} \end{aligned} \quad (3.14)$$

Similarly, if  $T \geq E$  then  $\tilde{R}$  becomes:

$$\tilde{R} = \sum_{k=0}^{m_1-1} (-1)^k \left(\frac{E}{T}\right)^k \frac{\binom{m_1-1}{k}}{\binom{m_2+k-1}{k}} \tag{3.15}$$

### 4 Bayes Estimator of R

In this section, the Bayes estimator of  $R = P(Y < X)$  is obtained based on the progressive type-II censoring. Let  $X_{1:m_1:n_1}, \dots, X_{m_1:m_1:n_1}$  be a progressive censored sample from  $Ray(\sigma_1)$  with progressive censoring scheme  $(n_1, m_1, r_1, \dots, r_{m_1})$ , and the progressive censored sample  $Y_{1:m_2:n_2}, \dots, Y_{m_2:m_2:n_2}$  from  $Ray(\sigma_2)$  with progressive censoring scheme  $(n_2, m_2, s_1, \dots, s_{m_2})$ . Assuming that the parameters  $\sigma_1, \sigma_2$  are random variables having a gamma prior distribution, for simplicity we assume that the parameters  $\lambda_1 = \sigma_1^2$  and  $\lambda_2 = \sigma_2^2$ . The conjugate priors for  $\lambda_1, \lambda_2$  are:

$$\pi(\lambda_1) = \frac{\beta_1^{\alpha_1}}{\Gamma(\alpha_1)} \lambda_1^{-\alpha_1+1} e^{-\beta_1/\lambda_1} \tag{4.1}$$

and

$$\pi(\lambda_2) = \frac{\beta_2^{\alpha_2}}{\Gamma(\alpha_2)} \lambda_2^{-\alpha_2+1} e^{-\beta_2/\lambda_2} \tag{4.2}$$

The likelihood function  $L(\lambda_1) = f(X|\lambda_1)$  for random variable X is:

$$f(X|\lambda_1) = \frac{k_1}{\lambda_1^{m_1}} (\prod_{i=1}^{m_1} x_i) \exp\left[-\frac{1}{2\lambda_1} \sum_{i=1}^{m_1} (1+r_i) x_i^2\right] \tag{4.3}$$

and the marginal distribution for  $X \geq 0$  is given by:

$$\begin{aligned} f(X) &= \int_0^\infty f(X|\lambda_1) \pi(\lambda_1) d\lambda_1 \\ &= \frac{k_1 \beta_1^{\alpha_1} \Gamma(\alpha_1+m_1-2)}{A^{\alpha_1+m_1-2} \Gamma(\alpha_1)} (\prod_{i=1}^{m_1} x_i) \end{aligned} \tag{4.4}$$

where  $A = \beta_1 + 0.5 \sum_{i=1}^{m_1} (1+r_i) x_i^2$ , then the posterior distribution of  $\lambda_1$  is given by:

$$\pi(\lambda_1|X) = \frac{A^{\alpha_1+m_1-2} e^{-A/\lambda_1}}{\lambda_1^{\alpha_1+m_1-1} \Gamma(\alpha_1+m_1-2)} \tag{4.5}$$

Similarly, for  $Y$  the posterior distribution of  $\lambda_2$  is given by:

$$\pi(\lambda_2|Y) = \frac{B^{\alpha_2+m_2-2} e^{-B/\lambda_2}}{\lambda_2^{\alpha_2+m_2-1} \Gamma(\alpha_2+m_2-2)} \tag{4.6}$$

where  $B = \beta_2 + 0.5 \sum_{j=1}^{m_2} (1+s_j) y_j^2$ . Now the joint posterior distribution for  $\lambda_1, \lambda_2$  is given by:

$$\pi(\lambda_1, \lambda_2|X, Y) = H \frac{e^{-A/\lambda_1} e^{-B/\lambda_2}}{\lambda_1^{\alpha_1+m_1-1} \lambda_2^{\alpha_2+m_2-1}} \tag{4.7}$$

where  $H = \frac{A^{\alpha_1+m_1-2} B^{\alpha_2+m_2-2}}{\Gamma(\alpha_1+m_1-2) \Gamma(\alpha_2+m_2-2)}$ . Now by taking the following transformations  $R = \frac{\lambda_1}{\lambda_1 + \lambda_2}$  and  $\xi = \lambda_1 + \lambda_2$

where  $0 \leq R \leq 1, \xi > 0$ , then the joint posterior distribution for  $R$  and  $\xi$  is derived as following:

$$\pi(R, \xi|X, Y) = H \xi^{3-m_1-m_2-\alpha_1-\alpha_2} \frac{\exp\left[-\frac{1}{\xi} \left(\frac{A}{R} + \frac{B}{(1-R)}\right)\right]}{R^{\alpha_1+m_1-1} (1-R)^{\alpha_2+m_2-1}} \tag{4.8}$$

Then by applying integration on equation 4.8 with respect to  $\xi$ , we get the posterior distribution for  $R$

$$\pi(R|X, Y) = H \times \Gamma[m_1 + m_2 + \alpha_1 + \alpha_2 - 4] \frac{R^{\alpha_2+m_2-3} (1-R)^{\alpha_1+m_1-3}}{[A(1-R)+B.R]^{m_1+m_2+\alpha_1+\alpha_2-4}} \tag{4.9}$$

Then under the squared error loss function the Bayes estimator of  $R$  is given by:

$$\begin{aligned} \hat{R}_3 &= \int_0^1 R \cdot \pi(R|X, Y) dR \\ &= H \cdot \Gamma[m_1 + m_2 + \alpha_1 + \alpha_2 - 4] \int_0^1 \frac{R^{\alpha_2+m_2-2} (1-R)^{\alpha_1+m_1-3}}{[A(1-R)+B.R]^{m_1+m_2+\alpha_1+\alpha_2-4}} dR \end{aligned} \tag{4.10}$$

### 5 Simulation Study

Within this section, the Monte Carlo simulation is performed to check the performance of the different estimators of R under several types of progressive censoring schemes. Samples are generated under progressive type-II censoring with many different schemes for the (n-m) removed items. These schemes are described as follows: Scheme I: complete sample (n=m) i.e there is no removed items. Scheme II: ( $r_1 = 0, \dots, r_{m-1} = 0, r_m = n - m$ ). Scheme III: ( $r_1 = n - m, \dots, r_{m-1} = 0, r_m = 0$ ). Scheme IV: The remaining items (n-m) are removed equally at each failure time. For example, if n=10 and m=5 then scheme IV become ( $r_1 = 1, r_2 = 1, \dots, r_5 = 1$ ). Simulation is performed 1000 times with different sample sizes  $n_1, n_2 = 10, 20, 30$  and the number of failures  $m_1, m_2 = 5, 10, 15, 20, 30$  for X and Y. Different values of parameters  $(\sigma_1, \sigma_2) = (1,1), (1,2), (2,1)$  for the distributions of X and Y are used. The average estimates of MLE for R and the average of MSE's are reported in Table 1, 2, 3 for each set of parameters. With each table the 95% exact confidence interval of R. Also, simulation is constructed 1000 times for UMVUE of R suggested in Section (3) at each set of values in table 4. Finally in table 5 the Bayes estimator of R is obtained using different values for the parameters  $\alpha_1, \alpha_2, \beta_1$  and  $\beta_2$  when  $n_1, n_2 = 20, 30$  and  $m_1, m_2 = 10, 15$ . All tables are founded in the appendix. We note that in such cases as the effective sample size increases the estimates of R become better. When  $n = m$  i.e in case of complete samples the biased is decreased. Also, when  $(n_1, m_1) = (n_2, m_2)$  the estimates are good. We note that MLE of R give results better than the UMVUE of R.

**Table 1:** MLE, MSE and Exact 95% C.I for R=0.5 when  $\sigma_1 = \sigma_2 = 1$  with different scheme types r, s.

$(n_1, m_1)$	$(n_2, m_2)$	r	s	MLE		95%C.I.	
				$\hat{R}_1$	MSE	Lower	Upper
(10,10)	(10,10)	Comp	Comp	0.499078	$1.1 \cdot 10^{-2}$	0.28789	0.71060
(10,5)	(10,5)	II	II	0.501915	$2.205 \cdot 10^{-2}$	0.213291	0.789268
		III	III	0.498651	$1.30 \cdot 10^{-2}$	0.211109	0.787089
		IV	IV	0.506229	$1.34 \cdot 10^{-2}$	0.216201	0.792125
(20,20)	(10,10)	Comp	Comp	0.50448	$7.95 \cdot 10^{-3}$	0.26163	0.677949
(10,10)	(20,20)	Comp	Comp	0.49126	$8.92 \cdot 10^{-3}$	0.315959	0.732939
(20,10)	(10,5)	II	II	0.516308	$1.79 \cdot 10^{-2}$	0.237949	0.74752
		III	III	0.512684	$9.64 \cdot 10^{-3}$	0.235328	0.744772
		IV	IV	0.527223	$1.10 \cdot 10^{-2}$	0.245972	0.755686
(10,5)	(20,10)	II	II	0.483445	$1.81 \cdot 10^{-2}$	0.252294	0.761872
		III	III	0.486932	$1.03 \cdot 10^{-2}$	0.254936	0.764395
		IV	IV	0.467997	$1.09 \cdot 10^{-2}$	0.240789	0.750452
(20,10)	(20,10)	II	II	0.502365	$1.18 \cdot 10^{-2}$	0.290589	0.713296
		III	III	0.500906	$6.55 \cdot 10^{-3}$	0.289388	0.712101
		IV	IV	0.499608	$6.44 \cdot 10^{-3}$	0.288321	0.711035
(20,20)	(20,20)	Comp	Comp	0.49969	$6.08 \cdot 10^{-3}$	0.347527	0.651923
(30,15)	(30,15)	II	II	0.49626	$8.27 \cdot 10^{-3}$	0.322042	0.671396
		III	III	0.50307	$4.16 \cdot 10^{-3}$	0.328014	0.677373
		IV	IV	0.498426	$4.58 \cdot 10^{-3}$	0.323934	0.673302
(30,30)	(30,30)	Comp	Comp	0.50059	$3.94 \cdot 10^{-3}$	0.37554	0.62557

**Table 2:** MLE, MSE and Exact 95% C.I for R=0.2 when  $(\sigma_1, \sigma_2) = (1,2)$  with different scheme types r, s.

$(n_1, m_1)$	$(n_2, m_2)$	r	s	MLE		95%C.I.	
				$\hat{R}_1$	MSE	Lower	Upper
(10,10)	(10,10)	Comp	Comp	0.21291	$5.89 \cdot 10^{-3}$	0.0989044	0.399994
(10,5)	(10,5)	II	II	0.21925	$1.29 \cdot 10^{-2}$	0.0702452	0.510697
		III	III	0.21216	$5.74 \cdot 10^{-3}$	0.0675582	0.500226
		IV	IV	0.208308	$6.16 \cdot 10^{-3}$	0.0661114	0.494427
(20,20)	(10,10)	Comp	Comp	0.21169	$4.33 \cdot 10^{-3}$	0.08547	0.35701
(10,10)	(20,20)	Comp	Comp	0.19994	$3.72 \cdot 10^{-3}$	0.10783	0.41797
(20,10)	(10,5)	II	II	0.22161	$9.23 \cdot 10^{-3}$	0.07687	0.44124
		III	III	0.21439	$4.98 \cdot 10^{-3}$	0.07392	0.430831
		IV	IV	0.22960	$6.47 \cdot 10^{-3}$	0.08018	0.452546
(10,5)	(20,10)	II	II	0.201719	$7.44 \cdot 10^{-3}$	0.08349	0.463472
		III	III	0.19692	$4.26 \cdot 10^{-3}$	0.08123	0.45601
		IV	IV	0.19005	$4.27 \cdot 10^{-3}$	0.07799	0.445103
(20,10)	(20,10)	II	II	0.20661	$5.19 \cdot 10^{-3}$	0.09557	0.39091
		III	III	0.20345	$3.04 \cdot 10^{-3}$	0.09390	0.38630
		IV	IV	0.20333	$2.77 \cdot 10^{-3}$	0.093843	0.38613
(20,20)	(20,20)	Comp	Comp	0.20485	$2.85 \cdot 10^{-3}$	0.12079	0.32573
(30,15)	(30,15)	II	II	0.20828	$3.79 \cdot 10^{-3}$	0.11257	0.35301
		III	III	0.20735	$2.01 \cdot 10^{-3}$	0.1120	0.35171
		IV	IV	0.20148	$1.91 \cdot 10^{-3}$	0.10846	0.34353
(30,30)	(30,30)	Comp	Comp	0.20296	$1.79 \cdot 10^{-3}$	0.13253	0.29797

**Table 3:** MLE, MSE and Exact 95% C.I for R=0.8 when  $(\sigma_1, \sigma_2) = (2,1)$  with different scheme types r, s.

$(n_1, m_1)$	$(n_2, m_2)$	r	s	MLE		95%C.I.	
				$\hat{R}_1$	MSE	Lower	Upper
(10,10)	(10,10)	Comp	Comp	0.78995	$5.84 \cdot 10^{-3}$	0.60411	0.90261
(10,5)	(10,5)	II	II	0.779165	$1.2 \cdot 10^{-2}$	0.48699	0.92915
		III	III	0.78699	$6.18 \cdot 10^{-3}$	0.4985	0.93212
		IV	IV	0.78757	$6.36 \cdot 10^{-3}$	0.49937	0.93234
(20,20)	(10,10)	Comp	Comp	0.79852	$3.86 \cdot 10^{-3}$	0.57973	0.89124
(10,10)	(20,20)	Comp	Comp	0.78893	$4.45 \cdot 10^{-3}$	0.64383	0.91482
(20,10)	(10,5)	II	II	0.79655	$7.60 \cdot 10^{-3}$	0.53386	0.91568
		III	III	0.79974	$4.12 \cdot 10^{-3}$	0.53878	0.917195
		IV	IV	0.81033	$4.22 \cdot 10^{-3}$	0.55551	0.92218
(10,5)	(20,10)	II	II	0.780636	$8.89 \cdot 10^{-3}$	0.56198	0.924043
		III	III	0.78366	$4.66 \cdot 10^{-3}$	0.56634	0.9246
		IV	IV	0.773707	$6.23 \cdot 10^{-3}$	0.55211	0.92119
(20,10)	(20,10)	II	II	0.78873	$5.14 \cdot 10^{-3}$	0.60236	0.90197
		III	III	0.79276	$2.81 \cdot 10^{-3}$	0.60818	0.9041
		IV	IV	0.79387	$3.31 \cdot 10^{-3}$	0.60979	0.90469





(20,20)	(20,20)	Comp	Comp	0.79693	$2.53 \cdot 10^{-3}$	0.67666	0.88037
(30,15)	(30,15)	II	II	0.795464	$3.42 \cdot 10^{-3}$	0.652201	0.889696
		III	III	0.798036	$1.69 \cdot 10^{-3}$	0.655795	0.891244
		IV	IV	0.79720	$1.68 \cdot 10^{-3}$	0.654632	0.89074
(30,30)	(30,30)	Comp	Comp	0.79717	$1.67 \cdot 10^{-3}$	0.70220	0.86756

**Table 4:** UMVUE  $\hat{R}_2$  and MSE for R at different sets of values of  $\sigma_1, \sigma_2$  with different scheme types r, s.

$(n_1, m_1)$	$(n_2, m_2)$	r	s	$(\sigma_1, \sigma_2) = (1,1)$ R=0.5 $\hat{R}_2$ MSE		$(\sigma_1, \sigma_2) = (1,2)$ R=0.2 $\hat{R}_2$ MSE		$(\sigma_1, \sigma_2) = (2,1)$ R=0.8 $\hat{R}_2$ MSE	
				(10,10)	(10,10)	Comp	Comp	0.50808	$1.35 \cdot 10^{-2}$
(10,5)	(10,5)	II	II	0.495515	$2.82 \cdot 10^{-2}$	0.196772	$1.13 \cdot 10^{-2}$	0.80204	$1.27 \cdot 10^{-2}$
		III	III	0.5045	$1.52 \cdot 10^{-2}$	0.18978	$6.19 \cdot 10^{-3}$	0.81112	$6.29 \cdot 10^{-3}$
		IV	IV	0.49617	$1.66 \cdot 10^{-2}$	0.19043	$5.48 \cdot 10^{-3}$	0.810735	$6.38 \cdot 10^{-3}$
(20,20)	(10,10)	Comp	Comp	0.49997	$8.62 \cdot 10^{-3}$	0.20129	$4.41 \cdot 10^{-3}$	0.80175	$3.78 \cdot 10^{-3}$
(10,10)	(20,20)	Comp	Comp	0.50002	$1.00 \cdot 10^{-2}$	0.19802	$3.78 \cdot 10^{-3}$	0.80355	$3.89 \cdot 10^{-3}$
(20,10)	(10,5)	II	II	0.50059	$2.15 \cdot 10^{-2}$	0.20457	$9.18 \cdot 10^{-3}$	0.80413	$7.97 \cdot 10^{-3}$
		III	III	0.5036	$1.15 \cdot 10^{-2}$	0.19251	$4.62 \cdot 10^{-3}$	0.80694	$4.18 \cdot 10^{-3}$
		IV	IV	0.52110	$1.21 \cdot 10^{-2}$	0.20198	$4.81 \cdot 10^{-3}$	0.81298	$4.13 \cdot 10^{-3}$
(10,5)	(20,10)	II	II	0.49943	$2.05 \cdot 10^{-2}$	0.20083	$8.82 \cdot 10^{-3}$	0.80030	$9.25 \cdot 10^{-3}$
		III	III	0.49673	$1.15 \cdot 10^{-2}$	0.19241	$4.18 \cdot 10^{-3}$	0.80489	$4.46 \cdot 10^{-3}$
		IV	IV	0.47983	$1.24 \cdot 10^{-2}$	0.18361	$4.48 \cdot 10^{-3}$	0.80132	$5.29 \cdot 10^{-3}$
(20,10)	(20,10)	II	II	0.50182	$1.29 \cdot 10^{-2}$	0.20285	$5.59 \cdot 10^{-3}$	0.79938	$5.39 \cdot 10^{-3}$
		III	III	0.49939	$7.09 \cdot 10^{-3}$	0.19574	$2.93 \cdot 10^{-3}$	0.80537	$3.09 \cdot 10^{-3}$
		IV	IV	0.50416	$6.80 \cdot 10^{-3}$	0.19113	$2.64 \cdot 10^{-3}$	0.80283	$2.91 \cdot 10^{-3}$
(20,20)	(20,20)	Comp	Comp	0.50026	$6.29 \cdot 10^{-3}$	0.20091	$2.67 \cdot 10^{-3}$	0.79727	$2.74 \cdot 10^{-3}$
(30,15)	(30,15)	II	II	0.50353	$8.53 \cdot 10^{-3}$	0.19523	$3.54 \cdot 10^{-3}$	0.79645	$3.56 \cdot 10^{-3}$
		III	III	0.49911	$4.64 \cdot 10^{-3}$	0.19582	$1.72 \cdot 10^{-3}$	0.80565	$1.88 \cdot 10^{-3}$
		IV	IV	0.50236	$4.88 \cdot 10^{-3}$	0.19517	$1.84 \cdot 10^{-3}$	0.80332	$1.89 \cdot 10^{-3}$
(30,30)	(30,30)	Comp	Comp	0.49885	$4.23 \cdot 10^{-3}$	0.20195	$1.68 \cdot 10^{-3}$	0.80127	$1.72 \cdot 10^{-3}$

**Table 5:** Bayes estimator of R,  $\hat{R}_3$  and MSE when  $\sigma_1 = 1, \sigma_2 = 2$

$(\alpha_1, \beta_1)$	$(\alpha_2, \beta_2)$	r	s	$(n_1, n_2, m_1, m_2)=(20,20,10,10)$		$(n_1, n_2, m_1, m_2)=(30,30,15,15)$	
				$\hat{R}_3$	MSE	$\hat{R}_3$	MSE
(0,0)	(0,0)	II	II	0.22119	$5.90.10^{-3}$	0.21197	$3.57.10^{-3}$
		III	III	0.21705	$3.27.10^{-3}$	0.21186	$1.97.10^{-3}$
		IV	IV	0.21461	$2.86.10^{-3}$	0.21050	$1.95.10^{-3}$
(0,1)	(0,1)	II	II	0.23210	$6.41.10^{-3}$	0.22299	$4.10.10^{-3}$
		III	III	0.22379	$3.25.10^{-3}$	0.21600	$2.02.10^{-3}$
		IV	IV	0.22554	$3.56.10^{-3}$	0.21363	$2.02.10^{-3}$
(1,1)	(1,1)	II	II	0.22805	$5.71.10^{-3}$	0.220073	$4.31.10^{-3}$
		III	III	0.21738	$2.86.10^{-3}$	0.21280	$1.92.10^{-3}$
		IV	IV	0.22286	$3.26.10^{-3}$	0.21337	$1.95.10^{-3}$
(2,1)	(2,1)	II	II	0.23215	$6.42.10^{-3}$	0.22391	$3.91.10^{-3}$
		III	III	0.22133	$3.25.10^{-3}$	0.21549	$2.12.10^{-3}$
		IV	IV	0.22160	$3.29.10^{-3}$	0.213616	$2.01.10^{-3}$
(1,2)	(1,2)	II	II	0.24156	$6.17.10^{-3}$	0.22672	$4.29.10^{-3}$
		III	III	0.228882	$3.36.10^{-3}$	0.21822	$2.12.10^{-3}$
		IV	IV	0.22876	$3.49.10^{-3}$	0.2177	$2.06.10^{-3}$
				$(n_1, n_2, m_1, m_2)=(20,20,20,20)$		$(n_1, n_2, m_1, m_2)=(30,30,30,30)$	
$(\alpha_1, \beta_1)$	$(\alpha_2, \beta_2)$	r	s	$\hat{R}_3$ MSE		$\hat{R}_3$ MSE	
(0,0)	(0,0)	Comp	Comp	0.21128	$2.80.10^{-3}$	0.20507	$1.66.10^{-3}$
(0,1)	(0,1)	Comp	Comp	0.21521	$2.78.10^{-3}$	0.21269	$2.03.10^{-3}$
(1,1)	(1,1)	Comp	Comp	0.21656	$2.86.10^{-3}$	0.20979	$1.72.10^{-3}$
(2,1)	(2,1)	Comp	Comp	0.21768	$2.84.10^{-3}$	0.20784	$1.73.10^{-3}$
(1,2)	(1,2)	Comp	Comp	0.219668	$2.84.10^{-3}$	0.21345	$1.92.10^{-3}$

## 6 Conclusions

We have presented some efficient estimators of the stress-strength parameter R using MLE and UMVUE methods. The methods are very efficient. We have found that, our estimates of R using progressive censoring schemes are very close to estimates in case of complete samples so these estimates are better to accelerate the life testing. This work gives a general estimate since the case when sample sizes equal the number of failures is a special case. The exact confidence intervals based on MLE of Rare obtained. The choice of removing schemes effect on the estimates. Numerical results are presented which exhibit the performance of the proposed methods. From the mean square error (MSE) we found that UMVUE of R is more effective than MLE and Bayes estimator of R in many cases.

## Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this article

## References

- [1] L. Rayleigh, On the resultant of a large number of vibrations of the same pitch and of arbitrary phase. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, **10**(60), 73-78 (1880).
- [2] N. Balakrishnan and R. Aggarwala, *Progressive Censoring: theory, methods, and applications*, Springer Science & Business Media, Boston: (2000).
- [3] N. Balakrishnan, Progressive censoring methodology: an appraisal, *TEST*, **16**(2), 211 – 296 (2007).
- [4] S. Kotz and M. Pensky, *The stress-strength model and its generalization: theory and applications*, World Scientific, Singapore: (2003).
- [5] Z. W. Birnbaum, *On a use of the Mann-Whitney statistic*. Proceedings of the Third Berkeley Symposium Math. Statistic Probability I, 13-17, (1956).
- [6] R. A. Johnson, *3 Stress-strength models for reliability*. Handbook of statistics, 7, 27-54, (1988).
- [7] A. M. Awad and M. K. Charraf, Estimation of  $P(Y < X)$  in the Burr case; A comparative study. *Commun. Statist. Simul.*, **15**(2), 389-403 (1986).
- [8] P. Fathipour, A. Abolhasani and H. J. Khamnei, Estimating  $R=P(Y < X)$  in the Generalized Rayleigh Distribution with Different Scale Parameters. *Applied Mathematical Sciences*, 7, 87-92 (2013).
- [9] K. E. Ahmed, M. E. Fakhry and Z. F. Jaheen, Empirical Bayes estimation of  $R=P(Y < X)$  and characterizations of Burr-type X model. *Journal of Statistical Planning and Inference*, **64**(2), 297-308 (1997).
- [10] B. Saraoğlu, I. Kinaci and D. Kundu, On estimation of  $R=P(Y < X)$  for exponential distribution under progressive Type-II censoring, *Journal of Statistics Computation and Simulation*, **82**(5), 729-744 (2012).
- [11] A. M. Abd-Elfattah, M. H. Abu-Moussa and M. M. El-Fahham, Estimation of Stress-Strength Parameter for Burr type XII distribution Based on progressive type-II Censoring, *ANGLISTICUM. Journal of the Association-Institute for English Language and American Studies*, 3, (2016).
- [12] A. M. Abd-Elfattah and R.M. Mandouh, *Estimation of  $PrY < X$  in Lomax case*, The 39th Annual Conference on Statistics, Computer Science and Operation Research, ISSR, Cairo University, Egypt, part 1, 156-166 (2004).
- [13] H. Panahi and S. Asadi. *Estimation of  $R=P[Y < X]$  for two-parameter Burr Type XII Distribution*, World Academy of Science, Engineering and Technology, **72**, 465-470 (2010).
- [14] A. Asgharzadeh, R. Valiollahi and M. Z. Raqab, Stress-strength reliability of Weibull distribution based on progressively censored samples. *Sort: Statistics and Operations Research Transactions*, **35**(2), 103-124 (2011).