

September 2014

Visible/Near infrared (VIS/NIR) spectroscopy and multivariate data analysis (MVDA) for identification and quantification of olive leaf spot (OLS) disease

Nawaf Abukhalaf

Palestine Technical University-Kadoorie, n.abukhalaf@ptuk.edu.ps

Mazen Salman

m.salman@ptuk.edu.ps

Follow this and additional works at: <https://digitalcommons.aaru.edu.jo/ptuk>



Part of the [Arts and Humanities Commons](#), [Business Commons](#), [Education Commons](#), [Engineering Commons](#), [Other Architecture Commons](#), and the [Social and Behavioral Sciences Commons](#)

Recommended Citation

Abukhalaf, Nawaf and Salman, Mazen (2014) "Visible/Near infrared (VIS/NIR) spectroscopy and multivariate data analysis (MVDA) for identification and quantification of olive leaf spot (OLS) disease," *Palestine Technical University Research Journal*: Vol. 2 : Iss. 1 , Article 1.
Available at: <https://digitalcommons.aaru.edu.jo/ptuk/vol2/iss1/1>

This Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in Palestine Technical University Research Journal by an authorized editor. The journal is hosted on [Digital Commons](#), an Elsevier platform. For more information, please contact rakan@aarj.edu.jo, marah@aarj.edu.jo, u.murad@aarj.edu.jo.

Visible/Near infrared (VIS/NIR) spectroscopy and multivariate data analysis (MVDA) for identification and quantification of olive leaf spot (OLS) disease

Nawaf Abu-Khalaf*, Mazen Salman

Technical and Applied Research Center (TARC), Palestine Technical University–Kadoorie (PTUK), P. O. Box: 7, Tulkarm, Palestine

E-mail: nawafu@hotmail.com

Received 16 January 2014, Accepted 31 January 2014, Published 9 February 2014

Abstract: Early detection of plant disease requires usually elaborating methods techniques and especially when symptoms are not visible. Olive Leaf Spot (OLS) infecting upper surface of olive leaves has a long latent infection period. In this work, VIS/NIR spectroscopy was used to determine the latent infection and severity of the pathogens. Two different classification methods were used, Partial Least Squared-Discrimination Analysis (PLS-DA) (linear method) and Support Vector Machine (SVM) (non-linear). SVM-classification was able to classify severity levels 0, 1, 2, 3, 4, and 5 with classification rates of 94, 90, 73, 79, 83 and 100%, respectively. The overall classification rate was about 86%. PLS-DA was able to classify two different severity groups (first group with severity 0, 1, 2, 3, and second group with severity 4, 5), with a classification rate greater than 95%. The results promote further researches, and the possibility of evaluation OLS in-situ using portable VIS/NIR devices.

Keywords: olives, Olive Leaf Spot (OLS), disease severity, VIS/NIR spectroscopy, Multivariate Data, Analysis (MVDA) (i.e. chemometrics), Partial Least Squared-Discrimination Analysis (PLS-DA), Support Vector Machine (SVM)-classification

Introduction

The olive tree (*Olea europaea L.*) is one of the most ancient trees cultivated in the Mediterranean regions. Olive trees play an important role in the social and economical life of this region. It is one of the oldest agricultural trees cultivated over large areas in Palestine (Qutub *et al.*, 2012). The number of olive trees is exceeding 10 million (about 67.3% of horticultural trees) and occupying more than 50% of the agricultural area (Palestinian Central Bureau of Statistics, 2012). The trees comprise one of the main sources of income, reaching its contribution to the good years to about 13% of the annual agricultural production.

The olive tree is affected by many pests and diseases (Sanei and Razavi, 2011). Olive Leaf Spot (OLS) is a foliar disease widespread in all olive growing regions of the world, and has been known in the Mediterranean areas (Obanor *et al.*, 2005). The disease, also known as peacock eye disease, is

caused by the fungus *Spillocaea oleaginea* (Cast.) Hughes (syn. *Cycloconium oleagina*) (Gonzalez-Lamothe *et al.*, 2002). Severely infected trees show defoliation, poor twig and growth. As a result of infection, yield losses may reach up to 20% (Azeri, 1993; Graniti, 1993; Rongai, 2012). Symptoms of the disease occur usually on the upper surface of the leaves, expanding and coalescing to cover a large proportion of leaf area (**Figure 1**). The symptoms form dark brown round spots (2-15 mm in diameter) which become necrotic and surrounded by concentric yellowish or pale brown haloes (Sanchez *et al.*, 1998). Leaf spot is usually more abundant on the lower parts of olive trees (Razavi and Jahany, 2009; Rongai, 2012).

Infection with *S. oleaginea* is normally associated with high humidity and winter conditions (cool and low light), where high temperatures restrict spore germination (Al-Khatib *et al.*, 2010; Obanor *et al.*, 2008a). OLS infection can occur at any time of the year, but usually during late autumn to early summer, if environmental conditions are favourable. In hot dry weather conditions, conidia remain viable

Visible/Near infrared (VIS/NIR) spectroscopy and multivariate data analysis (MVDA) for identification and quantification of olive leaf spot (OLS) disease

but inactive on infected leaves and start to germinate early in winter. Conidium production is optimal at 15°C and/or temperatures ranging from 2 to 25°C and high humidity (85%) (Obanor *et al.*, 2008a). Conidia of *S. oleagina* are dispersed by rain splash or wind-borne water droplets (Sanei and Ravazi 2011).



Figure 1: Symptoms of Olive Leaf Spot (OLS), defoliation (top) and poor twig (bottom).

In Palestine, the disease is wide common in many areas in southern and northern parts of the country including Hebron, Bethlehem, Ramallah, Nablus, Qalqilia, Tulkarm, Salfit, Jenin and Tubas districts (Abumsha *et al.*, 2013; Salman *et al.*, 2011). The Palestinian Ministry of Agriculture and several private institutions work on olive sector to provide guidance and information about improving olive crop production and protection.

Prevalence *S. oleaginea* in Palestine is found to be most prevalent in the period from late autumn to spring and of minor significance in the period from the beginning of July until the middle of November (Abumsha *et al.*, 2008).

The disease is chemically controlled by application of coppers fungicides directly after harvest (Sistani *et al.*, 2009). The most commonly used fungicides include copper hydroxide, copper oxide and copper oxychlorides, although some long-persisting preventative fungicides have also been used to control the disease (Sistani *et al.*, 2009). Chemical treatment appears to be rarely effective (Obanor *et al.*, 2008a). Moreover, using chemical fungicides leads to the appearance of resistant pathogen races to copper (Cu) (Vanneste *et al.*, 2003) as well as disturbance of the plant metabolism following Cu accumulation in the soil (Obanor *et al.*, 2008b). Application of chemical fungicides to control the disease in Palestine is however not acceptable. This is due to the negative effects of these fungicide on human and environment health (Rumpf *et al.*, 2010), as well as their effects on taste and quality of the olive oil which usually lead to low prices and low income of the yield crop.

Visible/Near infrared (VIS/NIR) spectroscopy is a mature sensor technology. It has a big success for non-invasive analysis detection of many agricultural commodities' parameters, e.g. quality, safety, stress and disease (Alander *et al.* 2013; Garrigues and de la Guardia, 2013; Levasseur-Garcia 2012; Mahlein *et al.* 2012; Rumpf *et al.* 2010; Sankaran *et al.* 2010). This sensor technology was used for some applications related to olives (De Luca *et al.*, 2011, 2012; Salguero-Chaparro *et al.*, 2013). Portables VIS/NIR spectrometers are marketable and used for different agricultural and food industries (dos Santos *et al.* 2013; Tiwari *et al.* 2013). Interpretation of VIS/NIR spectroscopy signals is carried out using multivariate data analysis (MVDA) (*i.e.* chemometrics). The latter used different modelling and classification methods for linear and non-linear models (Abu-Khalaf *et al.*, 2013; Natsheh *et al.*, 2013).

A feasibility study for using VIS/NIR spectroscopy to detect plant diseases was carried out in our biotechnology laboratory (Abu-Khalaf and Salman, 2013). In this contribution, further steps for investigating OLS disease were carried out, taking into consideration healthy leaves and the whole range of OLS disease severity. Two classification methods namely Partial Least Squared-Discrimination Analysis (PLS-DA) (linear method) and Support Vector Machine (SVM) (non-linear) were carried out. These classification methods were used in many different applications (Abu-Khalaf *et al.* 2004; Kalinowski *et al.* 2013; Luna *et al.* 2013; Peerapattana, 2013; Szymanska, 2012; Talens *et al.* 2013; Yang *et al.* 2013). The aim of this work was to

Visible/Near infrared (VIS/NIR) spectroscopy and multivariate data analysis (MVDA) for identification and quantification of olive leaf spot (OLS) disease

develop an early detection method of the latent infection with OLS disease using VIS/NIR spectroscopy.

Material and methods

Olive leaves

Leaves from different olive trees cultivar Nabali were collected randomly from orchards in Tulkarm district (32°18'25.80"N 35°07'01.00"E). The leaves were brought directly to the laboratory. Some leaves showed clear symptoms, while others were expected to be infected. Leaves were ready, without any pre-treatment, for spectra acquisition using VIS/NIR spectroscopy.

Spectroscopy

A VIS/NIR spectroscopy, a USB2000+ miniature fiber optic spectrometer (OceanOptics, USA) with Vivo light source, was used for spectra acquisition. Leaves were placed on the top of Vivo light source, and spectra were acquired in triplicates for each leaf, covering whole leaf area.

The spectroscopy has a 550-1100 nm wave length, and a resolution of 0.35 nm full width at half maximum (FWHM). Spectroscopy has 2-MHz analog-to-digital (A/D) converter, a 2048-element CCD-array detector, and a high-speed USB 2.0 port. The USB2000+ can be controlled by SpectraSuite software. Vivo system contains four tungsten halogen bulbs that can be turned on or off individually. The risk of overheating the sample is mitigated through active cooling. This protects the sample and ensures accuracy every time. The four halogen tungsten light sources make the Vivo a high-powered VIS/NIR source, which allows a shorter integration time than conventional methods (OceanOptics, USA). The integration time used in this investigation was 1 ms.

Evaluation of OLS severity

After spectra acquisition of each leaf, severity of OLS was determined using NaOH solution. Each leaf was immersed in 5% NaOH at 50°C for 2 min. Disease severity was calculated by counting the number of lesions per leaf. Severity was then graded; 1 (1 lesion), 2 (2 lesions), 3 (3–5 lesions), 4 (6–10 lesions) or 5 (greater than 11 lesions) (Abuamsha *et al.*, 2013; Salman *et al.*, 2011).

Multivariate Data Analysis (MVDA)

Classification efficiency of leaves according to their severity using VIS/NIR spectra was carried out using MDVA. Two supervised classification methods were used, namely Partial Least Squares-Discrimination Analysis (PLS-DA) (linear method) and Support Vector Machine (SVM) (non-linear method). Unscrambler software was used for analysis (version 10.3, CAMO Software AS, Oslo, Norway).

PLS-DA consists of a classical PLS regression, where VIS/NIR spectra represented X-data matrix, and the dependent variable (Y) is categorical and represented samples class membership (*i.e.* Y with values of -1 and 1, where 1 represents each sample belonging to the targeted class and -1 represents each sample belonging to the other classes). Random cross validation, with segmentations was used.

SVM method was firstly proposed by Vapnik (1995). SVM-classification algorithms try to find patterns in empirical data (*i.e.* training data) with regard to label classes. SVMs have many advantages; the main advantages are mainly referred to their generalization ability, which is achieved by using the maximum margin hyperplane for separation and the application of non-linear discriminant functions. In addition to that, SVM can handle convexity discrimination problems. This classification method used cross validation (Rumpf *et al.* 2010; Suphamitmongkol *et al.*, 2013).

In SVM-classification, VIS/NIR spectra represented X-data matrix, and leaves severity levels were used as the response factor (Y-data matrix, label classes). Unscrambler uses one column for a classifier factor. For SVM-classification, a 15-fold cross validation was performed for the training data. Polynomial degree three was used for classification. VIS/NIR spectra (X-data matrix) were normalized and auto-scaled before analysis.

Results and discussion

Severity levels were evaluated as mentioned above, and used as the classifier parameter in PLS-DA and SVM-classification methods. The number of the leaves used for building different models and their severity levels are shown in Table 1.

Visible/Near infrared (VIS/NIR) spectroscopy and multivariate data analysis (MVDA) for identification and quantification of olive leaf spot (OLS) disease

Table 1: Number of leaves with different severity levels. Zero severity means healthy leaves. Total number of leaves is 63 leaves.

	Leaves' severity levels						Total number of leaves
	0*	1	2	3	4	5	
Number of leaves	6	7	14	15	5	16	63

*: Healthy leaves

Trials were carried out using VIS, NIR and VIS/NIR spectra to get the ranges that give the best classification rates. It was found that the best classification rates' results were obtained using both VIS and NIR spectra.

Classification using PLS-DA

PLS-DA was used for classification of severity. It was found that PLS-DA could classify only leaves with the highest severity level (*i.e.* severity level 5) in a good classification rate (greater than 90%). This is might be due to the clear severe symptoms that appear on leaves' surface with severity degree 5, and the spectroscopy can sense this high severity.

Trials using PLS-DA to classify different groups of severity were carried out. Classification groups with severity 0, 1, 2 and group with severity 3, 4, 5, didn't give a good classification rate. However, it was found that classification of group consisted of all leaves with severity 0, 1, 2, 3 (*i.e.* 126 samples with triplicates) and group consisted of all leaves with severity 4 and 5 (*i.e.* 63 samples with triplicates) had a good classification rate, using three principals components (PCs). The former explained about 97 and 81% of the variance in X-data and Y-data, respectively. Random cross validation with 20 segments was used.

Table 2 and **Figure 2** show the classification rates (arranged as a confusion matrix), and scores plots of the validation set, respectively. The confusion matrix has information about the predicted and actual classifications of samples, with each row showing the instances in a predicted group (class), and each column representing the instances in an actual group (class). The sum of classification rates in each column should be 100%.

Table 2: Classification rates (%), explained in a confusion matrix, of different groups (group with severity 0, 1, 2, 3, and group with severity 4, 5) using PLS-DA, three principal components and random cross validation. Sum of each classification rate column is 100%.

Predicted groups	Actual groups	
	Group with severity 0, 1, 2, 3	Group with severity 4, 5
Group with severity 0, 1, 2, 3	95	2
Group with severity 4, 5	5	98

It can be seen from **Figure 2**, that a clear classification of different groups using three principal components (*i.e.* three factors). This classification result can be used for screening the leaves' severity, since it can predict the severity group of new testable leaves; groups with high severity (*i.e.* group with leaves' severity 4 and 5) or to with less severity (*i.e.* group with leaves' severity 0, 1, 2, 3).

Classification using SVM

SVM-classification method was able to classify disease severity using spectra data with a high accuracy and classification rates. The accuracy of training and validation sets (**Table 3**) were 87 and 71%, respectively.

A summary of the SVM-classification rates (%) are shown in a confusion matrix (**Table 4**).

The classification rates for 0, 1, 2, 3, 4, and 5 severity levels were 94, 90, 73, 79, 83 and 100%, respectively (**Figure 3**). The overall classification rate (*i.e.* average of all classification rates) is about 86%, which is relatively high and promising for predicting new severity classes.

Visible/Near infrared (VIS/NIR) spectroscopy and multivariate data analysis (MVDA) for identification and quantification of olive leaf spot (OLS) disease




Figure 2: Scores plots of validation set of samples with different severity levels (group with leaves' severity 0, 1, 2, 3, and group with leaves' severity 4, 5) using PLS-DA, three principal components (*i.e.* three factors) and random cross validation. A: scores plot for first and second principal components. B: scores plot for first and third principal components.

It can be noticed that the classification had high rates when the severities were either low (*i.e.* severity levels 0, 1) or high (severity levels 4, 5). This can be explained by that the spectroscopy sensor may detect the extreme situations, *i.e.* low and high severities. Getting high classification rate by increasing disease severity is in agreement with Rumpf *et al.* (2010). The leaves with severity levels between high and low still had good classification rates, despite that they were less than the other classification rates.

Table 3: Accuracy (%) of training and validation sets using SVM-classification.

	Training set	Validation set
Accuracy (%)	87	71

Tables 3 and 4 show that VIS/NIR spectroscopy was able to identify and quantify of OLS severity with a high accuracy; 71% for both accuracy and 86% for averaged classification rate. SVM-classification was used to classify disease severity into two groups (group with severity 0, 1, 2, 3, and group with severity 4, 5). The classification rate was greater than 99%, which is higher than the classification rate using PLS-DA. This is due to the fact the SVM-classification can handle non-linearity issues.




Figure 3: Classification rates (%) of different severity classes using SVM-classification.

Table 4: Classification rates (%) using SVM-classification of different severity class of olive leaf spot (OLS) severity explained in a confusion matrix. Sum of each classification rate column is 100%.

Predicted severity class	Actual severity class					
	0	1	2	3	4	5
0	94					
1	6	90	4			
2		5	73	19		
3		5	23	79		
4				2	83	
5					17	100

Further research is needed to investigate the possibility of identification and quantification of OLS in-situ using portable VIS/NIR spectroscopy.

Visible/Near infrared (VIS/NIR) spectroscopy and multivariate data analysis (MVDA) for identification and quantification of olive leaf spot (OLS) disease

Conclusion

VIS/NIR spectroscopy showed a high potential for sensing latent disease of olives. It could classify olives leave with different OLS' severities, with help of multivariate classification methods. Using PLS-DA (a linear classification method), with three principal components and random cross validation, a classification rate greater than 95% was achieved for different groups (group with severity 0, 1, 2, 3, and group with severity 4, 5). SVM-classification (a non-linear classification method) provided a high classification rates and high accuracy for OLS' severities. The classification rates for 0, 1, 2, 3, 4, and 5 severity levels were 94, 90, 73, 79, 83 and 100%, respectively. The overall classification rate was about 86%. This work revealed that VIS/NIR spectroscopy has a potential for evaluation of OLS under field conditions to detect infection of olive trees with OLS disease.

Acknowledgements

The authors would like to thank Palestine Technical University-Kadoorie (PTUK) for supporting this research. Special thanks are due to Miss Basima Abu Rmaileh and Eng. Mohammad Jawabreh for their technical assistance.

References

- Abuamsha R., Abueid M., Hajjeh H. and Salman M. (2013). Evaluation of the incidence and severity of olive leaf spot caused by *Spilocaea oleagina* in different olive cultivars in Palestine. *Journal of Agriculture and Environment for International Development*, 107 (2), 201–212.
- Abu-Khalaf N. and Salman M. (2013). Detecting plant diseases using visible/near infrared spectroscopy. *NIR news*. 24 (4), 12-28.
- Abu-Khalaf N., Bennedsen B., Bjorn G. (2004). Distinguishing carrot's characteristics by near infrared (NIR) reflectance and multivariate data analysis", International Commission of Agricultural Engineering (CIGR). Manuscript FP 03 012. March, 6.
- Abu-Khalaf N., Khayat S., and Natsheh B. (2013). Multivariate data analysis to identify the groundwater pollution sources in Tulkarm area / Palestine. *Science and Technology*. 3(4), 99-104.
- Alander J. A., Bochko V., Martinkauppi B., Saranwong S. and Mantere T. (2013). A review of optical nondestructive visual and near-infrared methods for food quality and safety. *International Journal of Spectroscopy*, Volume 2013, Article ID 341402, 36 pages.
- Al-Khatib M., Alhussaen K., El-Banna N. and Zyadeh M. (2010). Biological control of olive leaf spot (peacock spot disease) caused by *Cycloconium oleaginum* (*Spilocaea oleagina*), *Journal of Microbiology and Antimicrobials*. 2(6), 64-67.
- Azeri T. (1993). Research on olive leaf spot, olive-knot and *Verticillium* wilt of olive in Turkey. *EPPO Bulletin*, 23(3), 437-440.
- De Luca M., Terouzi W., Ioele G., Kzaiber F., Oussama A., Oliverio F. and Ragno G. (2011). Derivative FTIR spectroscopy for cluster analysis and classification of morocco olive oils. *Food Chemistry*, 124(3), 1113-1118.
- De Luca M., Terouzi W., Kzaiber F., Ioele G., Oussama A. and Ragno G. (2012). Classification of moroccan olive cultivars by linear discriminant analysis applied to ATR-FTIR spectra of endocarps. *International Journal of Food Science & Technology*, 47(6) 1286–1292.
- dos Santos C.A.T., Lopo M., Páscoa R.N. and Lopes J.A. (2013). A Review on the applications of portable near-infrared spectrometers in the agro-food industry. *Applied Spectroscopy*, 67(11), 1215-1233.
- Garrigues S. and de la Guardia M. (2013). Non-invasive analysis of solid samples. *TrAC Trends in Analytical Chemistry*, 43, 161-173.
- Gonzalez-Lamothe R., Segura R., Trapero A., Baldoni L., Botella M.A. and Valpuesta V. (2002). Phylogeny of the fungus *Spilocaea oleagina*, the causal agent of peacock leaf spot in olive. *FEMS Microbiology Letters*, 210(1), 149-155.
- Graniti A. (1993). Olive Scab: A Review. *EPPO Bulletin*, 23(3), 377-384.
- Kalinowski P., Wozniak L., Strzelczyk A., Jasinski P. and Jasinski G. (2013). Efficiency of linear and non-linear classifiers for gas identification from electrocatalytic gas sensor. *Metrology and Measurement Systems*. 20(3), 501-512.
- Levasseur-Garcia C. (2012). Infrared spectroscopy applied to identification and detection of microorganisms and their metabolites on cereals (corn, wheat, and barley)", in *Agricultural Sci-*

Visible/Near infrared (VIS/NIR) spectroscopy and multivariate data analysis (MVDA) for identification and quantification of olive leaf spot (OLS) disease

- ences, Ed. by Aflakpui G., Intech, New York, p. 185.
- Luna A.S., da Silva A.P., Pinho J.S., Ferré J. and Boqué R. (2013). Rapid characterization of transgenic and non-transgenic soybean oils by chemometric methods using NIR spectroscopy. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 100, 115-119.
- Mahlein A. K., Oerke E.C., Steiner U. and Dehne H.W. (2012). Recent advances in sensing plant diseases for precision crop protection. *European Journal of Plant Pathology*, 133(1), 197-209.
- Natsheh B., Abu-Khalaf N., Sayara T., Khayat S. and Salman M. (2013). Multivariate data analysis for bioremediation of contaminated soil through interactions between heavy metals microbes and plants. *Palestine Technical University Research Journal*. 1, 21-28.
- Obanor F.O., Jaspers M.V., Jones E.E. and Walter M., (2008 b). Greenhouse and field evaluation of fungicides for control of olive leaf spot in New Zealand. *Crop Prot.* 27, 1335-1342.
- Obanor F.O., Walter M., Jones E.E. and Jaspers M.V. (2005). Sources of variation in a field evaluation of the incidence and severity of olive leaf spot. *New Zealand Plant Protection*, 58, 273-277.
- Obanor F.O., Walter M., Jones E.E., Jaspers M.V., (2008 a). Effect of temperature, relative humidity, leaf wetness and leaf age on *Spilocaea oleagina* conidium germination on olive leaves. *Eur. J. Plant Pathol.*, Vol. 120: 211-222.
- PCBS, Palestinian Central Bureau of Statistics. Ramallah, Palestine, <http://www.pcbs.gov.ps> (2012).
- Peerapattana J., Shinzawa H., Otsuka K., Hattori Y. and Otsuka M. (2013). Partial least square discriminant analysis of mangosteen pericarp powder by near infrared spectroscopy", *Journal of Near Infrared Spectroscopy*, 21(3), 195-202.
- Qutub M., Ali S., Mutawea M., Abed M., Arabasi T., Pierini F. and Lodolini E. M. (2012). Characterisation of the main Palestinian olive cultivars and olive oil.
- Razavi S.E. and Jahany M. (2009). Olive scab, in olive phytopathology, Ed. by Sanei,S.J. Peike-Reihan publication, Gorgan, p. 147.
- Rongai D., Basti C. and di Marco C. (2012). A natural product for the control of olive leaf spot caused by *Fusicladium oleagineum* (Cast.) Ritschel & Braun. *Phytopathologia Mediterranea*, 51(2), 276-282.
- Rumpf T., Mahlein A.K., Steiner U., Oerke E.C., Dehne H.W. and Plumer L. (2010). Early detection and classification of plant diseases with Support Vector Machines based on hyperspectral reflectance. *Computers and Electronics in Agriculture*, 74(1), 91-99.
- Salguero-Chaparro L., Baeten V., Fernández-Piernaand J.A. and Pena-Rodriguez F. (2013). Near infrared spectroscopy (NIRS) for on-line determination of quality parameters in intact olives. *Food Chemistry*. 139 (1-4), 1121-1126.
- Salman M., Hawamda A., Amarni A.A., Rahil M., Hajjeh H., Natsheh B. and Abuamsha R. (2011). Evaluation of the incidence and severity of olive leaf spot caused by *Spilocaea oleagina* on olive trees in Palestine. *American Journal of Plant Science*, 2(3), 457-460.
- Sanchez M.E., Ruiz Davila A., Perez de Algaba A., Blanco Lopez M.A. and Trapero Casas A. (1998). Occurrence and etiology of death of young olive-trees in southern Spain. *European Journal of Plant Pathology*, 104(4), 347-357.
- Sanei S.J. and Razavi S.E. (2011). Survey of *Spilocaea oleagina*, causal agent of olive leaf spot, in North of Iran. *Journal of Yeast and Fungal Research* . 2(3), 33-38.
- Sankaran S., Mishra A., Ehsani R. and Davis C. (2010). A review of advanced techniques for detecting plant diseases. *Computers and Electronics in Agriculture*, 72(1), 1-13.
- Sistani F., Ramezanpour S.S and Nasrollanejad S. (2009). Field evaluation of different fungicides application to control olive leaf spot. *Australian Journal of Basic and Applied Sciences*, 3(4), 3341-3345.
- Suphamitmongkol W., Nie G., Liu R., Kasemsumran S. and Shi Y. (2013). An alternative approach for the classification of orange varieties based on near infrared spectroscopy. *Computer and Electronic in Agriculture*, 91, 87-93.
- Szymanska E., Saccenti E., Smilde A.K. and Westerhuis J.A. (2012). Double-check: validation of diagnostic statistics for PLS-DA models

Visible/Near infrared (VIS/NIR) spectroscopy and multivariate data analysis (MVDA) for identification and quantification of olive leaf spot (OLS) disease

- in metabolomics studies", *Metabolomics*, 8(1), 3-16.
- Talens P., Mora L., Morsy N., Barbin D. F., EIMasry G. and Sun D.W. (2013). Prediction of water and protein contents and quality classification of Spanish cooked ham using NIR hyperspectral imaging. *Journal of Food Engineering*, 117(3), 272–280.
- Tiwari G., Slaughter D.C. and Cantwell M. (2013). Nondestructive maturity determination in green tomatoes using a handheld visible and near infrared instrument", *Postharvest Biology and Technology* 86, 221–229.
- Vanneste J.L., Voyle M.D. and Zydenbos S.M. (2003). Genetic basis of copper resistance in New Zealand strains of *Pseudomonas syringae*. *New Zealand Plant Protection*, 109-112.
- Vapnik V.N. (1995). *The nature of statistical learning theory.*, Springer-Verlag, New York.
- Yang Y., Ferro M. D., Cavaco I. and Liang Y. (2013). Detection and identification of extra virgin olive oil adulteration by GC-MS combined with chemometrics. *Journal of agricultural and food chemistry*, 61(15), 3693-3702.