

2020

## On new approximations and expositions of reciprocal third power mappings

B. V. Senthil Kumar

*Department of Information Technology, Nizwa College of Technology, Nizwa - 611, Oman,*  
senthilkumar@nct.edu.om

Khalifa Al-Shaqsi

*Department of Information Technology, Nizwa College of Technology, Nizwa - 611, Oman,*  
senthilkumar@nct.edu.om

Hemen Dutta

*Department of Mathematics, Gauhati University, Guwahati - 781 014, Assam, India,*  
senthilkumar@nct.edu.om

Follow this and additional works at: <https://digitalcommons.aaru.edu.jo/isl>

---

### Recommended Citation

V. Senthil Kumar, B.; Al-Shaqsi, Khalifa; and Dutta, Hemen (2020) "On new approximations and expositions of reciprocal third power mappings," *Information Sciences Letters*: Vol. 9 : Iss. 2 , Article 5.  
Available at: <https://digitalcommons.aaru.edu.jo/isl/vol9/iss2/5>

This Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in Information Sciences Letters by an authorized editor. The journal is hosted on Digital Commons, an Elsevier platform. For more information, please contact [rakan@aarj.edu.jo](mailto:rakan@aarj.edu.jo), [marah@aarj.edu.jo](mailto:marah@aarj.edu.jo), [u.murad@aarj.edu.jo](mailto:u.murad@aarj.edu.jo).

# On new approximations and expositions of reciprocal third power mappings

B. V. Senthil Kumar<sup>1,\*</sup>, Khalifa Al-Shaqsi<sup>1</sup> and Hemen Dutta<sup>2</sup>

<sup>1</sup>Department of Information Technology, Nizwa College of Technology, Nizwa - 611, Oman

<sup>2</sup>Department of Mathematics, Gauhati University, Guwahati - 781 014, Assam, India

Received: 17 Feb. 2020, Revised: 17 April 2020, Accepted: 21 April 2020

Published online: 1 May 2020

**Abstract:** The intention of this work is to deal with new form of reciprocal third power functional equations for their solutions. The Ulam stabilities of these equations are determined in the setting of non-Archimedean fields. A proper instance is demonstrated to show the invalidity of stability result for a very critical case. The interpretation of the equations dealt in this study is associated with a significant hypothesis in electromagnetic theory and the relation of stiffness (or deflection) and length of diving board (or cantilever beam).

**Keywords:** Reciprocal functional equation, cubic functional equation, Generalized Hyers-Ulam stability, non-Archimedean field.

## 1 Introduction

The inception and the evolution of Ulam stability results of various equations in different versions are accessible in [1, 2, 3, 4, 5, 6]. For the past four decades, there are many interesting, new, motivating and important and remarkable results pertinent to various forms of functional, reciprocal or rational type or multiplicative inverse, difference, differential and integral equations, one can refer to [7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20]. Quite recently, the following reciprocal third power functional equation,

$$m_c(\lambda_1 + \lambda_2) = \frac{m_c(\lambda_1)m_c(\lambda_2)}{\left[m_c(\lambda_1)^{\frac{1}{3}} + m_c(\lambda_2)^{\frac{1}{3}}\right]^3} \quad (1)$$

is dealt in [21] to investigate its fuzzy stabilities and its application using inverse cubic law. It is also prove that the reciprocal third power mapping  $m_c(\lambda) = \frac{k}{\lambda^3}$  is a solution of (1). Inspired by the significant results available in the literature, we propose a new reciprocal third power difference functional equation:

$$m_c\left(\frac{1}{p}\sum_{k=1}^p\lambda_k\right) - m_c\left(\sum_{k=1}^p\lambda_k\right) = \frac{(p^3 - 1)\prod_{k=1}^p m_c(\lambda_k)}{\left[\sum_{k=1}^p m_c(\lambda_k)^{\frac{1}{3}}\right]^3} \quad (2)$$

and a reciprocal third power adjoint functional equation:

$$m_c\left(\frac{1}{p}\sum_{k=1}^p\lambda_k\right) + m_c\left(\sum_{k=1}^p\lambda_k\right) = \frac{(p^3 + 1)\prod_{k=1}^p m_c(\lambda_k)}{\left[\sum_{k=1}^p m_c(\lambda_k)^{\frac{1}{3}}\right]^3} \quad (3)$$

It is not hard to verify that the reciprocal third power mapping  $m_c(\lambda) = \frac{1}{\lambda^3}$  is a solution of equations (2) and (3). We illustrate the application of the above equations, for  $p = 2$  with  $(u_1, u_2)$  replaced with  $(u, v)$  in both equations. Hence, we can reduce them in the following forms for two variables:

$$m_c\left(\frac{\lambda_1 + \lambda_2}{2}\right) - m_c(\lambda_1 + \lambda_2) = \frac{7m_c(\lambda_1)m_c(\lambda_2)}{\left[m_c(\lambda_1)^{\frac{1}{3}} + m_c(\lambda_2)^{\frac{1}{3}}\right]^3} \quad (4)$$

and

$$m_c\left(\frac{\lambda_1 + \lambda_2}{2}\right) + m_c(\lambda_1 + \lambda_2) = \frac{9m_c(\lambda_1)m_c(\lambda_2)}{\left[m_c(\lambda_1)^{\frac{1}{3}} + m_c(\lambda_2)^{\frac{1}{3}}\right]^3} \quad (5)$$

respectively. We solve Ulam stability problems for the above equations (4) and (5) in the setting of non-Archimedean fields. We present a suitable counter-example for the invalidity of stability result in

\* Corresponding author e-mail: [senthilkumar@nct.edu.om](mailto:senthilkumar@nct.edu.om)

case of singularity. Then, we elucidate their implications in electromagnetism and in the property of diving board.

In order to prove our main results, we furnish below the fundamental definition of non-Archimedean field.

**Definition 1.** Suppose  $\mathbb{G}$  is a field provided with a valuation  $|\cdot| : \mathbb{G} \rightarrow [0, \infty)$ . Then  $\mathbb{G}$  is said to be a non-Archimedean field if the ensuing conditions hold:  $|k| = 0$  if and only if  $k = 0$ ,  $|k\ell| = |k||\ell|$  and  $|k + \ell| \leq \sup\{|k|, |\ell|\}$  for all  $k, \ell \in \mathbb{G}$ .

In the entire study, we assume that  $\mathbb{G}$  and  $\mathbb{H}$  is a non-Archimedean field and a complete non-Archimedean field, respectively. From now on, we use the symbol  $\mathbb{G}^*$  which excludes 0 from  $\mathbb{G}$ . For the intention of proving the major results in an easy manner, let us define the difference operator  $\Delta m_c : \mathbb{G}^* \times \mathbb{G}^* \rightarrow \mathbb{H}$  as follows:

$$\Delta m_c(\lambda_1, \lambda_2) = m_c\left(\frac{\lambda_1 + \lambda_2}{2}\right) - m_c(\lambda_1 + \lambda_2) - \frac{7m_c(\lambda_1)m_c(\lambda_2)}{\left[m_c(\lambda_1)^{\frac{1}{3}} + m_c(\lambda_2)^{\frac{1}{3}}\right]^3}$$

for all  $\lambda_1, \lambda_2 \in \mathbb{G}^*$ .

## 2 Similarity of equations (4) and (5)

In this section, let us prove that the equations (4) and (5) have reciprocal third power mapping as their solution. In the following result, we show that they are analogous to each other. In the following outcome, we assume that  $\lambda_1, \lambda_2, \lambda \in \mathbb{R}^*$ .

**Theorem 1.** Let  $m_c : \mathbb{R}^* \rightarrow \mathbb{R}$  be a mapping. Then the following assertions are identical to each other.

- (a)  $m_c$  is a solution of (1).
- (b)  $m_c$  is a solution of (4).
- (c)  $m_c$  is a solution of (5).

Thus, a reciprocal third power mapping is a solution of equations (4) and (5).

*Proof.* Let us first assume that  $m_c$  is a solution of (1). Now, switching  $(\lambda_1, \lambda_2)$  to  $(\frac{\lambda}{2}, \frac{\lambda}{2})$  in (1) and then multiplying by 8, one attains that

$$m_c\left(\frac{\lambda}{2}\right) = 8m_c(\lambda). \tag{6}$$

Then, reinstating  $(\lambda_1, \lambda_2)$  by  $(\frac{\lambda_1}{2}, \frac{\lambda_2}{2})$  in (1) and using the result of (6) in the resultant, one acquires that

$$m_c\left(\frac{\lambda_1 + \lambda_2}{2}\right) = \frac{8m_c(\lambda_1)m_c(\lambda_2)}{\left[m_c(\lambda_1)^{\frac{1}{3}} + m_c(\lambda_2)^{\frac{1}{3}}\right]^3}. \tag{7}$$

It is easy to achieve equation (4) by taking the difference between (7) and (1). Next, let us assume that  $m_c$  is a solution of (5). If  $(\lambda_1, \lambda_2)$  is changed as  $(\frac{\lambda}{2}, \frac{\lambda}{2})$  in (5) and on further simplification, one finds that

$$m_c\left(\frac{\lambda}{2}\right) = 8m_c(\lambda). \tag{8}$$

Employing the result of (8) in (4) and then simplifying further, one obtains that

$$m_c(\lambda_1 + \lambda_2) = \frac{m_c(\lambda_1)m_c(\lambda_2)}{\left[m_c(\lambda_1)^{\frac{1}{3}} + m_c(\lambda_2)^{\frac{1}{3}}\right]^3}. \tag{9}$$

Now, replacing  $(\lambda_1, \lambda_2)$  by  $(\frac{\lambda_1}{2}, \frac{\lambda_2}{2})$  in (9) and then utilizing (8), one gets that

$$m_c\left(\frac{\lambda_1 + \lambda_2}{2}\right) = \frac{8m_c(\lambda_1)m_c(\lambda_2)}{\left[m_c(\lambda_1)^{\frac{1}{3}} + m_c(\lambda_2)^{\frac{1}{3}}\right]^3}. \tag{10}$$

Adding up equations (10) and (9), one arrives at (5). Finally, let us assume that  $m_c$  is a solution of (5). Using similar reasoning applied in the aforementioned steps, if  $(\lambda_1, \lambda_2)$  is considered as  $(\frac{\lambda}{2}, \frac{\lambda}{2})$  in (5) and simplified further, one has

$$m_c\left(\frac{\lambda}{2}\right) = 8m_c(\lambda). \tag{11}$$

Using the outcome of (11) in (5), it produces (1). Therefore,  $m_c$  is a reciprocal third power mapping.

## 3 Approximation of reciprocal third power mapping

In the upcoming results, we determine the validity of various fundamental stabilities of equations (4) and (5) associated with Ulam, Hyers, T. Rassias, J. Rassias and Gavruta.

**Theorem 2.** Consider a fixed number  $\beta \neq \pm 1$ . Suppose  $m_c : \mathbb{G}^* \rightarrow \mathbb{H}$  is a mapping satisfies

$$|\Delta m_c(\lambda_1, \lambda_2)| \leq \xi(\lambda_1, \lambda_2) \tag{12}$$

where  $\xi : \mathbb{G}^* \times \mathbb{G}^* \rightarrow [0, \infty)$  is a function with the condition that

$$\lim_{m \rightarrow \infty} \left|\frac{1}{8}\right|^{\beta m} \xi\left(\frac{\lambda_1}{2^{\beta m + \frac{\beta+1}{2}}}, \frac{\lambda_2}{2^{\beta m + \frac{\beta+1}{2}}}\right) = 0 \tag{13}$$

for all  $\lambda_1, \lambda_2 \in \mathbb{G}^*$ . Then, there exists a unique reciprocal third power mapping  $M_c : \mathbb{G}^* \rightarrow \mathbb{H}$  which satisfies (4) with the result that

$$|m_c(u) - M_c(u)| \leq \sup \left\{ \left|\frac{1}{8}\right|^{p\beta + \frac{\beta-1}{2}} \xi\left(\frac{\lambda}{2^{p\beta + \frac{\beta+1}{2}}}, \frac{\lambda}{2^{p\beta + \frac{\beta+1}{2}}}\right) \right\} : p \in \mathbb{N} \cup \{0\} \tag{14}$$

for all  $\lambda \in \mathbb{G}^*$ .

*Proof.* Firstly, we construct a reciprocal third power mapping  $M_c$  satisfying (4). For this, let us replace  $(\lambda_1, \lambda_2)$  by  $(\frac{\lambda}{2}, \frac{\lambda}{2})$  in (12) to obtain

$$\left| m_c(\lambda) - \frac{1}{2^\beta} m_c\left(\frac{\lambda}{8^\beta}\right) \right| \leq |8|^{\frac{|\beta-1|}{2}} \xi\left(\frac{\lambda}{2^{\frac{\beta+1}{2}}}, \frac{\lambda}{2^{\frac{\beta+1}{2}}}\right) \tag{15}$$

for all  $\lambda \in \mathbb{G}^*$ . Now, again replace  $\lambda$  with  $\frac{\lambda}{2^{\beta m}}$  in (15) and multiply by  $\left|\frac{1}{8}\right|^{\beta m}$  in the resultant to acquire

$$\left| \frac{1}{8^{\beta m}} m_c\left(\frac{\lambda}{2^{\beta m}}\right) - \frac{1}{8^{(m+1)\beta}} m_c\left(\frac{\lambda}{2^{(m+1)\beta}}\right) \right| \leq \left|\frac{1}{8}\right|^{\beta m + \frac{\beta-1}{2}} \xi\left(\frac{\lambda}{2^{\beta m + \frac{\beta+1}{2}}}, \frac{\lambda}{2^{\beta m + \frac{\beta+1}{2}}}\right) \tag{16}$$

for all  $\lambda \in \mathbb{G}^*$ . Application of (13) and (16) produces that the sequence  $\left\{\frac{1}{8^{\beta m}} m_c\left(\frac{\lambda}{2^{\beta m}}\right)\right\}$  is Cauchy. In view of the fact that  $\mathbb{H}$  is complete, this sequence converges to a mapping  $M_c : \mathbb{G}^* \rightarrow \mathbb{H}$  given by

$$M_c(\lambda) = \lim_{m \rightarrow \infty} \frac{1}{8^{\beta m}} m_c\left(\frac{\lambda}{2^{\beta m}}\right). \tag{17}$$

Also, for each  $\lambda \in \mathbb{G}^*$  and for all integers  $k > 0$ , we have

$$\begin{aligned} \left| \frac{1}{8^{\beta m}} m_c\left(\frac{\lambda}{2^{\beta m}}\right) - m_c(\lambda) \right| &= \left| \sum_{p=0}^{m-1} \left\{ \frac{1}{8^{(p+1)\beta}} m_c\left(\frac{\lambda}{2^{(p+1)\beta}}\right) - \frac{1}{8^{p\beta}} m_c\left(\frac{\lambda}{2^{p\beta}}\right) \right\} \right| \\ &\leq \sup \left\{ \left| \frac{1}{8^{(p+1)\beta}} m_c\left(\frac{\lambda}{2^{(p+1)\beta}}\right) - \frac{1}{8^{p\beta}} m_c\left(\frac{\lambda}{2^{p\beta}}\right) \right| : 0 \leq p < m \right\} \\ &\leq \sup \left\{ \left| \frac{1}{8} \right|^{p\beta + \frac{\beta-1}{2}} \xi\left(\frac{\lambda}{2^{p\beta + \frac{\beta+1}{2}}}, \frac{\lambda}{2^{p\beta + \frac{\beta+1}{2}}}\right) : 0 \leq p < m \right\}. \end{aligned} \tag{18}$$

Now, employing (17) and allowing  $m \rightarrow \infty$  in the inequality (18), we observe that the inequality (14) exists. Using (13), (12) and (17), for all  $\lambda_1, \lambda_2 \in \mathbb{G}^*$  we have

$$|\Delta m_c(\lambda_1, \lambda_2)| = \lim_{m \rightarrow \infty} \left| \frac{1}{8} \right|^{\beta m} \left| \Delta m_c\left(\frac{\lambda_1}{2^{\beta m}}, \frac{\lambda_2}{2^{\beta m}}\right) \right| \leq \lim_{m \rightarrow \infty} \left| \frac{1}{8} \right|^{\beta m} \xi\left(\frac{\lambda_1}{2^{\beta m}}, \frac{\lambda_2}{2^{\beta m}}\right) = 0$$

which implies that  $M_c$  satisfies equation (4) and therefore it is a reciprocal third power mapping. Next, is to prove the uniqueness assertion of  $M_c$ . For this, let us presume that  $M'_c : \mathbb{G}^* \rightarrow \mathbb{H}$  be another reciprocal third power mapping satisfying the approximation (14). Then, we have

$$\begin{aligned} |M_c(\lambda) - M'_c(\lambda)| &= \lim_{k \rightarrow \infty} \left| \frac{1}{2} \right|^{\beta k} \left| M_c\left(\frac{\lambda}{2^{\beta k}}\right) - M'_c\left(\frac{\lambda}{2^{\beta k}}\right) \right| \\ &\leq \lim_{k \rightarrow \infty} \left| \frac{1}{2} \right|^{\beta k} \sup \left\{ \left| M_c\left(\frac{\lambda}{2^{\beta k}}\right) - m_c\left(\frac{\lambda}{2^{\beta k}}\right) \right|, \left| m_c\left(\frac{\lambda}{2^{\beta k}}\right) - M'_c\left(\frac{\lambda}{2^{\beta k}}\right) \right| \right\} \\ &\leq \lim_{k \rightarrow \infty} \lim_{m \rightarrow \infty} \sup \left\{ \sup \left\{ \left| \frac{1}{8} \right|^{(p+k)\beta + \frac{\beta-1}{2}} \xi\left(\frac{\lambda}{2^{(p+k)\beta + \frac{\beta+1}{2}}}, \frac{\lambda}{2^{(p+k)\beta + \frac{\beta+1}{2}}}\right) : k \leq p \leq m+k \right\} \right\} \\ &= 0 \end{aligned}$$

for all  $\lambda \in \mathbb{G}^*$ , which shows that  $M_c$  is unique. This completes the proof.

The ensuing corollaries are pertinent to other stabilities of equation (4) involving a positive constant, sum of powers of norms, product of different powers of norms and mixed product-sum of powers of norms as upper bounds and their proofs can be accomplished by the application of Theorem 2. Hence, we furnish only the

statements. In the following outcomes, let us assume that  $m_c : \mathbb{G}^* \rightarrow \mathbb{H}$  is a mapping.



**Corollary 1.** Let  $\mu > 0$  be a constant. If the mapping  $m_c$  satisfies  $|\Delta m_c(\lambda_1, \lambda_2)| \leq \mu$  for all  $\lambda_1, \lambda_2 \in \mathbb{G}^*$ , then a unique reciprocal third power mapping  $M_c : \mathbb{G}^* \rightarrow \mathbb{H}$  exists and satisfies (4) with  $|m_c(\lambda) - M_c(\lambda)| \leq \mu$  for all  $\lambda \in \mathbb{G}^*$ .

**Corollary 2.** Let  $\mu \geq 0$  and  $q \neq -3$ , be fixed constants. If the mapping  $m_c$  satisfies  $|\Delta m_c(\lambda_1, \lambda_2)| \leq \mu (|\lambda_1|^q + |\lambda_2|^q)$  for all  $\lambda_1, \lambda_2 \in \mathbb{G}^*$ , then there exists a unique reciprocal third power mapping  $M_c : \mathbb{G}^* \rightarrow \mathbb{H}$  satisfying (4) and

$$|m_c(\lambda) - M_c(\lambda)| \leq \begin{cases} \frac{|\mu|}{|2|^q} |\lambda|^q, & q > -3 \\ |\mu| 2^3 |\lambda|^q, & q < -3 \end{cases}$$

for all  $\lambda \in \mathbb{G}^*$ .

**Corollary 3.** Let  $a, b : q = a + b \neq -3$  and  $\mu \geq 0$  be real numbers. Suppose the mapping  $m_c$  satisfies  $|\Delta m_c(\lambda_1, \lambda_2)| \leq \mu |\lambda_1|^a |\lambda_2|^b$  for all  $\lambda_1, \lambda_2 \in \mathbb{G}^*$ . Then, a unique reciprocal third power mapping  $M_c : \mathbb{G}^* \rightarrow \mathbb{H}$  satisfying (4) exists with the result that

$$|m_c(\lambda) - M_c(\lambda)| \leq \begin{cases} \frac{\mu}{|2|^q} |\lambda|^q, & q > -3 \\ \mu |2|^3 |\lambda|^q, & q < -3 \end{cases}$$

for all  $\lambda \in \mathbb{G}^*$ .

**Corollary 4.** Let  $\mu \geq 0$  and  $q \neq -3$  be real numbers. Suppose the mapping  $m_c$  satisfies

$$|\Delta m_c(\lambda_1, \lambda_2)| \leq \mu \left( |\lambda_1|^{\frac{q}{2}} |\lambda_2|^{\frac{q}{2}} + (|\lambda_1|^q + |\lambda_2|^q) \right)$$

for all  $\lambda_1, \lambda_2 \in \mathbb{G}^*$ . Then, there exists a unique reciprocal third power mapping  $M_c : \mathbb{G}^* \rightarrow \mathbb{H}$  satisfying (4) with the result that

$$|m_c(\lambda) - M_c(\lambda)| \leq \begin{cases} \frac{|\mu|}{|2|^q} |\lambda|^q, & q > -3 \\ |\mu| 3 |2|^3 |\lambda|^q, & q < -3 \end{cases}$$

for all  $\lambda \in \mathbb{G}^*$ .

Motivated from the excellent counter-example presented in [22], we prove the failure of stability result of equation (4) for a very singular case  $q = -3$  in Corollary 2 in the setting of non-zero real numbers. Let us consider the following function:

$$\nu(\lambda) = \begin{cases} \frac{\beta}{\lambda^3}, & \text{for } \lambda \in (1, \infty) \\ \beta, & \text{otherwise} \end{cases} \quad (19)$$

where  $\nu : \mathbb{R}^* \rightarrow \mathbb{R}$ . Let  $m_c : \mathbb{R}^* \rightarrow \mathbb{R}$  be a mapping defined by

$$m_c(\lambda) = \sum_{k=0}^{\infty} 2^{-3k} \nu(2^{-k} \lambda) \quad (20)$$

for all  $\lambda \in \mathbb{R}^*$ . In the ensuing theorem, the mapping  $m_c$  becomes an example to prove that equation (4) is not stable for  $q = -3$  in Corollary 2.

**Theorem 3.** If the mapping  $m_c : \mathbb{R}^* \rightarrow \mathbb{R}$  defined in (19) satisfies the inequality

$$|\Delta m_c(\lambda_1, \lambda_2)| \leq \frac{184\beta}{7} \left( |\lambda_1|^{-3} + |\lambda_2|^{-3} \right) \quad (21)$$

for all  $\lambda_1, \lambda_2 \in \mathbb{R}^*$ , then a reciprocal third power mapping  $M_c : \mathbb{R}^* \rightarrow \mathbb{R}$  and a constant  $C > 0$  do not exist such that

$$|m_c(\lambda) - M_c(\lambda)| \leq C |\lambda|^{-3} \quad (22)$$

for all  $\lambda \in \mathbb{R}^*$ .

*Proof.* Firstly, let us show that  $m_c$  satisfies (21). From the definition of  $m_c$ , we have  $|m_c(\lambda)| = \left| \sum_{k=0}^{\infty} 2^{-3k} \nu(2^{-k} \lambda) \right| \leq \sum_{k=0}^{\infty} \frac{\beta}{2^{3k}} = \frac{8\beta}{7}$ , which implies that the real number  $\frac{8\beta}{7}$  is an upper bound for the mapping  $m_c$ . When  $|\lambda_1|^{-3} + |\lambda_2|^{-3} \geq 1$ , then  $|\Delta m_c(\lambda_1, \lambda_2)| < \frac{184\beta}{7}$ . Now, when we suppose that  $0 < |\lambda_1|^{-3} + |\lambda_2|^{-3} < 1$ , then there exists a positive integer  $j$  such that

$$\frac{1}{2^{3(j+1)}} \leq |\lambda_1|^{-3} + |\lambda_2|^{-3} < \frac{1}{2^{3j}}. \quad (23)$$

The above relation (23) yields  $2^{3j} (|\lambda_1|^{-3} + |\lambda_2|^{-3}) < 1$ , and further produces  $2^{3j} \lambda_1^{-3} < 1, 2^{3j} \lambda_2^{-3} < 1$ . Hence, we have  $\frac{\lambda_1}{2^j} > 1, \frac{\lambda_2}{2^j} > 1$ . From the last two inequalities, we find that  $\frac{\lambda_1}{2^{j-1}} > 2 > 1, \frac{\lambda_2}{2^{j-1}} > 2 > 1$  and as a result, we have,  $\frac{1}{2^{j-1}}(\lambda_1) > 1, \frac{1}{2^{j-1}}(\lambda_2) > 1, \frac{1}{2^{j-1}}(\lambda_1 + \lambda_2) > 1, \frac{1}{2^{j-1}} \left( \frac{\lambda_1 + \lambda_2}{2} \right) > 1$ . Thus, for every  $k = 0, 1, 2, \dots, j - 1$ , we obtain,

$\frac{1}{2^k}(\lambda_1) > 1, \frac{1}{2^k}(\lambda_2) > 1, \frac{1}{2^k}(\lambda_1 + \lambda_2) > 1, \frac{1}{2^k} \left( \frac{\lambda_1 + \lambda_2}{2} \right) > 1$

and  $\Delta \nu(2^{-k} \lambda_1, 2^{-k} \lambda_2) = 0$  for  $k = 0, 1, 2, \dots, j - 1$ . Using (19) and the definition of  $m_c$ , we obtain,

$$|\Delta m_c(\lambda_1, \lambda_2)| \leq \sum_{k=j}^{\infty} \frac{\beta}{2^{3k}} + \sum_{k=j}^{\infty} \frac{\beta}{2^{3k}} + \sum_{k=j}^{\infty} \frac{7\beta}{8 \cdot 2^{3k}} \leq \frac{184\beta}{8} \sum_{k=j}^{\infty} \frac{1}{2^{3k}} \leq \frac{184\beta}{7} \left( |\lambda_1|^{-3} + |\lambda_2|^{-3} \right)$$

for all  $\lambda_1, \lambda_2 \in \mathbb{R}^*$ . Hence, the inequality (21) holds. Now, we assert that the equation (4) is not stable for  $q = -3$  in Corollary 2. For this, let us consider a reciprocal third power mapping  $m_c : \mathbb{R}^* \rightarrow \mathbb{R}$  exists and satisfies (22). Then, we have

$$|m_c(\lambda)| \leq (C + 1) |\lambda|^{-3}. \quad (24)$$

But it is possible to choose a positive integer  $m$  with  $m\beta > C + 1$ . If  $\lambda \in (1, 2^{m-1})$  then  $2^{-k} \lambda \in (1, \infty)$  for all  $k = 0, 1, 2, \dots, m - 1$  and thus,

$$|m_c(\lambda)| = \sum_{k=0}^{\infty} \frac{\nu(2^{-k} \lambda)}{2^{3k}} \geq \sum_{k=0}^{m-1} \frac{2^{3k} \beta}{2^{3k}} = \frac{m\beta}{\lambda^3} > (C + 1) \lambda^{-3}$$

which contradicts (24) and hence this concludes that the equation (4) is not stable for  $q = -3$  in Corollary 2.

*Remark.* The stability results concerning equation (5) and an illustration of a counter-example for singular case to show non-stability of equation (5) can be achieved similar to the results of equation (4).

#### 4 Elucidation of equations (4) and (5) in real time

In this section, we interpret the insinuation of equations (4) and (5) in real time such as electromagnetic theory and construction of spring or diving board in a swimming pool.

- We associate the well-known magnetostatic reciprocal third power law with equations (4) and (5). This law

$$\begin{aligned}
 M_{\frac{\lambda_1+\lambda_2}{2}} - M_{\lambda_1+\lambda_2} &= \frac{7K}{(\lambda_1 + \lambda_2)^3} \\
 &= \frac{7K}{\lambda_1^3 + 3\lambda_1^2\lambda_2 + 3\lambda_1\lambda_2^2 + \lambda_2^3} \\
 &= \frac{\frac{7K}{\lambda_1^3\lambda_2^3}}{\left(\frac{1}{\lambda_1^3} + \frac{3}{\lambda_1^2\lambda_2} + \frac{3}{\lambda_1\lambda_2^2} + \frac{1}{\lambda_2^3}\right)} \\
 &= \frac{7M_{\lambda_1}M_{\lambda_2}}{M_{\lambda_1} + 3M_{\lambda_1}^{\frac{2}{3}}M_{\lambda_2}^{\frac{1}{3}} + 3M_{\lambda_1}^{\frac{1}{3}}M_{\lambda_2}^{\frac{2}{3}} + M_{\lambda_2}} = \frac{7M_{\lambda_1}M_{\lambda_2}}{\left[M_{\lambda_1}^{\frac{1}{3}} + M_{\lambda_2}^{\frac{1}{3}}\right]^3}. \quad (25)
 \end{aligned}$$

The aforementioned equation (25) connecting the field strength with different circumstances can be interpreted through equation (4). Analogous to this association, equation (5) can be expounded with the sum of field strengths  $M_{\frac{\lambda_1+\lambda_2}{2}}$  and  $M_{\lambda_1+\lambda_2}$ .

- The stiffness or deflection of a diving board or cantilever beam is proportional to the cube of its thickness and inversely proportional to the cube of its length. Thus, the solution of equations (4) and (5) can be related with the stiffness or deflection as it is a function of reciprocal third power of length. Suppose if  $S$  is the stiffness of a diving board with length  $\ell$ , then we have  $S = \frac{K}{\ell^3}$ , where  $K$  is constant of proportionality. Equation (4) interprets that it is a relation between difference of the stiffness of diving boards of lengths  $\frac{\lambda_1+\lambda_2}{2}$  and  $\lambda_1 + \lambda_2$  and the stiffness of diving boards of lengths  $\lambda_1$  and  $\lambda_2$ . Similarly, we can infer the meaning of equation (5) with the sum of stiffness of diving boards in different situation.

#### 5 Conclusion

In this study, we have introduced new reciprocal third power difference and adjoint functional equations (4) and (5). We have also proved the stability results pertinent to the results of Ulam, Hyers, T. Rassias, J. Rassias and Gavruta. We have illustrated that the equation (4) is unstable for a singular case by means of a counter-example. The relevance of equations (4) and (5) is interpreted through magnetic field strength in

states that the magnetic field strength  $M_{\lambda_1}$  due to a dipole from a point at a distance  $\lambda_1$  is proportional to the reciprocal third power of  $\lambda_1$ . Then,  $M_{\lambda_1} = \frac{K}{\lambda_1^3}$ , where  $K$  is a constant of proportionality. Similarly, suppose  $\lambda_2$  is the distance between them, then,  $M_{\lambda_2} = \frac{K}{\lambda_2^3}$ . Now, suppose a dipole is at a distance  $(\lambda_1 + \lambda_2)$  from a point, then,  $M_{\lambda_1+\lambda_2} = \frac{K}{(\lambda_1+\lambda_2)^3}$ . Suppose the distance  $\lambda_1 + \lambda_2$  is halved, then the magnetic field strength is  $M_{\frac{\lambda_1+\lambda_2}{2}} = \frac{8K}{(\lambda_1+\lambda_2)^3}$ . Now,

electromagnetic theory and relation between stiffness and length of a diving board in a swimming pool.

#### Acknowledgment

The first two authors are supported by the Research Council, Oman (Under Project proposal ID: BFP/RGP/CBS/18/099).

#### References

- [1] P. Găvruta, A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings, *J. Math. Anal. Appl.* **184** (1994) 431–436.
- [2] D. H. Hyers, On the stability of the linear functional equation, *Proc. Nat. Acad. Sci.*, **27** (1941) 222–224.
- [3] J. M. Rassias, On approximately of approximately linear mappings by linear mappings, *J. Funct. Anal.*, **46** (1982) 126–130.
- [4] T. M. Rassias, On the stability of the linear mapping in Banach spaces, *Proc. Amer. Math. Soc.* **72** (1978) 297–300.
- [5] K. Ravi, M. Arunkumar and J. M. Rassias, On the Ulam stability for the orthogonally general Euler-Lagrange type functional equation, *International Journal of Mathematical Sciences*, **3** (2008) 36–47.
- [6] S. M. Ulam, *Problems in Modern Mathematics*, Chapter VI, Wiley-Interscience, New York, 1964.
- [7] T. Aoki, On the stability of the linear transformation in Banach spaces, *J. Math. Soc. Japan*, **2** (1950) 64–66.
- [8] A. Bodaghi and B. V. Senthil Kumar, Estimation of inexact reciprocal-quintic and reciprocal-sextic functional equations, *Mathematica*, **49** (2017) 3–14.

- [9] A. Ebadian, S. Zolfaghari, S. Ostadbashi and C. Park, Approximation on the reciprocal functional equation in several variables in matrix non-Archimedean random normed spaces, *Adv. Diff. Equ.* **314** (2015) 2015. DOI10.1186/s13662-015-0656-7.
- [10] D. Kang and H. B. Kim, The generalized Hyers-Ulam-Rassias stability of cubic functional equations generalized by an automorphism on groups, *Global J. Pure Appl. Math.* **13** (2017) 4695–4707.
- [11] S. Owyed, M. Abdou, A.-H. Abdel-Aty, W. Alharbi and R. Nekhili, Numerical and approximate solutions for coupled time fractional nonlinear evolutions equations via reduced differential transform method, *Chaos, Solitons & Fractals* **131** (2020) 109474.
- [12] S. O. Kim, B. V. Senthil Kumar and A. Bodaghi, Stability and non-stability of the reciprocal-cubic and reciprocal-quartic functional equations in non-Archimedean fields, *Adv. Difference Equ.*, **77** (2017) 12 pages.
- [13] S. S. Kim, J. M. Rassias, N. Hussain and Y. J. Cho, Generalized Hyers-Ulam stability of general cubic functional equation in random normed spaces, *Filomat*, **30** (2016) 89–98.
- [14] A. K. Mirmostafae, Non-Archimedean stability of quadratic equations, *Fixed Point Theory*, **11** (2010) 67–75.
- [15] K. Ravi, J. M. Rassias and B. V. Senthil Kumar, Ulam stability of reciprocal difference and adjoint functional equations, *Aust. J. Math. Anal. Appl.* **8** (2011) 1–18.
- [16] K. Ravi and B. V. Senthil Kumar, Ulam-Gavruta-Rassias stability of Rassias reciprocal functional equation, *Global J. Appl. Math. Sci.*, **3** (2010) 57–79.
- [17] K. Ravi, E. Thandapani and B. V. Senthil Kumar, Stability of reciprocal type functional equations, *Pan American Math. J.*, **21** (2011) 59–70.
- [18] B. V. Senthil Kumar and H. Dutta, Non-Archimedean stability of a generalized reciprocal-quadratic functional equation in several variables by direct and fixed point methods, *Filomat*, **32** (2018) 3199–3209.
- [19] B. V. Senthil Kumar and H. Dutta, Fuzzy stability of a rational functional equation and its relevance to system design, *Int. J. General Syst.* **48** (2019) 157–169.
- [20] B. V. Senthil Kumar and H. Dutta, Approximation of multiplicative inverse undecic and duodecic functional equations, *Math. Meth. Appl. Sci.* **42** (2019) 1073–1081.
- [21] B. V. Senthil Kumar, H. Dutta and S. Sabarinathan, Fuzzy approximations of a multiplicative inverse cubic functional equation, *Soft Computing*, (2020), <https://doi.org/10.1007/s00500-020-04741-x>.
- [22] Z. Gajda, On the stability of additive mappings, *Int. J. Math. Math. Sci.*, **14** (1991) 431–434.