

April 2021

cystatin C serum level and radiological findings in acute ischemic stroke

adham mahmoud ismaeil

Neurology Department, Faculty of Medicine, Zagazig University, Egypt, adham_ismaeel2000@yahoo.com

Sawsan Abdelaziz

Neurology Department, Faculty of Medicine, Zagazig University, Egypt, sayousef@zu.edu.eg

ahmed el sayed badawy

Neurology Department, Faculty of Medicine, Zagazig University, Egypt, badawyemail@yahoo.com

rania yehia helal

Neurology Department, Faculty of Medicine, Zagazig University, Egypt, helal_rania87@yahoo.com

Follow this and additional works at: <https://digitalcommons.aaru.edu.jo/zumj>

Recommended Citation

ismaeil, adham mahmoud; Abdelaziz, Sawsan; badawy, ahmed el sayed; and helal, rania yehia (2021) "cystatin C serum level and radiological findings in acute ischemic stroke," *Zagazig University Medical Journal*: Vol. 27 : Iss. 1 , Article 37.

Available at: <https://digitalcommons.aaru.edu.jo/zumj/vol27/iss1/37>

This Original Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in Zagazig University Medical Journal by an authorized editor. The journal is hosted on [Digital Commons](#), an Elsevier platform. For more information, please contact rakan@aarj.edu.jo, marah@aarj.edu.jo, dr_ahmad@aarj.edu.jo.



Manuscript ID ZUMJ-1905-1229 (R3)

DOI 10.21608/zumj.2019.12875.1229

ORIGINAL ATRICLE**Cystatin C serum level and radiological findings in acute ischemic stroke**Adham Mahmoud Mohamad Ismail¹, Sawsan Abd El Aziz Youssef², Ahmed El Sayed Badawy³, Rania Yehia Helal Mohamed⁴¹ Professor of Neurology, Faculty of Medicine, Zagazig University, Egypt.² Professor of Neurology, Faculty of Medicine, Zagazig University, Egypt.³ Associate Professor of Neurology, Faculty of Medicine, Zagazig University, Egypt.⁴ Assisstant lecturer at Neurology Department, Faculty of medicine, Zagazig University, Zagazig, Egypt.***Corresponding Author:**

Rania Yehia Helal Mohamed

Assisstant lecturer at
Neurology Department,
Faculty of medicine, Zagazig
University, Zagazig, Egypt.
Tel: 01270998497

Submit Date: 2019-05-16

Revise Date: 2019-07-28

Accept Date: 2019-07-30

ABSTRACT

Background: Cerebrovascular stroke is a costly disease. Cystatin C is an important biomarker for acute ischemic stroke due to the background of accumulating evidence indicating a link between vascular disease of the kidney and brain. We investigated the association of serum cystatin C levels with radiological findings of first ever acute ischemic stroke. **Methods:** We included in this prospective cohort study 58 adult patients with first ever acute cerebrovascular ischemic stroke of not more than 72 hours duration with normal kidney functions (26 males and 32 females with age ranged from 33 to 90 years). Serum cystatin C level was measured by enzyme-linked immune-sorbent assay (ELISA). All patients were subjected to brain computed tomography (CT) scan to confirm the diagnosis of acute ischemic stroke and it was repeated after 48 hours. **Results:** We found statistically significant positive correlation between cystatin C serum level and early signs of middle cerebral artery (MCA) infarction and size of infarction. **Conclusion:** There is positive correlation between serum cystatin C level and size of infarction in acute ischemic stroke.

Keywords: Cystatin C, Stroke, Infarct size.**INTRODUCTION**

Cystatin C is a 122-amino acid, 13-kDa protein that is a member of the family of cysteine proteinase inhibitors[1]. It is freely filtered by the glomerulus and is largely reabsorbed and metabolised in the proximal tubules[2]. Cystatin C is a better estimate of renal function, particularly within the normal range of kidney function[3]. The attractiveness of cystatin C as a biomarker for acute ischemic stroke is obvious, in the background of the association between vascular disease of the kidney and brain. Vascular supply to kidney and brain have a lot of similarities. Both are exposed to high-volume blood flow throughout the cardiac cycle and are low resistance end-organs. It therefore seems logical that

microvascular disease in the kidney and brain might be associated[4].

This study was undertaken to evaluate the association of serum cystatin C levels with first ever acute ischemic stroke, and to correlate it with the radiological findings in brain CT scan.

METHODS

We included in this prospective cohort study 58 adult patients with first ever acute cerebrovascular ischemic stroke with normal kidney functions.

Inclusion criteria:

Patients with first ever acute cerebrovascular ischemic stroke who were diagnosed according to the World Health Organization (WHO) criteria^[5], confirmed by brain computed

tomography (CT) , Patients were assessed in the first 72 hours of stroke , Age \geq 18 years.

Exclusion criteria :

Those who suffered from hemorrhagic stroke (intracerebral hemorrhage or subarachnoid hemorrhage),any central nervous system (CNS) disease other than acute cerebral arterial infarction ,patients under thrombolytic therapy,patients with history of head injury ,patients with chronic kidney disease, patients with liver diseases.

Patients were subjected to the following:

Complete history taking, full general and neurological examination.

Laboratory investigations: Both routine and special laboratory investigations were done at Clinical pathology department, Zagazig university hospitals.

-Routine laboratory investigations : complete blood count , liver and kidney function tests , erythrocyte sedimentation rate and lipid profile.

-Special laboratory investigations :measurement of cystatin C serum level in three ml of venous blood which were drawn from all ischemic stroke patients using standard venipuncture techniques, within 72 h of stroke onset. Blood samples were left to clot for 4 hours at room temperature, then centrifuged to obtain the serum which was stored frozen .The levels of cystatin C were measured by the use of double antibody sandwich enzyme-linked immune-sorbent assay technology.

Radiological investigations : Brain CT was done initially for detection of early signs of middle cerebral artery infarction and later on after 48 h for calculation of the infarction volume which was divided as small (less than 1.5 cm³), moderate (1.5 cm³ to 3 cm³) and large (more than 3 cm³)[5]

Follow up:

* All patients were followed up for 1 week by National Institutes of Health Stroke Scale (NIHSS)

*** We classify patients into two groups:**

a) Group with early neurological deterioration (END): in which there is decrease in NIHSS by 2 points or more during the first 7 days

following symptom onset including patients who died[6]

b) Group without early neurological deterioration (Non END)[6].

Written informed consent was obtained from all participants and the study was approved by the research ethical committee of Faculty of Medicine, Zagazig University. The work has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for studies involving humans.

Statistical analysis:

The data were coded and entered using the statistical package SPSS. The data were summarized using mean and SD and median and IQ range for quantitative data, and number and percentage for qualitative data. Student's t-test was used to assess statistical differences between the two groups of quantitative data. Nonparametric Mann–Whitney (MW) and Kruskal–Wallis (KW) tests were used for quantitative variables, which were not normally distributed. P values less than or equal to 0.05 were considered statistically significant.

RESULTS

We included in this prospective cohort study 58 adult patients with first ever acute cerebrovascular ischemic stroke of not more than 72 hours duration with normal kidney functions (26 males and 32 females with age ranged from 33 to 90 years).The mean age was 61.2 \pm 14.5 [Table 1].

Regarding vascular risk factors, 69% of our patients were hypersensitive, 56.9% were diabetic, 27.6% had cardiac risk factors, 17.2% were current smokers [Figure 1]

Table 2 shows that 17.2% of our ischemic stroke patients had early signs of middle cerebral artery (MCA) infarction. Ischemic stroke of anterior circulation had a high percentage, left sided brain lesion are more frequent than right sided lesion and regarding size of infarction, we found that medium sized infarctions are the most prevalent followed by small sized lesions.

We found statistically significant positive correlation between cystatin C serum level and

early signs of middle cerebral artery (MCA) infarction ($p=0.03$) and there was statistically significant positive correlation between cystatin C serum level and size of infarction ($p<0.001$) [Table 3]

There is highly statistically significant association between END and early signs of middle cerebral artery (MCA) infarction and size of lesion. END is associated with large size of infarction [Table 4]

Table 1. Age and sex distribution of studied patients

Variables	Studied patients (n=58)
Age (years):	
Mean \pm SD	61.2 \pm 14.5
Range	33.0 – 90.0
Sex:	
Males	26 (44.8%)
Females	32 (55.2%)

Table 2. Characteristics of brain imaging among stroke patients.

	Frequency	Percentage
Early signs of middle cerebral artery (MCA) infarction:		
Absent	48	40.8%
Present	10	17.2%
Site of lesion:		
Anterior circulation	46	79.3%
Posterior circulation	3	5.2%
Multiple territory	9	15.5%
Side of lesion:		
Left	22	37.9%
Right	36	62.1%
Size of lesion:		
Small	18	31.0%
Medium	23	39.7%
Large	17	29.3%

Table 3. Association between cystatin C serum level and brain imaging.

Size of lesion	Cystatin C ($\times 10^3$ ng/ml)			P
	Median	IQ-Range		
Early signs of middle cerebral artery (MCA) infarction:			MW	
Absent	8.2	5.5-16.3	300	0.03
Present	14.4	11.0-19.5		
Site of lesion:			KW	
Anterior circulation	4.2	3.5-9.0	94	0.2
Posterior circulation	7.3	3.0-11.0		
Multiple territory	5.5	4.5-13.0		
Side of lesion:			MW	
Left	7.3	4.5-13.5	260	0.1
Right	5.9	3.0-15.2		
size of lesion :			KW	
Small	4.2	3.4 – 9.4	204	<0.001
Medium	7.5	6.2 – 14.2		
Large	19.8	12.6 – 30.8		

Table 4. Association between END and brain imaging in the studied patients.

Size of lesion	END cases (n=13)		Non-END cases (n=45)		χ^2	P
	No	%	No	%		
Early signs of middle cerebral artery (MCA) infarction:						
Absent	4	8.3	44	91.6	fisher	<0.001
Present	9	90	1	10		
Site of lesion:						
Anterior circulation	10	21.7	36	78.3	0.1	0.8
Posterior circulation	1	33.3	2	66.7	Fisher	
Multiple territory	2	22.2	7	77.8	Fisher	
Side of lesion:						
Left	7	31.8	15	68.2	1.8	0.1
Right	6	16.7	30	83.3		
Size of lesion:						
Small (n=18)	0	0.0	18	100	16.3	<0.001
Medium (n=23)	3	13.0	20	87.0		
Large (n=17)	10	58.8	7	41.2		




Figure 1. Risk factors in studied patients.

DISCUSSION

As one organ may tell us about the other, there is association between stroke and impaired renal functions as observed in previous various studies and this was based on evidence indicating a link between vascular disease of the brain and kidney[7].

Our study included 58 first ever acute cerebrovascular ischemic stroke patients. Their mean age was 61.2 ± 14.5 years with a range (33-90) which approximately near other studies [8-10] who stated that the mean age of their patients was (60 ± 16.7 , 63.35 ± 11.18 and 61.5 ± 10.5) respectively.

Regarding gender difference as a stroke risk, we found that the risk was higher among females than males and represents 55.2% and 44.8% of our stroke patients. Other studies[10-12] are in agreement with our results .

With regard to risk factors, 69% of our patients were hypertensive, 56.9% were diabetic, 27.6% had cardiac risk factors, 17.2% were current smokers. These results were consistent with a previous study [13] who reported that 76.6% of their patients were hypertensive and 48.4% of them were diabetic.

Our study showed that 17.2% of our ischemic stroke patients had early signs of middle cerebral artery (MCA) infarction, while the percentage was 20.8% by another previous research[10].

In our study, ischemic stroke of anterior circulation had a higher percentage (79.3%) ,These results were consistent with previous studies[14-19].

However, A previous study [20] found that among patients with acute ischemic stroke (AIS), 48.1% patients suffered anterior circulation stroke; 23.8% patients suffered posterior circulation stroke and the remainder suffered multiple infarction including anterior and posterior circulation stroke.

Another study[21] found that among patients with AIS, 62% patients suffered anterior circulation stroke; 23% posterior circulation stroke and 15% multiple infarction including anterior and posterior circulation stroke.

In our study, left sided weakness (LSW) were more frequent than right sided weakness (RSW) .This is in agreement with a study [22] who reported 53.1 % of ischemic stroke patients was with LSW, 46.8 % with RSW and other study [17] in which 50% of cases presented with LSW, in comparison with 38% presented with RSW .

Regarding size of infarction, we found that medium sized infarctions were the most prevalent (39.7%) followed by small sized lesions (31.0%) . A previous research[9] also found that out of the studied ischemic stroke patients 36.6% had large size infarction , 40 % had medium size infarction and 23.3% had small size infarction ,while Other research [23] found that 44% of their patients had small size infarction , 27.9 % had medium size infarction and 27.6% had large size infarction . This variation between our study and their study may be caused by differences in the sample size

and the type of imaging as their patients were imaged with magnetic resonance imaging [9].

As regards association between cystatin C serum level and brain imaging. We found statistically significant positive correlation between cystatin C serum level and early signs of middle cerebral artery (MCA) infarction and there was statistically significant positive correlation between cystatin C serum level and size of infarction. Cystatin C serum level is higher with larger size of infarction. This is in agreement with a study[24] who reported that patients with higher cystatin C serum level had larger size of infarction. Under ischemic stimuli, neuronal cells, including neurons and astrocytes, may release high levels of cystatin C[25].

Higher cystatin C level was consistently associated with a larger white matter lesion volume [26].

Elevated level of serum cystatin C is associated with the total burden of cerebral small vessel disease in patients with acute lacunar stroke independent of conventional risk factors[4]

In the present study, There was no statistically significant correlation between cystatin C serum level and site or side of the lesion. This is in agreement with a previous study [27] who found that serum cystatin C was not associated with the location of ischemic area.

Early neurological deterioration (END) after acute ischemic stroke is associated with several clinical, biochemical and radiological markers. This deterioration may be due to different processes, for example development of cerebral edema, progression of thrombus and hemorrhagic transformation; it affects up to 25% of all acute stroke patients and raises the risk of death or disability two to three-fold [28].

Identifying those markers for END will lead to target curative interventions for prevention [

Regarding association between END and brain imaging. We found statistically significant positive correlation between END and early signs of middle cerebral artery (MCA) infarction and there was highly statistically

significant association between END and size of lesion. END is associated with large size of infarction. This is in agreement with a previous study[21]

However, There was no statistically significant correlation between END and site or side of the lesion.

CONCLUSION

In view of our results, we can conclude that there is statistically significant positive correlation between serum cystatin C level and early signs of middle cerebral artery (MCA) infarction and size of infarction.

We recommend that specialists should consider adding the use of cystatin C serum level to their routine admission testing in patients with acute ischemic stroke, since it is significantly correlated with size of infarction and for prediction of early neurological deterioration of acute ischemic stroke.

We recommend further studies with larger sample size and longer duration of follow up, with measurement of cystatin C during follow up period to assess the fluctuation and variability in its value with time in stroke patients.

Conflict of Interest: Nothing to declare.

Financial disclosure: Nothing to declare.

REFERENCES

1. **Newman DJ. Cystatin C.** Ann Clin Biochem 2002; 39: 89-104.
2. **Makin SD, Cook FA, Dennis MS and Wardlaw JM.** Cerebral small vessel disease and renal function: systematic review and meta-analysis. Cerebrovasc Dis. 2015;39:39-52.
3. **Sur A, Mishra P, Swain M and Mohapatra N.** Study of Serum Cystatin C, Lipid Profile and Electrolytes in Acute Stroke Patients. International Journal of Clinical Biochemistry and Research 2015;2(4):194-197.
4. **Yang S, Cai J, Lu R, Wu J, Zhang M and Zhou X.** Association Between Serum Cystatin C Level and Total Magnetic Resonance Imaging Burden of Cerebral Small Vessel Disease in Patients With Acute Lacunar Stroke. J. Stroke Cerebrovasc. Dis. 2017; 26 : 186-191.
5. **Bogousslavsky J, Diserens K and Rothacher G.** Stroke selected topics in Seminars in clinical neurology. New York 2006; 4: 1-65.
6. **Kim T, Kang M, Jeong H, Kim C, Kim Y,**

- Nam K et al.** Cystatin C is a useful predictor of early neurological deterioration following ischaemic stroke in elderly patients with normal renal function. *European Stroke Journal* 2015;2:23-30.
7. **Oksala N, Salonen T, Strandberg T, Oksala A, Pohjasvaara T, Kaste M, et al.** Cerebral small vessel disease and kidney function predict long-term survival in patients with acute stroke. *Stroke* 2010; 41:1914–1920.
 8. **Laskowitz D, Kasner S, Saver J, Remmel K and Jauch E.** Clinical usefulness of a biomarker-based diagnostic test for acute stroke: the Biomarker Rapid Assessment in Ischemic Injury (BRAIN) study. *Stroke* 2010 ;(40):77–85.
 9. **Arafa A, Sarhan A, El Sharkawy k and Mohamed T.** Red blood cell distribution width as a predictor of clinical outcome in acute stroke patients. MSc thesis 2017. Zagazig university.
 10. **Fathy S, Abd El Gawad E, El Motayam A and Mohammed W.** Blood Biomarkers ;Brain Natriuritic peptide, d-Dimer and creatine kinase-MB in patients with cardioembolic stroke. MD thesis 2018. Zagazig University.
 11. **Al-Shammri S, Shahid Z, Ghali A, Mehndiratta MM, Swaminathan TR, Chadha G, et al.** Risk factors, subtypes and outcome of ischaemic stroke in Kuwait— A hospital-based study. *Med Princ Pract* 2003 ;12: 218–223.
 12. **Somay G, Topaloğlu P, Somay H, Sato M, Iso H, Sato S et al.** Cerebrovascular risk factors and stroke subtypes in different age groups: a hospital-based study. *Turk J Med Sci.* 2006 ; 36(1) : 23-39.
 13. **Ibrahim G, Abd El Gawaad E , Mohammad H and Mohammad T .** Assessment of Extracranial Carotid Arteries In Acute Ischemic Stroke: Correlation With Risk Factors. MSc thesis 2017. Zagazig University.
 14. **Kang DW, Yoo SH, Chun S, Kwon KY, Kwon SU, Koh JY et al.** Inflammatory and hemostatic biomarkers associated with early recurrent ischemic lesions in acute ischemic stroke. *Stroke* 2009 ; (40): 1653–1658.
 15. **Abdel Ghani A, Zaitoun A and Gawish H.** Prognostic value of d-Dimer in diffusion weighted-MRI defined early ischemic stroke recurrence. *Egypt Journal of Neurology, Psychiatry and Neurosurgery* 2011; (48): 215-222.
 16. **Jung KW, Shon YM, Yang DW, Kim BS and Cho AH.** Coexisting carotid atherosclerosis in patients with intracranial small- or large-vessel disease. *Journal of Clinical Neurology* 2012 ;(8): 104-108.
 17. **Danthala M and Lakshmaiah V.** Correlative study of serum cystatin C levels with severity of acute ischemic stroke. *J Clin Biomed Sci* 2013 ; 3(4):188-92.
 18. **Wiseman S, Marlborough F, Doubal F, Webb DJ and Wardlaw J.** Blood markers of coagulation, fibrinolysis, endothelial dysfunction and Inflammation in lacunar stroke versus non-lacunar stroke and non-stroke: systematic review and meta-analysis. *Cerebrovascular Disorders* 2014; (37):64–75.
 19. **Zi WJ and Shuai J .** plasma d-dimer levels are associated with stroke subtypes and infarction volume in patients with acute ischemic stroke. *PLoS ONE* 2014 ; 9(1): 1-8.
 20. **Yang X, Gao S, Ding J, Chen Y, Zhou X and Wang J.** Plasma D-dimer predicts short-term poor outcome after acute ischemic stroke. *PLOS ONE* 2014 ;(2):1-7.
 21. **Huang Z, Wang Q, Dai Y, Lu H , Liang X, Hu H and Liu X.** Early neurological deterioration in acute ischemic stroke: A propensity score analysis *Journal of the Chinese Medical Association* 2018;81: 865-870.
 22. **Pan SL, Wu SC, Lee TK, Canada RW and Newman LU.** Location and size of infarct on functional outcome of noncardioembolic ischemic stroke. *Disability and Rehabilitation* 2005; 28(16): 977 – 983.
 23. **Mustanoja S, Putaala J, Haapaniemi E, Strbian D, Kaste M and Tatlisumak T.** Multiple brain infarcts in young adults: clues for etiologic diagnosis and prognostic impact. *Eur J Neurol* 2012 ;20:216-222.
 24. **Xiao D, Liu H, Zhang H and Luo Y.** Impact of cystatin C levels on infarct size and hemorrhage volume in acute cerebral stroke. *J Neurol* 2012 ; 259:2053–2059.
 25. **Gauthier S, Kaur G, Mi W, Tizon B and Levy E.** Protective mechanisms by cystatin C in neurodegenerative diseases .*Front Biosci (Schol Ed).* 2012; 3: 541–554.
 26. **Akoudad S, Sedaghat S, Hofman A, Koudstaal P, Lugt A, Ikram M et al.** Kidney function and cerebral small vessel disease in the general population . *World Stroke Organization Am J Physiol Renal Physiol* 2015; 294:F1–9.
 27. **Jeong H, Kim B, Yang M, Han M and Bae H.** Neuroimaging Markers for Early Neurologic

Deterioration in Single Small Subcortical Infarction. *Stroke*.2015 ;46:687-691.

- 28. Helleberg BH, Ellekjær H, Rohweder G and Indredavik B** .Mechanisms, predictors and clinical impact of early neurological deterioration: the protocol of the Trondheim early neurological deterioration study. *MCNeurology* 2014, 14:201-222.

- 29. Zhang X, Sun Z, Ding C, Tang Y, Jiang X, Xie Y et al.** Metabolic Syndrome Augments the Risk of Early Neurological Deterioration in Acute Ischemic Stroke Patients Independent of Inflammatory Mediators: A Hospital-Based Prospective Study .*Oxidative Medicine and Cellular Longevity*.2016, 83:1-6.

Cite This Article

ismaeil, A., yousef, S., badawy, A., helal, R. cystatin C serum level and radiological findings in acute ischemic stroke. *Zagazig University Medical Journal*, 2021; (140-147): -. doi: 10.21608/zumj.2019.12875.1229