

1-27-2012

Global prevalence of periodontitis: A literature review

Shaju JACOB

Follow this and additional works at: <https://digitalcommons.aaru.edu.jo/iajd>

Recommended Citation

JACOB, Shaju (2012) "Global prevalence of periodontitis: A literature review," *International Arab Journal of Dentistry*. Vol. 3: Iss. 1, Article 6.

Available at: <https://digitalcommons.aaru.edu.jo/iajd/vol3/iss1/6>

This Original Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in International Arab Journal of Dentistry by an authorized editor. The journal is hosted on [Digital Commons](#), an Elsevier platform. For more information, please contact rakan@aarj.edu.jo, marah@aarj.edu.jo, u.murad@aarj.edu.jo.

GLOBAL PREVALENCE OF PERIODONTITIS: A LITERATURE REVIEW

Shaju Jacob*

Abstract

The prevalence of periodontitis is dependent on the studied population and the case definition adopted. Periodontal diseases assume a greater global importance as the senior population is on the rise in most countries.

The aim of this article is to review the global prevalence of periodontitis. Studies describing the prevalence of periodontitis in the world were searched through various indices using the key words 'periodontitis, national survey, prevalence' and reviewed. Many national surveys have used the WHO method for assessing prevalence. Yet several studies have not used the community periodontal index (CPI) for periodontitis case definition. Most surveys have used partial recording which can underestimate the periodontitis prevalence. There is a need for a standardized and valid population-based case definition of periodontitis.

Keywords: prevalence – periodontitis - survey.

IAJD 2012;3(1):26-30.

PRÉVALENCE MONDIALE DE LA PARODONTITE: REVUE DE LITTÉRATURE

Résumé

La prévalence de la parodontite dépend de la population étudiée et des critères qui la définissent. Les maladies parodontales dans le monde gagnent en importance avec la tendance qu'a la population dans la plupart des pays développés à vieillir davantage. L'objectif de cette étude est d'évaluer la prévalence mondiale de la parodontite. Les études rapportant la prévalence de la parodontite dans le monde ont été recherchées en utilisant les mots clés « parodontite, enquête nationale, prévalence » et analysées. De nombreuses enquêtes nationales ont utilisé la méthode de l'OMS pour évaluer la prévalence. Pourtant, plusieurs études n'ont pas utilisé l'indice de la communauté parodontale pour la définition de cas de parodontite. La plupart des enquêtes ont utilisé l'enregistrement partiel qui sous-estime la prévalence de la parodontite. Il est nécessaire d'établir une définition de la maladie parodontale qui soit standard et valide quelle que soit la population étudiée.

Mots- clés: prévalence – parodontite - enquête.

IAJD 2012;3(1):26-30.

* MDS Periodontics
Associate Professor
Faculty of Dentistry,
Ayush and Health Sciences University of Chhattisgarh
shajujacob@yahoo.com

Introduction

Periodontal disease is one of the two major dental diseases that affect human populations worldwide at high prevalence rates. The World Health Organization (WHO) [1] reported that 10 – 15% of the world populations suffer from severe periodontitis. Yet any information on the periodontitis prevalence must be interpreted in light of the population studied and the case

definition used for periodontitis [2]. Available population-based data originate from studies encompassing a wide range of objectives, designs and measurement criteria [3]. This can hamper the comparability of the prevalence's estimates among various populations.

The aim of this study is to evaluate the prevalence of periodontitis among different populations by reviewing the published literature.

Materials and Methods

Various indices including PubMed, MedInd, WHO, DAOJ were searched using the key words "periodontitis, national survey and prevalence". Published material reporting prevalence on a national level or on a major section (population) of the country were selected. This resulted in the identification of 31 publications of 19 countries.

Results

Prevalence of periodontitis estimated in a representative sample of a population depends on the case definition used and the studied population. WHO recommends the use of the Community Periodontal Index (CPI) for prevalence studies so that the results can be compared among different populations. With a partial recording, disease is defined (periodontitis; CPI=3) when the CPI probe records between 3.5mm and 5.5mm in at least one of the teeth while severe periodontitis (CPI=4) is defined when the probing depth is >5.5mm.

Many national surveys have used the WHO criteria in the case definition of periodontitis.

In a cross-sectional analysis of the Japanese National data [4], 42.5% exhibited periodontitis at CPI≥3, and the prevalence dropped to 9.5% at the CPI=4. In the national survey of Korea [5] done on a proportionately region-stratified clustered sampling, 10.3% had CPI≥3 and 3% CPI=4. The authors observed that the prevalence was lower than that reported by other studies performed in industrialized nations like the United States, Australia and Japan.

Prevalence of periodontitis was significantly higher in males and increased with increasing age groups, as reported by a national survey in India [6] using CPI for disease assessment. Moderate periodontitis was seen in 17.5% of 35-44 years-old and in 21.4% in 65-74 years-old, whereas severe disease defined as at least one tooth with >6mm probing depth was observed in 7.8% and 18.1% of the two age groups, respectively. Similar conclusions were drawn from a CPITN survey involving 1150 Chileans [7] (2 age groups: 35-44 years-old and 65-74 years-old) though the percentages were higher since the prevalence of chronic inflammatory periodontal disease (codes 3 + 4) was 91% in subjects aged 35-44 years-old and 100% in subjects aged 65-74 years-old. The total prevalence for both age cohorts

was 92.2%. Prevalence of periodontal disease was slightly lower in females and severity was significantly higher in males.

The prevalence of severe periodontal disease is lower than the chronic periodontitis, a hypothesis highlighted by many studies. A survey among Canadian adults [8] included 2110 adults aged 35-44 years in Quebec and used a stratified sample of randomly selected census areas and households in Quebec. Periodontal pockets of ≥4mm were observed in 73.6% persons, and those ≥6mm were prevalent in 21.4% of the examined persons. Also, in a national study [9] carried out in France in 1993 to assess the periodontal status of the population aged 35-44 years, a representative sample of 1000 subjects was studied using the CPITN index. Gingivitis prevalence was high (80.4%) while 26.6% of dentate subjects had shallow pockets (4-5 mm). Deep pockets (> 6 mm) were rare (1.6%). In the national survey of Finland [10], 5255 persons were examined for periodontitis from the total sample of 8028 adults (≥30 years). 64% had periodontitis (CPI≥3) while 21% had severe periodontitis (CPI=4). The age group of 35-44 years-old reported a prevalence of 61% and 14% of moderate and severe periodontitis, respectively. In another study in Southern Finland [11], 325 workers (aged 38-65 years) with access to subsidized dental care and 174 controls without access were periodontally examined. The authors [11] stated that deep pockets ≥ 6 mm were observed in 5% for the health care subsidized workers and 11% of the controls.

Krustrup and Petersen [12] did a cross-sectional study on a random sample of 1,115 Danish adults aged 35-44 years and 65-74 years. 42% of the younger adults had moderate periodontitis while 6.2% had severe periodontitis. In the older participants group, the prevalence of moderate and severe diseases was 82% and 20%, respectively. There was no difference in prevalence between males and females in both age groups.

In a national study of United Kingdom adults, Morris et al. [13] reported that pockets greater than 5.5mm were uncommon (5%) in the total adult population. The prevalence in the 35-44 years group was 5% and thereafter it increased to 17% in the 55-64 years group and 15% among those ≥ 65 years. In the latter group, only 4% had pockets of 8.5mm.

WHO [14] compiled data from prevalence studies conducted in many countries on periodontal status. Severe periodontitis was considered when a subject had at least one site with ≥6mm. The lowest prevalence of severe periodontitis was seen in Madagascar and Hungary (3%). China, Brazil, Denmark, French Polynesia, Pakistan, Poland, Japan, Norway and Malaysia reported a prevalence of less than 10% of severe periodontitis [14]. Prevalence of more than 20% was seen in the populations of Bangladesh, Canada, Germany, India, Belarus and Chile. In India, according to WHO data, the prevalence of severe periodontitis was in the range of 19 to 32%.

Corbet et al. [15], in a cross sectional survey of a province in Southern China, selected 1572 ages 35-44 years and 1286 aged 65-74 years by a two-stage cluster sampling method. A complex case definition modified from CPI was used for prevalence estimates separately for the two age groups to account for the age influence on the prevalence estimates. The periodontal disease was defined as having at least 2 sextants with a clinical attachment level (CAL) ≥4mm; 40% of 35-44 years-old had the disease, while 65-74 years reported a prevalence of 34.5% for their case definition of at least two sextants with CAL ≥6mm. 4% of 35-44 years and 6.5% of 65-74 years had severe disease (CPI=4). The periodontal status of 8462 residents of Keelung, Taiwan, was studied by Lai et al. [16] in a cross-sectional survey, using the CPI and CAL indexes at subject (prevalence) and sextant levels (severity). 95% had some signs of periodontal disease, of whom 29.7% had periodontal pockets >3mm

and 35% had attachment loss >3 mm. In a Thai survey [17] of high income retired persons of 50–73 years, severe disease was diagnosed when a person had a mean CAL ≥ 4 mm. Severe disease was prevalent in 16% of the population.

The United-States was the first to recognize the importance of surveys in the determination of periodontal disease's prevalence with their landmark NHANES I [18] conducted between 1971 and 1974 on a probability sample of approximately 28,000 subjects. The periodontal conditions were assessed by the Periodontal Index (PI) on all teeth present. The average PI scores increased steadily with age, and were higher in males than females (0.96 vs. 0.7) and in blacks than in whites (1.28 vs. 0.76). These differences between the four subgroups were consistent across all age groups.

The Dental Health Outcomes Survey [19] conducted in 1981 was a household-based survey in which 1792 participants, aged ≥ 19 years were selected for periodontal examination using a multistage probability sampling design from all American states. Thirty-six percent of the participants had periodontitis (at least one tooth with probing depth ≥ 4 mm) and only 15% of the participants had no periodontal disease. In the whole population, 28% had moderate periodontitis (≥ 1 tooth with a probing depth of 4–6 mm) and 8% had advanced periodontitis (≥ 1 tooth with a probing depth ≥ 7 mm).

The National Survey of Oral Health [19] was conducted by the NIDR in 1985–1986 to assess the oral health status of adults in the United States. The targeted population was employed adults (aged 18–64 years) and seniors (aged ≥ 65 years). The sampling frame included a multistage sampling design from which 15,132 persons 18–64 years of age and 5,686 persons aged 65–85 years were examined. 14.3% of employed adults and 22.2% of seniors had one or more teeth with ≥ 4 mm probing depth. The prevalence of ≥ 4 mm probing depth increased with age, from 4% in the 18–19 years group to 22% in

the 60–64 years group. However, in the seniors group, it decreased slightly with age. The prevalence of attachment loss of ≥ 3 mm, ≥ 4 mm, and ≥ 5 mm were 44.6%, 24.1% and 13.6%, respectively, in employed adults and 86.3%, 68.2%, and 51.7%, respectively, in seniors.

The NHANES III [20] survey was conducted between 1988–1994; 3.1% had advanced periodontitis, 9.5% had moderate periodontitis and 22% had mild periodontitis. The survey used a complex case definition combining probing depth and clinical attachment level assessment as well as the determination of furcation involvement. The prevalence of periodontitis increased steadily with increasing age. However, moderate and advanced periodontitis increased in prevalence between 30 and 70 years of age and then levelled off to slightly decline thereafter.

In the NHANES IV [21], the periodontitis was defined as the presence of at least 3 sites with CAL > 4 mm and at least 2 sites with PD > 3 mm. Black (6.8%) exhibited the highest prevalence of periodontitis, followed by Mexican-Americans (4.6%) and White (3.8%).

In the national survey [22] conducted between 1986 and 1987 to evaluate the oral health of 14,013 American children aged 13 to 19 years, CAL ≥ 3 mm was used as a threshold for periodontitis. Aggressive periodontitis defined by having attachment loss ≥ 3 mm in ≥ 4 teeth including two or fewer canines, premolars and second molars was present in 0.4% of 13–15 years-old, 0.8% of 16–19 years old, 0.06% of White, 2.6% of Black and 0.5% of Hispanic. The prevalence rates for chronic (at least one tooth with ≥ 3 mm attachment loss) periodontitis were higher and were 2.3% and 3.2% in the two age groups, respectively.

In a cross sectional survey [23] as part of the Erie County study in New-York, prevalence was estimated by mean CAL. 8.6% had severe disease (mean CAL ≥ 4 mm). Prevalence ranged from 14.7% in the 35–44 years group to 23% in the 65–74 years group.

The Piedmont Health study [24] is a longitudinal study based on a stratified, clustered, random sample of people aged ≥ 65 years in five contiguous North Carolina counties. Severe periodontitis was defined as the presence of ≥ 4 sites with loss of attachment ≥ 5 mm and ≥ 1 of those same sites had a pocket depth of more than 3 mm; the estimates were 16% for the White and 46% among the Black.

Bial & Mellonig [25] used clinical and radiologic (panoramic and bitewings) examinations to screen a large group comprising 49,380 male naval recruits in Okinawa, Seattle, 17 to 32 years old. Cases demonstrating greater bone loss on the permanent first molars and/or incisors than on other teeth were classified as having juvenile periodontitis. It was estimated that 0.55% of the subjects had bone loss, 0.37% subjects had early onset (juvenile) periodontitis, 0.13% subjects with rapidly progressing periodontitis and 0.05% with isolated bone defects.

In Australia, the National Dental Telephone Interview Survey 2002 was conducted in Adelaide [26]. It comprised 709 persons, aged 45–54 years. The prevalence of CAL ≥ 4 mm, ≥ 5 mm and ≥ 6 mm was 66%, 37% and 19%, respectively. The prevalence of PD ≥ 6 mm was 10.2%.

In the National Survey of Australia 2007 [27], approximately one of five Australian adults had moderate (20.5%) or severe (2.4%) periodontitis based on the Center of Disease Control (CDC) classification [28]. The prevalence for age group 35–54 years was 24.5%; males 30.4%, females 18.6%. The percentage of the Australian population with CAL ≥ 4 mm was 42.5%. The prevalence of CAL ≥ 4 mm was lowest in the most recent generation (17.4%) and increased across the generations, with a prevalence of 49% in the 1950–69 generation, 73.0% in the 1930–49 generation and 80.5% in the pre-1930 generation.

The national survey [29] of France was conducted on persons between 35 and 64 years, living in all 22 administrative regions of metropolitan France

(N= 2144). The sample of 2144 subjects was 51% women, with a median age of 49.6 years (Q1 = 42.44 years; Q2 = 58.37 years). Population prevalence estimates indicated that loss of attachment ≥ 5 mm was 46.68% and probing depth (> 5 mm) was 10.21%. One or more deep pocket (> 5 mm) was present in 10.2% of the sample. 1.73% presented probing depths of >7 mm. Schürch et al. [30] found the prevalence of deep pockets (≥ 6 mm) was low, 2–3%, in a sample of 206 persons randomly selected from Canton of Berne, Switzerland while 25% persons of all age groups had moderate periodontal disease.

Discussion

The CPITN is a treatment index commonly used for estimating the prevalence of periodontitis in population; it is simple and easily reproducible. WHO modified it as CPI by introducing the CAL measurement. Yet many drawbacks were pointed, mainly the use of a partial recording. CPITN overestimated prevalence and severity of periodontal diseases in younger age groups and underestimated them among older subjects [28]. The performance of a partial recording system is affected by the actual prevalence of periodontal disease in the population in question. The less frequent the disease, the more difficult it becomes for a partial recording system to detect it and thus may lead to a greater underestimation of its prevalence. A full-mouth examination remains the best method for accurately assessing the prevalence and severity of periodontal disease in a population. Yet in a large survey, it becomes practically difficult to have a whole mouth periodontal recording.

The lack of standardized study design, definition of periodontal disease, methods for disease detection and measurement and criteria for subject selection markedly limit interpretation and analysis of available population-based periodontal disease data from around the world [3]. Significant disparities appear to exist in the level

of periodontitis among young, adult and senior populations in the world. Subjects of African ethnicity seem to have the highest prevalence of periodontitis, followed by Hispanics and Asians. Disparities in periodontal status appear to be related to socio-economic levels.

Conclusions

After reviewing the most relevant published studies reporting the prevalence of periodontal diseases, the following conclusions can be drawn:

- CPI/CPITN was the most used index for assessing periodontitis in many populations.

- The use of partial recording underestimates the prevalence of the disease, thus reduces the accuracy of the results.

- The use of a variety of case definitions in the different surveys makes comparison difficult.

These results highlight the need for a valid standardised case definition of the periodontal disease, a well-defined study design and calibrated operators.

References

- Petersen PE, Ogawa H. Strengthening the prevention of periodontal disease: the WHO approach. *J Periodontol.* 2005;76:2187-93.
- Burt B. Position paper: Epidemiology of periodontal diseases. *J Periodontol.* 2005;76:1406-19.
- Kingman A, Albandar JM. Methodological aspects of epidemiological studies of periodontal diseases. *Periodontol* 2000 2002;29:11-30.
- Ojima M, Hanioka T, Tanaka K, Inoshita E, Aoyama H. Relationship between smoking status and periodontal condition, findings from national databases in Japan. *J Periodontal Res.* 2006; 41:573-79.
- Choi Y, Baek H, Song K, Han J, Kwon H, Lee SG. Prevalence of periodontitis and associated risk factors in Korean adults: Korean National Oral Health Survey 2006. *J Korean Acad Periodontol* 2009;39:261-68.
- Bali. National Oral health survey 2001-2003. Dental Council of India 2004.
- Gamonal JA, Lopez NJ, Aranda W. Periodontal conditions and treatment needs, by CPITN, in the 35-44 and 65-74 year-old population in Santiago, Chile. *Int Dent J.* 1998;48:96-03.
- Brodeur JM, Payette M, Benigeri M, Charbonneau A, Olivier. Periodontal diseases among Quebec adults aged 35-44 years. *Can Dent Assoc.* 2001;67:34-41.
- Bourgeois D, Hescot P, Doury J. Periodontal conditions in 35-44-yr-old adults in France, 1993. *J Periodontal Res* 1997;32:570-74.
- Suominen-Taipale L, Nordblad A, Vehkalahti M, Aromaa A, eds. Oral health in the Finnish adult population. Health 2000 Survey. Publications of the National Public Health Institute 2008.
- Ahlberg J, Tuominen R, Murtomaa H. Periodontal status among male industrial workers in Southern Finland with or without access to subsidized dental care. *Acta Odontol Scand* 1996;54:166-70.
- Krustrup U, Erik Petersen P. Periodontal conditions in 35-44 and 65-74-year-old adults in Denmark. *Acta Odontol Scand.* 2006;64:65-73.
- Morris AJ, Steele J, White DA. The oral cleanliness and periodontal health of UK adults in 1998. *Br Dent J.* 2001;191:186-92.
- World Health Organization. The WHO Global Oral Health Data Bank. Geneva World Health Organization; 2003.
- Corbet E, Wong M, Lin H. Periodontal conditions in adult Southern Chinese. *J Dent Res.* 2001;80:1480-85.
- Lai H, Lo M, Wang P, Wang T, Chen T, Wu G. A community-based epidemiological study of periodontal disease in Keelung, Taiwan: A model from Keelung community-based integrated screening programme (KCIS No. 18). *J Clin Periodontol.* 2007;34:851-59.
- Torrunguang K, Tamsailom S, Rojanasomsith K, Sutdhibhisal S, Nisapakultorn K, Vanichjakvong O et al. Risk indicators of periodontal disease in older Thai adults. *J Periodontol.* 2005;76:558-65.
- Albandar JM. Periodontal diseases in North America. *Periodontol* 2000. 2002;29:31-69.
- Miller A, Brunelle J, Carlos J, Brown L, Löe H. Oral health of United States adults. The national survey of oral health in U.S. employed adults and seniors: 1985-86. NIH Publication no. 87-2868. Bethesda, Maryland: U.S. Department of Health and Human Services, National Institutes of Health, 1987.
- Albandar JM, Brunelle J, Kingman A. Destructive periodontal disease in adults 30 years of age and older in the United States, 1988-1994. *J Periodontol.* 1999;70:13-29.
- Borrell L, Burt B, Taylor G. Prevalence and trends in periodontitis in the USA: the [corrected] NHANES, 1988 to 2000. *J Dent Res.* 2005;84:924-30.
- Albandar JM, Brown L, Löe H. Clinical features of early onset periodontitis. *J Am Dent Assoc.* 1997;128:1393-99.
- Machtei E, Hausmann E, Dunford R, Grossi S, Ho A, Davis G et al. Longitudinal study of predictive factors for periodontal disease and tooth loss. *J Clin Periodontol.* 1999;26:374-80.
- Beck J, Koch G, Rozier R, Tudor G. Prevalence and risk indicators for periodontal attachment loss in a population of older community-dwelling blacks and whites. *J Periodontol.* 1990;61:521-28.
- Bial J, Mellonig J. Radiographic evaluation of juvenile periodontitis (periodontosis). *J Periodontol.* 1987;58:321-26.
- Bourgeois D, Bouchard P, Mattout C. Epidemiology of periodontal status in dentate adults in France, 2002-2003. *J Periodontal Res.* 2007;42:219-27.
- Slade G, Spencer A, Roberts-Thomson K (Editors) Australia's dental generations: the National Survey of Adult Oral Health 2004-06. AIHW cat. no. DEN 165. Canberra: Australian Institute of Health and Welfare (Dental Statistics and Research Series No. 34).
- Page R, Eke P. Case definitions for use in population-based surveillance of periodontitis. *J Periodontol.* 2007;78:1387-99.
- Bourgeois D, Hescot P, Doury J. Periodontal conditions in 35-44-yr-old adults in France, 1993. *J Periodontal Res* 1997;32:570-574.
- Schürch E Jr, Minder C, Lang N, Geering A. Periodontal conditions in a randomly selected population in Switzerland. *Community Dent Oral Epidemiol.* 1988;16:181-86.
- Do G, Spencer A, Roberts-Thomson K, Ha H. Smoking as a risk indicator for periodontal disease in the middle-aged Vietnamese population. *Community Dent Oral Epidemiol.* 2003;31:437-46.