

2020

EFFECT OF ARM CYCLING CROSS TRAINING AND VISUAL FEEDBACK ON MODULATION OF LOWER LIMB SPASTICITY IN DIAPLEGIC CEREBRAL PALSY CHILDREN

Hala Ibrahim Kassem

Delta University for Science and Technology, halakassem146@gmail.com

Follow this and additional works at: <https://digitalcommons.aaru.edu.jo/dusj>



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Kassem, Hala Ibrahim (2020) "EFFECT OF ARM CYCLING CROSS TRAINING AND VISUAL FEEDBACK ON MODULATION OF LOWER LIMB SPASTICITY IN DIAPLEGIC CEREBRAL PALSY CHILDREN," *Delta University Scientific Journal*. Vol. 3 : No. 2 , Article 6.

Available at: <https://digitalcommons.aaru.edu.jo/dusj/vol3/iss2/6>

This Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in Delta University Scientific Journal by an authorized editor. The journal is hosted on [Digital Commons](#), an Elsevier platform. For more information, please contact rakan@aar.edu.jo, marah@aar.edu.jo, u.murad@aar.edu.jo.



EFFECT OF ARM CYCLING CROSS TRAINING AND VISUAL FEEDBACK ON MODULATION OF LOWER LIMB SPASTICITY IN DIAPLEGIC CEREBRAL PALSY CHILDREN

Hala I. Kassem

Prof. of Physical Therapy for Pediatrics, Dean of the Faculty of Physical Therapy, Delta University for Science and Technology, Egypt

Abstract

Spastic diaplegia is characterized by motor inco-ordination primarily in the lower extremities, that impairs many functional abilities, most notably ambulation. This study was conducted to determine the effect of arm cycling and visual feedback on controlling spasticity of the lower limbs, in spastic diaplegic cerebral palsied children. Thirty spastic diaplegic cerebral palsied children from both sexes, ranging in age from six to eight years represented the sample of the study. All subjects were ambulant with an abnormal gait pattern characterized by hip and knee flexion and ankle planter flexion. The subjects were divided randomly into two groups of equal number (A and B). Group A (control) received a designed physical therapy program in addition to arm pedaling using a reciprocate bicycle with an unloaded cycling for thirty ramps / minute for five minutes, repeated three times with five minutes rest in between. Group B (study) received the same program given to group A with applying the arm pedaling in front of a large mirror to produce visual feedback. Evaluation using 3D Motion Analysis was conducted to determine hip, knee and ankle joints' angles before and after three months of treatment. The results revealed significant improvement in all measuring variables of the two groups, when comparing their pre and post treatment mean values. Highly significant improvement was observed in favor of group B when comparing the post treatment results of the two groups. Improvement observed in the lower limbs joints' angles may be due to enhancement of the effect of cross education training using arm cycling via visual feedback, amelioration of motor learning which led to controlling spasticity of the lower limbs, in the diaplegic cerebral palsied children.

Key Words: Arm Cycling, Cross Training, Visual Feedback, Diaplegic Cerebral Palsy.

Introduction

Cerebral palsy (CP) is a number of non-progressive syndromes of disability which results from various insults to different areas within the developing nervous system¹. Impairment in CP may include malalignment, contractures, deformities, insufficient force generation, abnormal

muscle tone, abnormal tissue extensibility, exaggerated or hyperactive reflexes, poor selective control and regulation of activity of muscle groups, abnormal timing and decreased ability to learn unique movement².

Spastic diaplegia is the most prevalent form of cerebral palsy, with an incidence as high as 70% to 80% of the premature infants with CP³. The term diaplegia is used when the

legs are more affected than the arms. The severity and extent to which the arms are affected may vary from mild to moderate affections⁴. Spastic paraplegia is characterized by motor impairment, inco-ordination, primarily in the lower extremities that impairs many functional abilities, most notably ambulation⁵. It is clinically characterized by varying degrees of spasticity and /or inco-ordination in the lower extremities³. Lesion associated with spastic paraplegia may produce any or all of the following symptoms; increase muscle tone, loss of selective muscle control, deficient equilibrium reactions, and imbalance of muscle force across the joints, particularly in the lower extremities. Imbalance of muscle function in those children result in abnormal gait pattern described as crouch gait⁶.

For those patients whose motor skill may be absent, impaired, insufficient or abnormal, motor skill learning is considered one of the important aspects to be included in their rehabilitation program. Several attempts have been conducted to find out the conditions and variables of practice that may be used to facilitate, enhance and promote motor learning⁷. Practice is essential for motor learning and development of motor programs⁸. Well organized structured practice in related movement skill may improve performance in other desired skills⁹. Practice of one part of the body in performing a skilled act may increase performance of the same act in other part of the body. This phenomenon is called cross education or intermanual transfer¹⁰, bilateral transfer⁸, contra-lateral transfer of training¹¹.

A number of case reports and series reported benefits of mirror therapy in hemiparesis^{12,13,14}.

Although the majority of paraplegic children are able to ambulate, the acquisition of this skill is delayed and differs qualitatively from normal pediatric gait⁶.

This compromises major problems that hinder such children from practicing most of their daily living activities.

This study was conducted to determine the efficacy of facilitation of alternative upper limb movement, in addition to visual feedback, on improving lower limbs joints' angles during initial contact of gait in spastic paraplegic children.

Thirty spastic paraplegic cerebral palsied children from both sexes, ranging in age from 6 to 8 years (\bar{x} 7.60 \pm 0.39 years) represented the sample of the study.

They were selected from the outpatient clinic of the Faculty of Physical Therapy, Delta University for Science and Technology, according to the predetermined criteria as follows:

- All subjects were ambulant with an abnormal gait pattern (crouch gait), characterized by hip and knee flexion and ankle plantar flexion.
- The upper limbs were almost free with a degree of spasticity of the lower limbs ranged between grade 4 and 3 according to the Oswestry scale¹⁵.
- None of them had any fixed deformity or surgery of the lower extremities, however, few degrees of tightness was present in the Tendo-Achillis.
- They were able to understand and follow instructions given to them, with an IQ within normal average range.
- The subjects were divided randomly into two groups of equal number (A and B)

Instrumentation

For evaluation

Gait evaluation was done by using a clinical 3DMA 6.11 software version, video based gait assessment system (CLIMA) STT Systems Company, Spain which consists of

- 4 infrared cameras.
- Calibration tool
- Reflective Markers

- Large screen (24 inch)

For treatment

- Reciprocate unit:

An electronically braked bicycle ergometer, with its power read in watts till 50 pedal revolutions per minute. It is equipped with an electronic meter showing total pedal revolutions and time function.

- Mat, ball, roll wedge and tilting board.
- Table with adjustable height
- Chair with back support
- A large mirror placed in front of the child

Procedures

For evaluation

For capturing motion data in children some special modifications were made to the system configuration to improve visibility.

1. Camera System

The camera system was adapted to capture a much smaller person by lowering and moving it closer to the center of the capture volume.

The recommended measurements were:

- Camera height: 1.4 m.
- Length of the narrow side of the capture area: 3 m.
- Length of the wide side of the capture area: 3.6 m.

2. Calibration

-The calibration rod (8 point rod with 125 mm separation distance) was used.

3. Markers

These set contains 5 mm diameter reflective markers which were used for accurately detecting the child during the capture

1. Preparation

1- The first step before performing any capture, a calibration test was performed by placing a rod in vertical on the capture area (same where the motion was captured) and

without any other markers but the rod ones; as the results of the test were “Correct” we moved to the next step; where each child was asked to wear an easy fit, skin tight leg wear, then tracking tools (a simplified reflective markers consisting of 15 markers) were placed- according to Helen Hayes for children protocol¹⁶ - at the following points on the child’s body:

Right and left anterior superior iliac spine, right and left lateral aspect of the thigh, right and left knee, right and left calf ,right and left ankle, right and left 2nd toe, 2nd sacral vertebra , right and left heel.

2. Measurement

The capture process was then done with the child being instructed to walk with a straight back without turning

For treatment:

Group A received physical therapy program, for 60 minutes, including;

Improving posture control and balance, facilitation of righting, equilibrium and protective reactions, sustained stretching for the spastic muscles (Iliopsoas, Hamstrings and calf muscles), closed and open loops gait training exercises using parallel bars, obstacles, wedges, rolls and wooden stairs, arm pedaling on bicycle with unloaded cycling for three times as warming up and then with an intensity of thirty ramps/minute, for five minutes, repeated for three times, with five minutes rest in between.

Group B received the same program given to group A with applying the program in front of a large mirror to produce visual feedback.

Results

The raw data of the joints’ angles of the affected lower limbs were statistically treated to determine the mean and standard deviation of each measuring variable, for the two groups, before and after three months of treatment, t-test was then applied to examine

the significance of treatment conducted for each group.

The obtained results revealed no significant differences when comparing the pre-treatment mean values of the two groups. Significant improvement was observed in all the measuring variables of the two groups A and B, when comparing their pre and post treatment mean values. High significant improvement was observed in group B, when comparing its post treatment mean values with the post treatment mean values of group A.

Joint's angles during initial contact

For group A

As shown in table (1) and figure (1), the mean values of hip, knee and ankle joints' angles pre and post treatment were as follows:

Hip joints

The mean values of hip flexion angle pre and post treatment showed significant improvement. The mean value of right hip flexion angle pre -treatment was $42.79 \pm 3.39^\circ$ and became $39.27 \pm 2.99^\circ$ post treatment ($P < 0.01$). While the mean values of the left hip

flexion angles pre and post treatment were $43.11 \pm 3.26^\circ$ and $40 \pm 3.62^\circ$ respectively ($P < 0.05$).

Knee joints

The mean values of knee joints flexion angle pre and post treatment showed significant improvement. The mean value of right knee joint flexion angle pre and post treatment were $35.02 \pm 3.21^\circ$ and $32.3 \pm 3.07^\circ$ respectively ($P < 0.05$). While the mean values of the left knee joint flexion angles pre and post treatment were $35.77 \pm 3.34^\circ$ and $32 \pm 2.98^\circ$ respectively ($P < 0.01$).

Ankle joints

The mean values of ankle joints planter flexion angle pre and post treatment showed significant improvement. The mean value of right ankle joints planter flexion angle pre and post treatment were $30.12 \pm 3.01^\circ$ and $27.8 \pm 2.55^\circ$ respectively ($P < 0.05$). While the mean values of the left ankle joints planter flexion angles pre and post treatment were $30.65 \pm 3.04^\circ$ and $26.9 \pm 2.23^\circ$ respectively ($P < 0.01$).

Table 1: Pre and post treatment mean values of lower limbs joints' angles (hip, knee and ankle) in degrees for group A

Joints	X± SD	t-test	P-value	Significance
Hip joints				
Rt. side				
Pre treatment	$42.79 \pm 3.39^\circ$	3.016	< 0.01	Sig.
Post treatment	$39.27 \pm 2.99^\circ$			
Lt. side				
Pre treatment	$43.11 \pm 3.26^\circ$	2.47	< 0.05	Sig.
Post treatment	$40 \pm 3.62^\circ$			
Knee joints				

Rt. side				
Pre treatment	35.02±3.21°			
Post treatment	32.3±3.07°	2.37	< 0.05	Sig.
Lt. side				
Pre treatment	35.77±3.34°			
Post treatment	32±2.98°	3.184	< 0.01	Sig.
Ankle joints				
Rt. side				
Pre treatment	30.12±3.01°			
Post treatment	27.8±2.55°	2.278	< 0.05	Sig.
Lt. side				
Pre treatment	30.65±3.04°			
Post treatment	26.9±2.23°	3.85	< 0.01	Sig.

For group B:

As shown in table (2), the mean values of hip, knee and ankle joints' angles pre and post treatment were as follows:

Hip joints:

The mean values of hip flexion angle pre and post treatment showed significant improvement. The mean value of right hip flexion angle pre- treatment was 43.62±3.05° and became 35.05±2.44° post treatment (P< 0.0001). While the mean values of the left hip flexion angles pre and post treatment were 43.32±3.01° and 36.09±2.88° respectively (P< 0.0001).

Knee joints:

The mean values of knee joints flexion angle pre and post treatment showed significant improvement. The mean value of right knee

joint flexion angle pre and post treatment were 36.24±2.66° and 31.04±2.52° respectively (P< 0.0001). While the mean values of the left knee joint flexion angles pre and post treatment were 36.75±3.23° and 30.86±2.08° respectively (P< 0.0001).


Ankle joints:

The mean values of ankle joints planter flexion angle pre and post treatment showed significant improvement. The mean value of right ankle joints planter flexion angle pre and post treatment were 29.75±3.22° and 24.17±2.87° respectively (P< 0.0001). While the mean values of the left ankle joints planter flexion angles pre and post treatment were 29.9±3.14° and 25.0±2.63° respectively (P< 0.0001).

Table 2: Pre and post treatment mean values of lower limbs joints' angles (hip, knee and ankle) in degrees for group B

Joints	X± SD	t-test	P-value	Significance
Hip joints				
Rt. side Pre treatment Post treatment	42.79±3.39° 39.27±2.99°	3.016	< 0.01	Sig.
Lt. side Pre treatment Post treatment	43.11±3.26° 40±3.62°	2.47	< 0.05	Sig.
Knee joints				
Rt. side Pre treatment Post treatment	35.02±3.21° 32.3±3.07°	2.37	< 0.05	Sig.
Lt. side Pre treatment Post treatment	35.77±3.34° 32±2.98°	3.184	< 0.01	Sig.
Ankle joints				
Rt. side Pre treatment Post treatment	30.12±3.01° 27.8±2.55°	2.278	< 0.05	Sig.
Lt. side Pre treatment Post treatment	30.65±3.04° 26.9±2.23°	3.85	< 0.01	Sig.

Comparing the post treatment mean values of lower limbs' joints angles (hips, knees and ankles) in degrees for groups A and B, revealed significant improvement in favor of group B



Post treatment mean values of lower limbs joints' angles for groups A and B

Discussion

The gait of children with spastic diplegia is typically described as crouch gait, which is characterized by increased trunk flexion, hip flexion and ankle plantar flexion¹⁷. In addition to the obvious cosmetic disadvantages to this gait pattern, there is potential risk of increasing hip and ankle flexion contractures, which may ultimately result in loss of independent ambulation¹⁸. Thus, physical therapy interventions concerned with learning of this skill is important for those children.

This study was conducted to determine the effect of arm cycling, in addition to visual feedback, on controlling spasticity of the lower limbs in spastic diplegic children.

The results of the present study, after the suggested period of treatment revealed significant decrease in the range of measured angles in the two groups A and B, when comparing their pre and post treatment mean values.

High significant improvement was noticed in favor of group B, when comparing the post treatment mean values of the

measuring variables of groups A and B. However, no significant differences were observed when comparing the pre-treatment mean values of the two groups.

The pre-treatment results confirm the findings of Galli et al.,¹⁹ and Zhou et al.,²⁰, who reported that at initial contact, in diplegia, there is disturbance of the joints' angles, in the form of loss of dorsi- flexion of the ankle and generally reduced excursions.

O'Sullivan⁸ stated that the abnormal gait pattern seen in spastic diplegia results from the presence of spasticity and lack of practice. It may result from changes in the intrinsic properties of the muscle fibers themselves²¹. After the suggested period of treatment, it has been observed that the hip, knee flexion and also ankle plantar flexion angles improved. This post treatment improvement may be due to the effect of bicycle ergometer, in addition to the exercise program which helped in the motor learning process and in turn helped in the acquisition of motor control.

On basis of the literature review, little information is available regarding the role of cross education between the upper and lower extremities in spastic diplegic cerebral

palsied children. However, most of the studies were conducted to determine the effects of unilateral side training on the contralateral side in hemiplegic children.

Significant improvement observed in group B come in agreement with Stephen and Duncan²² who reported significant reduction in maximum force generated in ipsilateral and contralateral legs following continuous vibration at 30 Hz and 120Hz to the ipsilateral Quadriceps muscle. They attributed their findings to be due to the effect of activation of the central descending derived acting on the contralateral muscles. They established that cross education through inter limb transfer can be incorporated into treatment.

Accordingly, in diaplegic children, the uninvolved upper extremities can be engaged in practice to enhance the necessary motor programs, which can be used to control movements in the involved lower extremities.

The post treatment results agree with Farmer et al.,²³ who investigated neurophysiological evidence for a reorganization of central motor pathways in hemiplegic children. They reported that the motoneuron pools of homologous left and right hand muscles receive common synaptic input from abnormally branched pre-synaptic axon.

Weir et al.,²⁴ examined the effects of unilateral eccentric leg extension weight training and detraining on joint angle, cross education and bilateral deficit. The results indicated that, unilateral training resulted in increased bilateral strength, indicating cross training effect.

The results of the study agree with Wang et al.,²⁵ who established that there was a great influence of isometric knee flexion of the non-involved side on activation of the Rectus Femoris and Tibialis anterior muscles of the paralytic leg in stroke patients.

Andree and Maitra¹⁰ reported that an already learned task by one hand can positively influence learning and improve performance of the same task by the other hand, than activities comprising previously unlearned tasks.

The post treatment results confirm the findings of Helen and Daniel²⁶ who investigated neuromuscular recruitment during self-driven and externally driven lower limb motion. Healthy subjects exercised on a recumbent stepper using different combinations of upper and lower limb exertions. The recumbent stepper mechanically coupled the upper and lower limbs, allowing users to drive the stepping motion with upper and/or lower limbs. The subjects were instructed to step with: (1) active upper and lower limbs at an easy resistance level (active arms and legs), (2) active upper limbs and relaxed lower limbs at easy, medium and hard resistance levels (self-driven), (3) relaxed upper and lower limbs while another person drove the stepping motion. They recorded surface electromyography (EMG) from six lower limb muscles. The results revealed that self-driven EMG amplitudes were higher than externally driven EMG amplitudes. As resistance and upper limb exertion increased, self-driven EMG amplitudes also increased. According to the results of this study and literature review, it can be stated that improvement observed in the study group may be due to:

*Active upper limb movement increased neuromuscular activation of lower limbs during cyclic motions and enhanced activity dependent plasticity.

*Using visual feedback during active upper limb movement in addition to the exercise program increased neuromuscular activation of lower limbs and enhanced activity dependent plasticity.

*Self-assistance i.e. using upper limbs to assist the lower limbs could be beneficial due to inherent inter-limb neural coupling, in reflex response.

*Effect of activation of motor cortex during inter-limb movement and visual feedback.

*Interneuron connection between the upper and lower limbs and reorganization of motor output in the non-affected brain centers.

*Development of control over the tonic reflexes that are responsible about the distribution of muscle tone all over the body, through diffusion of motor impulses from practiced to unpracticed body parts through irradiation.

*The motor learning effect in skill acquisition and performance which is enhanced through practice and visual feedback.

*Modulation of abnormal muscle tone through improvement of sensory and motor awareness, enhancement of postural readjustment and motor learning process.

*Disruption of abnormal ratio between excitatory and inhibitory impulses from the afferent nerves in spastic cerebral palsy.

*The motor cortex sends its efferent corticospinal tracts that project ipsi- laterally as well as contra laterally.

Conclusion:

Cross education through inter limb transfer and Visual Feedback via mirroring can be incorporated into the physical therapy program to improve functional recovery (gait) through modulation of lower limb spasticity in Diaplegic Cerebral Palsy Children

References

- 1- Miller F. and Bachrach SJ. : Cerebral palsy; a complete guide for caregiving. 3rd ed. Baltimore: Johns Hopkins University Press, 2017; 3-5

- 2- Frank M. Chang, Jason T. Rhodes, Katherine M. Flynn and James J. Carollo: The role of gait analysis in treating gait abnormalities in cerebral palsy. *Orthopedic Management of Cerebral Palsy*, 2010 Volume 41, Issue 4, p441-612
- 3- Oskoui M., Coutinho F., Dykeman J., Jette N. and Pringsheim T.: An update on the prevalence of cerebral palsy: a systematic review and meta-analysis. *Dev. Med. Child Neurol.* 2013; 55; 509–519.
- 4- Lin CJ., Guo LY., Su FC., Chou YL., and Cherng RJ.: Common abnormal kinetic patterns of the knee in gait in spastic diplegia of cerebral palsy. *Gait Posture*. 2000 Jun.; 11(3):224–32.
- 5- Wren TA., Rethlefsen S. and Kay RM.: Prevalence of specific gait abnormalities in children with cerebral palsy: influence of cerebral palsy subtype, age, and previous surgery. *Journal of pediatric orthopedics*. 2005 Jan-Feb; 25(1):79–83.
- 6- Gunther M. and Blickhan R.: Joint stiffness of the ankle and the knee in running. *Journal of biomechanics*. 2002 Nov; 35 (11):1459–74.
- 7- Lark SD., Buckley JG., Bennett S., Jones D. and Sargeant AJ. : Joint torques and dynamic joint stiffness in elderly and young men during stepping down. *Clinical biomechanics*. 2003 Nov.; 18 (9):848–55.
- 8- O’Sullivan SB.: Strategies to improve motor control and motor learning. In. A. Davis Company, Philadelphia, 111, 132, 1995.
- 9- Muratori LM., Lamberg EM., Quinn L. and Duff SV.: Applying principles of motor learning and control to upper

- extremity rehabilitation. *J Hand Ther.* 2013; 26(2): 94–103.
- 10- Andree ME. and Maitra KK.: Intermanual transfer of a new writing occupation in young adults without disability, *Occupational Therapy International*, 9(1), 41–56, 2002.
- 11- Lara A. Green and David A. Gabriel: The effect of unilateral training on contralateral limb strength in young, older, and patient populations: a meta-analysis of cross education, *Physical Therapy Reviews*, 23:4-5, 238-249, 2018.
- 12- Thieme H, Morkisch N, Mehrholz J. et al.: Mirror therapy for improving motor function after stroke. *Cochrane Database Syst. Rev.* 2018; 7(7)
- 13- Tosi G, Romano D, Maravita A.: Mirror box training in hemiplegic stroke patients affects body representation. *Front Hum Neurosci.* 2018; 11: 617.
- 14- Suzanne S. Kuys, Tara Edwards, and Norman R. Morris : Effects and adherence of mirror therapy in people with chronic upper limb hemiparesis: A Preliminary Study, *ISRN Rehabilitation*, Volume 2012, Article ID 926784, 9 pages
- 15- Gaff B.: Grading of spasticity and its effect on voluntary movements. *Physiotherapy.* 62(11): 358-361, 1976.
- 16- Helen Hayes: A comparison of 3D gait models based on the Helen Hayes marker set Article, January 2000, <https://www.researchgate.net/publication/06253523>
- 17- Wei IP., Hsu WC., Chien HL., Chang CF., Liu YH., Ho TJ., et al.: Leg and joint stiffness in patients with bilateral medial knee osteoarthritis during level walking. *J Mech.* 2009 ; 25(3):279–87.
- 18- DeVita P and Hortobagyi T.: Age increases the skeletal versus muscular component of lower extremity stiffness during stepping down. *The Journals of gerontology Series A, Biological sciences and medical sciences.* 2000 Dec; 55 (12):B593–600.
- 19- Galli M, Cimolin V, Crivellini M, and Albertini G.: Gait analysis before and after gastrocnemius fascia lengthening in children with cerebral palsy. *Journal of Applied Biomaterials and Biomechanics.* 2005; 3(2):98–105.
- 20- Zhou J, Butler EE, and Rose J.: Neurologic correlates of gait abnormalities in cerebral palsy: Implications for treatment. *Front Hum Neurosci.* 2017; 11:103.
- 21- Thelen DD, Riewald SA, Asakawa DS, Sanger TD, and Delp SL.: Abnormal coupling of knee and hip moments during maximal exertions in persons with cerebral palsy. *Muscle and Nerve.* 2003 Apr; 27(4):486–93.
- 22- Stephen W. and Duncan LT.: Prolonged muscle vibration reduces maximal voluntary knee extension performance in both the ipsi-lateral and the contra-lateral limb in man. *European Journal of Applied Physiology*, 88(5)280-286, 2003
- 23- Farmer SF., Harrison LM., Ingram DA., and Stephens JA.: Plasticity of central motor pathways in children with hemiplegic cerebral palsy, *Neurology*, 41(9), 1505-10, 1991.
- 24- Weir JP., Harsh DJ., Housh TJ. And Weir LL: The effect of unilateral eccentric training and detraining on joint ankle specificity, cross training and the bilateral deficit. *J. Orthop. Sports Phys. Ther.* 22(5):207-15, 1995.

- 25- Wang W., Li K., Yue S., Yin C. and Wei N.: Associations between lower-limb muscle activation and knee flexion in post-stroke individuals: A study on the stance-to-swing phases of gait. PLoS One. 2017; 12(9):. Helen J. Huang and Daniel Ferris: Neural coupling between upper and lower limbs during recumbent stepping. Journal of Applied Physiology 97(4):1299-308, October 2004
- 26- Training and Detraining on Specificity, Cross-Training, The Effect of Unilateral Concentric Weight Joint Angle and the Bilateral Deficit) Joseph P. Weir, PhD ' Dona). Housh, Ph D Terry). Housh, Ph D loree 1. Weir, P~Journal of Orthopaedic & Sports Physical Therapy® Downloaded from www.jospt.org at on January 7, 2020. For personal use only. No other uses without permission. Copyright © 1997 Journal of Orthopaedic & Sports Physical Therapy®. All rights reserved.