

2019

The Caputo-Fabrizio Fractional Integral to Generate Some New Inequalities

Gustavo Asumu Mboro Nchama

Departamento de Ciencias Técnicas, Universidad Nacional de Guinea Ecuatorial(UNGE), Malabo, Guinea Ecuatorial. \\ Departamento de Matemática Aplicada, Universidad de La Habana, La Habana, Cuba., becquerr10@hotmail.com

Angela Leon Mecias

Departamento de Matemática Aplicada, Universidad de La Habana, La Habana, Cuba., becquerr10@hotmail.com

Mariano Rodriguez Richard

Departamento de Matemática Aplicada, Universidad de La Habana, La Habana, Cuba., becquerr10@hotmail.com

Follow this and additional works at: <https://digitalcommons.aaru.edu.jo/isl>

Recommended Citation

Asumu Mboro Nchama, Gustavo; Leon Mecias, Angela; and Rodriguez Richard, Mariano (2019) "The Caputo-Fabrizio Fractional Integral to Generate Some New Inequalities," *Information Sciences Letters*: Vol. 8 : Iss. 2 , Article 4.

Available at: <https://digitalcommons.aaru.edu.jo/isl/vol8/iss2/4>

This Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in Information Sciences Letters by an authorized editor. The journal is hosted on Digital Commons, an Elsevier platform. For more information, please contact rakan@aarj.edu.jo, marah@aarj.edu.jo, dr_ahmad@aarj.edu.jo.

The Caputo-Fabrizio Fractional Integral to Generate Some New Inequalities

Gustavo Asumu Mboro Nchama ^{1,2,*}, Ángela León Mecías ² and Mariano Rodríguez Richard ²

¹ Departamento de Ciencias Técnicas, Universidad Nacional de Guinea Ecuatorial(UNGE), Malabo, Guinea Ecuatorial.

² Departamento de Matemática Aplicada, Universidad de La Habana, La Habana, Cuba.

Received: 12 Jan. 2019, Revised: 22 Feb. 2019, Accepted: 12 Apr. 2019

Published online: 1 May 2019

Abstract: In this paper, we use the Caputo-Fabrizio (CF) fractional integral to establish some new integral inequalities in the case of functions with the same sense of variation.

Keywords: Caputo-Fabrizio fractional operators, Functions with the same sense of variation, Caputo-Fabrizio Fractional Integral Inequalities.

1 Introduction

The Fractional Calculus can be defined as a generalization of the integer-order differentiation. Its history goes back to seventeenth century, when in 1695 the derivative of order $\alpha = 1/2$ was described by Leibnitz in his letter to L'Hospital [25]. The Fractional Calculus has become more popular and useful due to its ability to describe some natural phenomena in numerous fields of engineering [21-24]. Moreover, it is well-known the importance of fractional calculus in obtaining fractional integral inequalities which are often used to prove the existence and uniqueness of fractional differential equations. In this line, there are several known forms of the fractional integrals which have been used to obtain fractional inequalities. The first is the Riemann-Liouville fractional integral [5]:

Theorem 1. Let f, g and h be positive and continuous functions on $[0, \infty)$, such that

$$(g(\tau) - g(\rho)) \left(\frac{f(\rho)}{h(\rho)} - \frac{f(\tau)}{h(\tau)} \right) \geq 0; \quad \tau, \rho \in [0, t], \quad t > 0.$$

Then we have

$$\frac{J^\alpha(f(t))}{J^\alpha(h(t))} \geq \frac{J^\alpha(gf(t))}{J^\alpha(gh(t))},$$

for all $\alpha > 0, t > 0$.

Theorem 2. Let f, g and h be positive and continuous functions on $[0, \infty)$, such that

$$(g(\tau) - g(\rho)) \left(\frac{f(\rho)}{h(\rho)} - \frac{f(\tau)}{h(\tau)} \right) \geq 0; \quad \tau, \rho \in [0, t], \quad t > 0.$$

Then for all $\alpha > 0, w, t > 0$, we have

$$\frac{J^\alpha(f(t)) \cdot J^w(gh(t)) + J^w(f(t)) \cdot J^\alpha(gh(t))}{J^\alpha(h(t)) \cdot J^w(gf(t)) + J^w(h(t)) \cdot J^\alpha(gf(t))} \geq 1.$$

Theorem 3. Let f and h be two positive continuous functions and $f \leq h$ on $[0, \infty)$. If $\frac{f}{h}$ is decreasing and f is increasing on $[0, \infty)$, then for any $p \geq 1, \alpha > 0, t > 0$, the inequality

$$\frac{J^\alpha(f(t))}{J^\alpha(h(t))} \geq \frac{J^\alpha(f^p(t))}{J^\alpha(h^p(t))},$$

is valid.

Theorem 4. Let f and h be two positive continuous functions and $f \leq h$ on $[0, \infty)$. If $\frac{f}{h}$ is decreasing and f is increasing on $[0, \infty)$, then for any $p \geq 1, \alpha > 0, w > 0, t > 0$, we have

$$\frac{J^\alpha(f(t)) \cdot J^w(h^p(t)) + J^w(f(t)) \cdot J^\alpha(h^p(t))}{J^\alpha(h(t)) \cdot J^w(f^p(t)) + J^w(h(t)) \cdot J^\alpha(f^p(t))} \geq 1.$$

The second is the Hadamard fractional integral [8, 20]:

* Corresponding author e-mail: becquerr10@hotmail.com



Theorem 5. Let f be an integrable function on $[1, \infty)$ such that there exist two integrable functions ϕ_1, ϕ_2 on $[1, \infty)$ with $\phi_1(t) \leq f(t) \leq \phi_2(t), \forall t \in [1, \infty)$. Then, for $t > 1, \alpha, \beta > 0$, one has

$$\begin{aligned} & {}^H I^\beta \phi_1(t) \cdot {}^H I^\alpha f(t) + {}^H I^\alpha \phi_2(t) \cdot {}^H I^\beta f(t) \\ & \geq {}^H I^\alpha \phi_2(t) \cdot {}^H I^\beta \phi_1(t) + {}^H I^\alpha f(t) \cdot {}^H I^\beta f(t). \end{aligned}$$

Theorem 6. Let f be an integrable function on $[1, \infty)$ and $p, q > 0$ satisfying $1/p + 1/q = 1$. Suppose that there exist two integrable functions ϕ_1, ϕ_2 on $[1, \infty)$ such that $\phi_1(t) \leq f(t) \leq \phi_2(t)$ for all $t \in [1, \infty)$. Then, for $t > 1, \alpha, \beta > 0$, one has

$$\begin{aligned} & \frac{1}{p} \cdot \frac{(\log t)^\beta}{\Gamma(\beta + 1)} \cdot {}^H I^\alpha ((\phi_2 - f)^p)(t) \\ & + \frac{1}{q} \cdot \frac{(\log t)^\alpha}{\Gamma(\alpha + 1)} \cdot {}^H I^\beta ((f - \phi_1)^q)(t) \\ & + {}^H I^\alpha \phi_2(t) \cdot {}^H I^\beta \phi_1(t) + {}^H I^\alpha f(t) \cdot {}^H I^\beta f(t) \\ & \geq {}^H I^\alpha \phi_2(t) \cdot {}^H I^\beta f(t) + {}^H I^\alpha f(t) \cdot {}^H I^\beta \phi_1(t). \end{aligned}$$

Theorem 7. Let f be an integrable function on $[1, \infty)$ and $p, q > 0$ satisfying $p + q = 1$. Suppose that there exist two integrable functions ϕ_1, ϕ_2 on $[1, \infty)$ such that $\phi_1(t) \leq f(t) \leq \phi_2(t)$ for all $t \in [1, \infty)$. Then, for $t > 1, \alpha, \beta > 0$, one has

$$\begin{aligned} & p \cdot \frac{(\log t)^\beta}{\Gamma(\beta + 1)} \cdot {}^H I^\alpha \phi_2(t) + q \cdot \frac{(\log t)^\alpha}{\Gamma(\alpha + 1)} \cdot {}^H I^\beta f(t) \\ & \geq {}^H I^\alpha (\phi_2 - f)^p(t) \cdot {}^H I^\beta (f - \phi_1)^q(t) \\ & + p \cdot \frac{(\log t)^\beta}{\Gamma(\beta + 1)} \cdot {}^H I^\alpha f(t) + q \cdot \frac{(\log t)^\alpha}{\Gamma(\alpha + 1)} \cdot {}^H I^\beta \phi_1(t). \end{aligned}$$

Another kind of fractional integral that appears in integral inequalities is the Saigo fractional integral [3, 4, 12, 16, ??]:

Theorem 8. Let $p \geq 1$ and let f, g be two positive functions on $[0, \infty)$ such that for all $x > 0, I_{0,x}^{\alpha,\beta,\eta}[f^p(x)] < \infty, I_{0,x}^{\alpha,\beta,\eta}[g^q(x)] < \infty$. If $0 < m \leq \frac{f(\tau)}{g(\tau)} \leq M < \infty, \tau \in (0, x)$ we have

$$\begin{aligned} & \left[I_{0,x}^{\alpha,\beta,\eta}[f^p(x)] \right]^{\frac{1}{p}} + \left[I_{0,x}^{\alpha,\beta,\eta}[g^q(x)] \right]^{\frac{1}{q}} \\ & \leq \frac{1 + M(m + 2)}{(m + 1)(M + 1)} \cdot \left[I_{0,x}^{\alpha,\beta,\eta}[(f + g)^p(x)] \right]^{\frac{1}{p}} \end{aligned}$$

for any $\alpha > \max\{0, -\beta\}, \beta < 1, \beta - 1 < \eta < 0$.

Theorem 9. Let $p > 1, \frac{1}{p} + \frac{1}{q} = 1$ and f, g be two positive functions on $[0, \infty)$ such that $I_{0,x}^{\alpha,\beta,\eta}[f(x)] < \infty, I_{0,x}^{\alpha,\beta,\eta}[g(x)] < \infty$. If $0 < m \leq \frac{f(\tau)}{g(\tau)} \leq M < \infty, \tau \in [0, x]$,

we have

$$\begin{aligned} & \left[I_{0,x}^{\alpha,\beta,\eta}[f(x)] \right]^{\frac{1}{p}} + \left[I_{0,x}^{\alpha,\beta,\eta}[g(x)] \right]^{\frac{1}{q}} \\ & \leq \left(\frac{M}{m} \right)^{\frac{1}{pq}} \cdot \left[I_{0,x}^{\alpha,\beta,\eta} \left[[f(x)]^{\frac{1}{p}} \cdot [g(x)]^{\frac{1}{q}} \right] \right], \end{aligned}$$

for all $x > 0, \alpha > \max\{0, -\beta\}, \beta < 1, \beta - 1 < \eta < 0$.

Theorem 10. Let f, g be two positive function on $[0, \infty)$ such that $I_{0,x}^{\alpha,\beta,\eta}[f^p(x)] < \infty, I_{0,x}^{\alpha,\beta,\eta}[g^q(x)] < \infty, x > 0$. If $0 < m \leq \frac{f(\tau)^p}{g(\tau)^q} \leq M < \infty, \tau \in [0, x]$. Then we have

$$\begin{aligned} & \left[I_{0,x}^{\alpha,\beta,\eta}[f^p(x)] \right]^{\frac{1}{p}} + \left[I_{0,x}^{\alpha,\beta,\eta}[g^q(x)] \right]^{\frac{1}{q}} \\ & \leq \left(\frac{M}{m} \right)^{\frac{1}{pq}} \cdot \left[I_{0,x}^{\alpha,\beta,\eta}[f(x) \cdot g(x)] \right], \end{aligned}$$

where $p > 1, \frac{1}{p} + \frac{1}{q} = 1$, for all $x > 0, \alpha > \max\{0, -\beta\}, \beta < 1, \beta - 1 < \eta < 0$.

Theorem 11. Let $f \geq 0, g \geq 0$ be two functions defined on $[0, \infty)$ such that g is non-decreasing. If

$$I_{0,x}^{\alpha,\beta,\eta} f(x) \geq I_{0,x}^{\alpha,\beta,\eta} g(x), \quad x > 0,$$

then for all $\alpha > \max\{0, -\beta\}, \beta < 1, \beta - 1 < \eta < 0, \gamma > 0, \delta > 0, \gamma - \delta > 0$, we have

$$I_{0,x}^{\alpha,\beta,\eta} f^{\gamma-\delta}(x) \leq I_{0,x}^{\alpha,\beta,\eta} f^\gamma(x) g^{-\delta}(x).$$

Theorem 12. Suppose that f, g and h be positive and continuous functions on $[0, \infty)$ such that

$$[g(\tau) - g(\rho)] \left(\frac{f(\rho)}{h(\rho)} - \frac{f(\tau)}{h(\tau)} \right) \geq 0; \quad \tau, \rho \in [0, x], x > 0.$$

Then for all $x > 0, \alpha > \max\{0, -\beta\}, \beta < 1, \beta - 1 < \eta < 0$, we have

$$\frac{I_{0,x}^{\alpha,\beta,\eta}[f(x)]}{I_{0,x}^{\alpha,\beta,\eta}[h(x)]} \geq \frac{I_{0,x}^{\alpha,\beta,\eta}[(gf)(x)]}{I_{0,x}^{\alpha,\beta,\eta}[(gh)(x)]}.$$

Theorem 13. Suppose that f, g and h be positive and continuous functions on $[0, \infty)$ such that

$$[g(\tau) - g(\rho)] \left(\frac{f(\rho)}{h(\rho)} - \frac{f(\tau)}{h(\tau)} \right) \geq 0; \quad \tau, \rho \in (0, t), t > 0.$$

Then for all $x > 0, \alpha > \max\{0, -\beta\}, \psi > \max\{0, -\phi\}, \beta < 1, \beta - 1 < \eta < 0, \phi < 1, \phi - 1 < \xi < 0$, we have

$$\frac{I_{0,x}^{\alpha,\beta,\eta}[f(x)] I_{0,x}^{\psi,\phi,\xi}[(gh)(x)] + I_{0,x}^{\psi,\phi,\xi}[f(x)] I_{0,x}^{\alpha,\beta,\eta}[(gh)(x)]}{I_{0,x}^{\alpha,\beta,\eta}[h(x)] I_{0,x}^{\psi,\phi,\xi}[(gf)(x)] + I_{0,x}^{\psi,\phi,\xi}[h(x)] I_{0,x}^{\alpha,\beta,\eta}[(gh)(x)]} \geq 1.$$

In 2015, Caputo and Fabrizio introduced a new fractional derivative [13]. The interest for this approach was due to the necessity of using a model describing structures with different scales [13]. In the literature there is no evidence of the use of the Caputo fractional derivative and Caputo-Fabrizio fractional integral to obtain fractional integral inequalities. The propose of this paper is to use the CF fractional integral to establish some new integral inequalities which have been obtained before by using the Riemann-Liouville, Hadamard and Saigo fractional operators [1, 5, 9]. The obtained Caputo-Fabrizio fractional inequalities could be helpful to prove the existence and uniqueness of some ordinary Caputo-Fabrizio fractional differential equations. The paper has been organized as follows, in Section 2, we define basic definitions related to fractional integrals. In Section 3, appears the main results. Finally, conclusion is summarized in Section 4.

2 Preliminaries

Here we give the following definitions:

Definition 1. Let $\alpha \in \mathbb{R}$ such that $0 < \alpha < 1$. The Caputo-Fabrizio fractional integral of order α of a function f is defined by [10]

$$I_{0,t}^\alpha f(t) = \frac{1}{\alpha} \int_0^t e^{-\frac{1-\alpha}{\alpha}(t-s)} f(s) ds.$$

Definition 2. Let $\alpha, a \in \mathbb{R}$ such that $0 < \alpha < 1$. The Caputo-Fabrizio fractional derivative of order α of a function f is defined by

$$D_{a,t}^\alpha f(t) = \frac{1}{1-\alpha} \int_a^t e^{-\frac{\alpha}{1-\alpha}(t-s)} f'(s) ds.$$

Definition 3. Let $\alpha > 0, \beta, \eta \in \mathbb{R}$, then the Saigo fractional integral $I_{0,t}^{\alpha,\beta,\eta} \{f(t)\}$ of order α for a real-valued continuous function $f(t)$ is defined by

$$I_{0,t}^{\alpha,\beta,\eta} \{f(t)\} = \frac{t^{-\alpha-\beta}}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} F_1(\alpha+\beta, -\eta; \alpha; 1-\frac{\tau}{t}) f(\tau) d\tau,$$

where the function $F_1(-)$ is the Gaussian hypergeometric function defined by

$$F_1(a, b; c; t) = \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n} \cdot \frac{t^n}{n!},$$

and $(a)_n$ is the Pochhammer symbol

$$(a)_n = a(a+1) \cdots (a+n-1), (a)_0 = 1.$$

Definition 4. The Hadamard fractional integral is defined by

$${}^H I^\alpha f(t) = \frac{1}{\Gamma(\alpha)} \int_1^t \left(\log \frac{t}{\tau}\right)^{\alpha-1} f(\tau) \frac{d\tau}{\tau}$$

for $Re(\alpha) > 0, t > 1$.

Definition 5. The Riemann-Liouville fractional integral is defined by

$${}^{RL} I_{0,t}^\alpha f(t) = \frac{1}{\Gamma(\alpha)} \int_0^t (t-\tau)^{\alpha-1} f(\tau) d\tau.$$

Definition 6. We say that two functions f and g have the same sense of variation (synchronous) on $[0, \infty)$ if [16]

$$(f(\tau) - f(\rho))(g(\tau) - g(\rho)) \geq 0 \quad \tau, \rho \in (0, t), t > 0.$$

For more details, see [1, 6, 7, 8, 11-15, 17-20].

3 Main Results

Here, we give some interesting inequalities concerning the Caputo-Fabrizio fractional integral.

Theorem 14. Let p be a positive function on $[0, \infty)$ and let f and g be two differentiable functions having the same sense of variation on $[0, \infty)$. If $f', g' \in L_\infty([0, \infty))$, then for all $t > 0, \alpha \in (0, 1)$, we have

$$0 \leq I_{0,t}^\alpha p(t) I_{0,t}^\alpha p f g(t) - I_{0,t}^\alpha p f(t) \cdot I_{0,t}^\alpha p g(t) \leq \|f'\|_{L_\infty} \cdot \|g'\|_{L_\infty} \cdot [I_{0,t}^\alpha p(t) \cdot I_{0,t}^\alpha t^2 p(t) - (I_{0,t}^\alpha t p(t))^2]. \quad (1)$$

Proof. Let f and g be two functions satisfying the conditions of theorem 14 and let p be a positive function on $[0, \infty)$. Define

$$H(\tau, \rho) := (f(\tau) - f(\rho))(g(\tau) - g(\rho)) \quad (2)$$

for all $\tau, \rho \in (0, t)$. We have

$$H(\tau, \rho) \geq 0. \quad (3)$$

Thanks to (3), we have

$$\begin{aligned} & \frac{1}{2\alpha^2} \int_0^t \int_0^t e^{-\frac{1-\alpha}{\alpha}(t-\tau)} \cdot e^{-\frac{1-\alpha}{\alpha}(t-\rho)} \cdot p(\tau)p(\rho)H(\tau, \rho) d\tau d\rho \\ & = I_{0,t}^\alpha p(t) \cdot I_{0,t}^\alpha p f g(t) - I_{0,t}^\alpha p f(t) \cdot I_{0,t}^\alpha p g(t) \geq 0. \end{aligned} \quad (4)$$

The relation (3) can be written as follows:

$$H(\tau, \rho) = \int_\tau^\rho \int_\tau^y f'(y)g'(z) dy dz. \quad (5)$$



Hence, we can write

$$\begin{aligned}
 H(\tau, \rho) &\leq \left| \int_{\tau}^{\rho} f'(y)dy \right| \left| \int_{\tau}^{\rho} g'(z)dz \right| \\
 &\leq \|f'\|_{L^{\infty}} \cdot \|g'\|_{L^{\infty}} \cdot (\tau - \rho)^2.
 \end{aligned} \tag{6}$$

Therefore

$$\begin{aligned}
 &\frac{1}{2\alpha^2} \int_0^t \int_0^t e^{-\frac{1-\alpha}{\alpha}(t-\tau)} \cdot e^{-\frac{1-\alpha}{\alpha}(t-\rho)} \cdot p(\tau)p(\rho)H(\tau, \rho)d\tau d\rho \\
 &\leq \frac{\|f'\|_{L^{\infty}} \|g'\|_{L^{\infty}}}{2\alpha^2} \int_0^t \int_0^t e^{-\frac{1-\alpha}{\alpha}(t-\tau)} e^{-\frac{1-\alpha}{\alpha}(t-\rho)} \cdot (\tau^2 - 2\tau\rho + \rho^2)p(\tau)p(\rho)d\tau d\rho.
 \end{aligned} \tag{7}$$

Consequently

$$\begin{aligned}
 &\frac{1}{2\alpha^2} \int_0^t \int_0^t e^{-\frac{1-\alpha}{\alpha}(t-\tau)} \cdot e^{-\frac{1-\alpha}{\alpha}(t-\rho)} \cdot p(\tau)p(\rho)H(\tau, \rho)d\tau d\rho \\
 &\leq \|f'\|_{L^{\infty}} \cdot \|g'\|_{L^{\infty}} \cdot [I_{0^+}^{\alpha}p(t) \cdot I_{0^+}^{\alpha}t^2p(t) - (I_{0^+}^{\alpha}tp(t))^2].
 \end{aligned} \tag{8}$$

According to (4) and (8), we get (1), and thus theorem 14 is proved.

Now, we are ready to generalize Theorem 14 to the following theorem:

Theorem 15. Let p be a positive function on $[0, \infty)$ and let f and g be two differentiable functions having the same sense of variation on $[0, \infty)$. If $f', g' \in L^{\infty}([0, \infty))$, then for all $t > 0, \alpha, \beta \in (0, 1)$, we have

$$\begin{aligned}
 0 &\leq I_{0^+}^{\alpha}pf g(t) \cdot I_{0^+}^{\beta}p(t) + I_{0^+}^{\alpha}p(t) \cdot I_{0^+}^{\beta}p f g(t) \\
 &\quad - I_{0^+}^{\alpha}p f(t) \cdot I_{0^+}^{\beta}p g(t) - I_{0^+}^{\alpha}p g(t) \cdot I_{0^+}^{\beta}p f(t) \\
 &\leq \|f'\|_{L^{\infty}} \cdot \|g'\|_{L^{\infty}} \{ I_{0^+}^{\alpha}t^2p(t) \cdot I_{0^+}^{\beta}p(t) \\
 &\quad + I_{0^+}^{\alpha}p(t) \cdot I_{0^+}^{\beta}t^2p(t) - 2I_{0^+}^{\alpha}tp(t) \cdot I_{0^+}^{\beta}tp(t) \}
 \end{aligned} \tag{9}$$

Proof. It is easy to see that

$$\begin{aligned}
 &\frac{1}{1-\alpha} \cdot \frac{1}{1-\beta} \int_0^t \int_0^t e^{-\frac{\alpha}{1-\alpha}(t-\tau)} \cdot e^{-\frac{\beta}{1-\beta}(t-\rho)} p(\tau)p(\rho)H(\tau, \rho)d\tau d\rho \\
 &= I_{0^+}^{\alpha}p f g(t) \cdot I_{0^+}^{\beta}p(t) + I_{0^+}^{\alpha}p(t) \cdot I_{0^+}^{\beta}p f g(t) \\
 &\quad - I_{0^+}^{\alpha}p f(t) \cdot I_{0^+}^{\beta}p g(t) \\
 &\quad - I_{0^+}^{\alpha}p g(t) \cdot I_{0^+}^{\beta}p f(t) \geq 0.
 \end{aligned} \tag{10}$$

From the relation (6), we obtain the following estimate

$$\begin{aligned}
 &\frac{1}{1-\alpha} \frac{1}{1-\beta} \int_0^t \int_0^t e^{-\frac{\alpha}{1-\alpha}(t-\tau)} \cdot e^{-\frac{\beta}{1-\beta}(t-\rho)} \cdot p(\tau)p(\rho)H(\tau, \rho)d\tau d\rho \\
 &\leq \|f'\|_{L^{\infty}} \cdot \|g'\|_{L^{\infty}} [I_{0^+}^{\alpha}t^2p(t) \cdot I_{0^+}^{\beta}p(t) \\
 &\quad + I_{0^+}^{\alpha}p(t) \cdot I_{0^+}^{\beta}t^2p(t) - 2I_{0^+}^{\alpha}tp(t) \cdot I_{0^+}^{\beta}tp(t)].
 \end{aligned} \tag{11}$$

Combining (10) with (11), inequality (9) follows.

Theorem 16. Let f, g and h be positive and continuous functions on $[0, \infty)$, such that

$$(g(\tau) - g(\rho)) \left(\frac{f(\rho)}{h(\rho)} - \frac{f(\tau)}{h(\tau)} \right) \geq 0; \quad \tau, \rho \in [0, t], t > 0, \tag{12}$$

then we have

$$\frac{I_{0^+}^{\alpha}f(t)}{I_{0^+}^{\alpha}h(t)} \geq \frac{I_{0^+}^{\alpha}g f(t)}{I_{0^+}^{\alpha}g h(t)} \tag{13}$$

for any $\alpha \in (0, 1), t > 0$.

Proof. Suppose that f, g and h are positive and continuous functions on $[0, \infty)$. Using (12), we can write

$$\begin{aligned}
 &g(\tau) \cdot \frac{f(\rho)}{h(\rho)} + g(\rho) \cdot \frac{f(\tau)}{h(\tau)} \\
 &\quad - g(\rho) \cdot \frac{f(\rho)}{h(\rho)} - g(\tau) \cdot \frac{f(\tau)}{h(\tau)} \geq 0
 \end{aligned}$$

for all $\tau, \rho \in [0, t], t > 0$. That is

$$\begin{aligned}
 &g(\tau)f(\rho)h(\tau) + g(\rho)f(\tau)h(\rho) \\
 &\quad - g(\rho)f(\rho)h(\tau) - g(\tau)f(\tau)h(\rho) \geq 0
 \end{aligned} \tag{14}$$

for all $\tau, \rho \in [0, t], t > 0$.

Now, multiplying both sides of (14) by $\frac{1}{\alpha^2} \cdot e^{-\frac{1-\alpha}{\alpha}(t-\tau)} \cdot e^{-\frac{1-\alpha}{\alpha}(t-\rho)}$, then integrating the resulting inequality with respect to τ and ρ over $(0, t) \times (0, t)$, we get

$$I_{0^+}^{\alpha}gh(t) \cdot I_{0^+}^{\alpha}f(t) - I_{0^+}^{\alpha}gf(t) \cdot I_{0^+}^{\alpha}h(t) \geq 0 \tag{15}$$

From (15), inequality (13) follows.

Theorem 17. Let f, g and h be positive and continuous functions on $[0, \infty)$, such that

$$(g(\tau) - g(\rho)) \left(\frac{f(\rho)}{h(\rho)} - \frac{f(\tau)}{h(\tau)} \right) \geq 0; \quad \tau, \rho \in [0, t], t > 0, \tag{16}$$

then for all $\alpha, \beta \in (0, 1), t > 0$, we have

$$\frac{I_{0^+}^{\alpha}gh(t) \cdot I_{0^+}^{\beta}f(t) + I_{0^+}^{\beta}gh(t) \cdot I_{0^+}^{\alpha}f(t)}{I_{0^+}^{\alpha}h(t) \cdot I_{0^+}^{\beta}g f(t) + I_{0^+}^{\alpha}g f(t) \cdot I_{0^+}^{\beta}h(t)} \geq 1 \tag{17}$$

Proof. Suppose that f, g and h are positive and continuous functions on $[0, \infty)$. Now, multiplying both sides of (14) by $\frac{1}{\alpha\beta} \cdot e^{-\frac{1-\alpha}{\alpha}(t-\tau)} \cdot e^{-\frac{1-\beta}{\beta}(t-\rho)}$, then integrating the resulting inequality with respect to τ and ρ over $(0, t) \times (0, t)$, we get

$$\begin{aligned}
 &I_{0^+}^{\alpha}gh(t) \cdot I_{0^+}^{\beta}f(t) + I_{0^+}^{\beta}gh(t) \cdot I_{0^+}^{\alpha}f(t) \\
 &\geq I_{0^+}^{\alpha}h(t) \cdot I_{0^+}^{\beta}g f(t) + I_{0^+}^{\alpha}g f(t) \cdot I_{0^+}^{\beta}h(t).
 \end{aligned}$$

Hence, we obtain (17).

Note 1. It is obviously to see that (16) holds true for f, g, h such that either

(i) g is increasing and $\frac{f}{h}$ is decreasing

or

(ii) g is decreasing and $\frac{f}{h}$ is increasing.

Theorem 18. Let f and h be two positive continuous functions and $f \leq h$ on $[0, \infty)$. If $\frac{f}{h}$ is decreasing and f is increasing on $[0, \infty)$, then for any $p \geq 1, \alpha \in (0, 1), t > 0$, the inequality

$$\frac{I_{0^+}^\alpha f(t)}{I_{0^+}^\alpha h(t)} \geq \frac{I_{0^+}^\alpha f^p(t)}{I_{0^+}^\alpha h^p(t)}, \quad (18)$$

is valid.

Proof. Since $p \geq 1$ and f is increasing, then $g := f^{p-1}$ is also increasing. Then by applying the theorem 16, we get

$$\frac{I_{0^+}^\alpha f(t)}{I_{0^+}^\alpha h(t)} \geq \frac{I_{0^+}^\alpha (f \cdot f^{p-1}(t))}{I_{0^+}^\alpha (h \cdot f^{p-1}(t))}, \quad (19)$$

now, since $f \leq h$ on $(0, \infty]$, then

$$\frac{1}{\alpha} \cdot e^{-\frac{1-\alpha}{\alpha}(t-\tau)} \cdot h(\tau) \cdot f^{p-1}(\tau) \leq \frac{1}{\alpha} \cdot e^{-\frac{1-\alpha}{\alpha}(t-\tau)} \cdot h^p(\tau) \quad (20)$$

for all $\tau \in [0, t], t > 0$. Integrating both sides of (20) with respect to τ over $(0, t)$, yields

$$I_{0^+}^\alpha (h f^{p-1}(t)) \leq I_{0^+}^\alpha (h^p(t)). \quad (21)$$

Consequently

$$\frac{I_{0^+}^\alpha (f \cdot f^{p-1}(t))}{I_{0^+}^\alpha (h \cdot f^{p-1}(t))} = \frac{I_{0^+}^\alpha f^p(t)}{I_{0^+}^\alpha (h \cdot f^{p-1}(t))} \geq \frac{I_{0^+}^\alpha f^p(t)}{I_{0^+}^\alpha h^p(t)}. \quad (22)$$

Combining (19) with (22), we obtain (18).

Theorem 19. Let f and h be two positive and continuous functions and $f \leq h$ on $[0, \infty)$. If $\frac{f}{h}$ is decreasing and f is increasing on $[0, \infty)$, then for any $p \geq 1, \alpha, \beta \in (0, 1), t > 0$, we have

$$\frac{I_{0^+}^\beta f(t) \cdot I_{0^+}^\alpha h^p(t) + I_{0^+}^\alpha f(t) \cdot I_{0^+}^\beta h^p(t)}{I_{0^+}^\alpha h(t) \cdot I_{0^+}^\beta f^p(t) + I_{0^+}^\beta h(t) \cdot I_{0^+}^\alpha f^p(t)} \geq 1. \quad (23)$$

Proof. Taking $g := f^{p-1}$, then by theorem 17, yields

$$\frac{I_{0^+}^\alpha [h \cdot f^{p-1}(t)] \cdot I_{0^+}^\beta f(t) + I_{0^+}^\alpha f(t) \cdot I_{0^+}^\beta [h \cdot f^{p-1}(t)]}{I_{0^+}^\alpha h(t) \cdot I_{0^+}^\beta f^p(t) + I_{0^+}^\beta h(t) \cdot I_{0^+}^\alpha f^p(t)} \geq 1. \quad (24)$$

Using the fact that $f \leq h$ on $[0, \infty)$, we can write

$$\frac{1}{\beta} e^{-\frac{1-\beta}{\beta}(t-\rho)} \cdot h \cdot f^{p-1}(\rho) \leq \frac{1}{\beta} \cdot e^{-\frac{1-\beta}{\beta}(t-\rho)} \cdot h^p(\rho) \quad (25)$$

for all $\rho \in [0, t], t > 0$. Integrating both sides of (25) with respect to ρ over $(0, t)$, we obtain

$$I_{0^+}^\beta h f^{p-1}(t) \leq I_{0^+}^\beta h^p(t). \quad (26)$$

Multiplying (21) by $I_{0^+}^\beta f(t)$ and (26) by $I_{0^+}^\alpha f(t)$, we can write

$$\begin{aligned} & I_{0^+}^\beta f(t) \cdot I_{0^+}^\alpha (h f^{p-1}(t)) + I_{0^+}^\alpha f(t) \cdot I_{0^+}^\beta (h f^{p-1}(t)) \\ & \leq I_{0^+}^\beta f(t) \cdot I_{0^+}^\alpha (h^p(t)) + I_{0^+}^\alpha f(t) \cdot I_{0^+}^\beta (h^p(t)). \end{aligned} \quad (27)$$

Now, using (24) and (27), we deduce (23).

Theorem 20. Let $\alpha \in (0, 1), p \geq 1$ and f, g be two positive continuous functions on $[0, \infty)$. If $0 < m \leq \frac{f(\tau)}{g(\tau)} \leq M, \tau \in (0, t)$, then we have

$$\begin{aligned} & [I_{0^+}^\alpha f^p(t)]^{1/p} + [I_{0^+}^\alpha g^p(t)]^{1/p} \\ & \leq \frac{M(m+2)+1}{(M+1)(m+1)} [I_{0^+}^\alpha (f+g)^p(t)]^{1/p}. \end{aligned} \quad (28)$$

Proof. Using the condition $\frac{f(\tau)}{g(\tau)} \leq M, \tau \in (0, t), t > 0$, we can write

$$(M+1)^p \cdot f^p(\tau) \leq M^p \cdot (f+g)^p(\tau). \quad (29)$$

Multiplying both sides of (29) by $\frac{1}{\alpha} \cdot e^{-\frac{1-\alpha}{\alpha}(t-\tau)}$, then integrating resulting identity with respect to τ from 0 to x , we get

$$(M+1)^p \cdot I_{0^+}^\alpha f^p(t) \leq M^p \cdot I_{0^+}^\alpha (f+g)^p(t)$$

Hence, we can write

$$[I_{0^+}^\alpha f^p(t)]^{1/p} \leq \frac{M}{M+1} \cdot [I_{0^+}^\alpha (f+g)^p(t)]^{1/p}, \quad (30)$$

on the other hand, using condition $m \leq \frac{f(\tau)}{g(\tau)}$, we obtain

$$\left(1 + \frac{1}{m}\right) g(\tau) \leq \frac{1}{m} (f(\tau) + g(\tau)),$$

therefore

$$\left(1 + \frac{1}{m}\right)^p g^p(\tau) \leq \left(\frac{1}{m}\right)^p (f(\tau) + g(\tau))^p. \quad (31)$$

Now, multiplying both sides of (31) by $\frac{1}{\alpha} \cdot e^{-\frac{1-\alpha}{\alpha}(t-\tau)}$, then integrating resulting identity with respect to τ from 0 to t , we have

$$[I_{0^+}^\alpha g^p(t)]^{1/p} \leq \frac{1}{m+1} \cdot [I_{0^+}^\alpha (f+g)^p(t)]^{1/p}. \quad (32)$$

The inequality (28) follows on adding the inequalities (30) and (32)

Theorem 21. Let $\alpha \in (0, 1), p > 1, \frac{1}{p} + \frac{1}{q} = 1$ and f, g be two positive and continuous functions on $[0, \infty)$. If

$$0 < m \leq \frac{f(\tau)}{g(\tau)} \leq M < \infty, \quad \tau \in [0, t], \quad (33)$$



then the inequality

$$\begin{aligned}
 & [I_{0r}^\alpha f(t)]^{1/p} \cdot [I_{0r}^\alpha g(t)]^{1/q} \\
 & \leq \left(\frac{M}{m}\right)^{1/pq} \cdot I_{0r}^\alpha [f^{1/p}(t) \cdot g^{1/q}(t)]
 \end{aligned} \tag{34}$$

holds.

Proof. Since $\frac{f(\tau)}{g(\tau)} \leq M, \tau \in [0, t], t > 0$, therefore

$$[g(\tau)]^{1/q} \geq M^{-1/q} \cdot [f(\tau)]^{1/q},$$

and so,

$$\begin{aligned}
 [f(\tau)]^{1/q} \cdot [g(\tau)]^{1/q} & \geq M^{-1/q} \cdot [f(\tau)]^{1/q} \cdot [f(\tau)]^{1/q} \\
 & = M^{-1/q} \cdot f(\tau).
 \end{aligned} \tag{35}$$

Multiplying both sides of (35) by $\frac{1}{\alpha} \cdot e^{-\frac{1-\alpha}{\alpha}(t-\tau)}$, then integrating resulting identity with respect to τ from 0 to t , we have

$$I_{0r}^\alpha [f^{1/p}(t) \cdot g^{1/q}(t)] \geq M^{-1/q} \cdot I_{0r}^\alpha f(t),$$

and consequently

$$(I_{0r}^\alpha [f^{1/p}(t) \cdot g^{1/q}(t)])^{1/p} \geq M^{-1/pq} \cdot (I_{0r}^\alpha f(t))^{1/p}. \tag{36}$$

On the other hand, since $mg(\tau) \leq f(\tau), \tau \in [0, t], t > 0$, then we have

$$[f(\tau)]^{1/p} \geq m^{1/p} \cdot [g(\tau)]^{1/p},$$

and so,

$$\begin{aligned}
 [g(\tau)]^{1/q} \cdot [f(\tau)]^{1/p} & \geq m^{1/p} \cdot [g(\tau)]^{1/p} \cdot [g(\tau)]^{1/q} \\
 & = m^{1/p} \cdot g(\tau).
 \end{aligned} \tag{37}$$

Now, multiplying both sides of (37) by $\frac{1}{\alpha} \cdot e^{-\frac{1-\alpha}{\alpha}(t-\tau)}$, then integrating resulting identity with respect to τ over $(0, t)$, we obtain

$$I_{0r}^\alpha [g^{1/q}(t) \cdot f^{1/p}(t)] \geq m^{1/p} \cdot I_{0r}^\alpha g(t).$$

Hence, it follows

$$(I_{0r}^\alpha [g^{1/q}(t) \cdot f^{1/p}(t)])^{1/q} \geq m^{1/pq} \cdot (I_{0r}^\alpha g(t))^{1/q}. \tag{38}$$

Thanks to (36) and (38), we obtain (34).

Theorem 22. Let $0 < \alpha < 1, p > 1, \frac{1}{p} + \frac{1}{q} = 1$, f and g be two positive and continuous functions on $[0, \infty)$. If

$$0 < m \leq \frac{f^p(\tau)}{g^q(\tau)} \leq M < \infty, \quad \tau \in [0, t],$$

then we have

$$[I_{0r}^\alpha f^p(t)]^{1/p} \cdot [I_{0r}^\alpha g^q(t)]^{1/q} \leq \left(\frac{M}{m}\right)^{1/pq} \cdot I_{0r}^\alpha [f(t) \cdot g(t)]. \tag{39}$$

Proof. Replacing $f(\tau)$ and $g(\tau)$ respectively by $(f(\tau))^p$ and $(g(\tau))^q, \tau \in [0, t], t > 0$ in theorem 21, we obtain (39).

Theorem 23. Let $0 < \alpha < 1, p > 1, \frac{1}{p} + \frac{1}{q} = 1$, f and g be two positive and continuous functions on $[0, \infty)$. If

$$0 < m \leq \frac{f(\tau)}{g(\tau)} \leq M < \infty, \tag{40}$$

then we have

$$I_{0r}^\alpha \left(\frac{f^p(t)}{g^{p/q}(t)} \right) \leq \left(\frac{M}{m}\right)^{1/q} \cdot \frac{(I_{0r}^\alpha f(t))^p}{(I_{0r}^\alpha g(t))^{p/q}}. \tag{41}$$

Proof. Using theorem 21, we obtain

$$\begin{aligned}
 I_{0r}^\alpha f(t) & = I_{0r}^\alpha \left[\left(\frac{f^p(t)}{g^{p/q}(t)} \right)^{1/p} \cdot (g(t))^{1/q} \right] \\
 & \geq \left(\frac{m}{M}\right)^{1/pq} \cdot \left[I_{0r}^\alpha \frac{f^p(t)}{g^{p/q}(t)} \right]^{1/p} \cdot [I_{0r}^\alpha g(t)]^{1/q}.
 \end{aligned}$$

Hence, we can write

$$[I_{0r}^\alpha f(t)]^p \geq \left(\frac{m}{M}\right)^{1/q} \cdot \left[I_{0r}^\alpha \frac{f^p(t)}{g^{p/q}(t)} \right] [I_{0r}^\alpha g(t)]^{p/q}. \tag{42}$$

Thanks to (42), we obtain (41).

Theorem 24. Let $0 < \alpha < 1, p > 1, \frac{1}{p} + \frac{1}{q} = 1$ and f be a positive and continuous function on $[0, \infty)$. If $0 < m \leq f(\tau) \leq M < \infty$ and

$$I_{0r}^\alpha f(t) \geq \left(\frac{1}{1-\alpha} \left[1 - e^{-\frac{1-\alpha}{\alpha}t} \right] \right)^{-p/q}, \tag{43}$$

then the inequality

$$I_{0r}^\alpha [f^p(t)] \leq \left(\frac{M}{m}\right)^{1/q} \cdot (I_{0r}^\alpha f(t))^{p+1}$$

holds.

Proof. Using theorem 23 and condition (43), we obtain

$$\begin{aligned}
 & I_{0r}^\alpha [f^p(t)] \\
 & = I_{0r}^\alpha \left[\frac{f^p(t)}{(1)^{p/q}} \right] \\
 & \leq \left(\frac{M}{m}\right)^{1/q} \cdot \frac{(I_{0r}^\alpha f(t))^p}{(I_{0r}^\alpha 1)^{p/q}} \\
 & = \left(\frac{M}{m}\right)^{1/q} \cdot \left(\frac{1}{1-\alpha} \left[1 - e^{-\frac{1-\alpha}{\alpha}t} \right] \right)^{-p/q} \cdot (I_{0r}^\alpha f(t))^p \\
 & \leq \left(\frac{M}{m}\right)^{1/q} \cdot (I_{0r}^\alpha f(t))^{p+1}
 \end{aligned}$$

as required.

Remark. For any $p > 1$, with f a positive and continuous function, we note that

$$0 < m \leq f^p(\tau) \leq M < \infty \Leftrightarrow 0 < m^{1/p} \leq f(\tau) \leq M^{1/p} < \infty. \tag{44}$$

Theorem 25. Let $0 < \alpha < 1, p > 1$ with $\frac{1}{p} + \frac{1}{q} = 1$, and f be a positive and continuous function on $[0, \infty)$. If

$$0 < m \leq f^p(\tau) \leq M < \infty, \quad \tau \in [0, t],$$

then

$$\begin{aligned} & [I_{0r}^\alpha f^p(t)]^{1/p} \\ & \leq \left(\frac{M}{m}\right)^{2/pq} \left(\frac{1}{1-\alpha} \left[1 - e^{-\frac{1-\alpha}{\alpha}t}\right]\right)^{-\frac{p+1}{q}} \left(I_{0r}^\alpha f^{1/p}(t)\right)^p. \end{aligned} \tag{45}$$

Proof. Putting $g(\tau) = 1$ into theorem 22, yields

$$[I_{0r}^\alpha f^p(t)]^{1/p} \cdot [I_{0r}^\alpha(1)]^{1/q} \leq \left(\frac{M}{m}\right)^{1/pq} \cdot I_{0r}^\alpha[f(t)],$$

which is equivalent to

$$\begin{aligned} & [I_{0r}^\alpha f^p(t)]^{1/p} \\ & \leq \left(\frac{M}{m}\right)^{1/pq} \cdot \left(\frac{1}{1-\alpha} \left[1 - e^{-\frac{1-\alpha}{\alpha}t}\right]\right)^{-1/q} \cdot I_{0r}^\alpha[f(t)]. \end{aligned} \tag{46}$$

Substituting $g(\tau) = 1$ into theorem 21 and using (44), we obtain

$$[I_{0r}^\alpha f(t)]^{1/p} \cdot [I_{0r}^\alpha(1)]^{1/q} \leq \left(\frac{M}{m}\right)^{1/p^2q} \cdot I_{0r}^\alpha[f^{1/p}(t)],$$

that is

$$\begin{aligned} & [I_{0r}^\alpha f(t)]^{1/p} \\ & \leq \left(\frac{M}{m}\right)^{1/p^2q} \cdot \left(\frac{1}{1-\alpha} \left[1 - e^{-\frac{1-\alpha}{\alpha}t}\right]\right)^{-1/q} \cdot I_{0r}^\alpha[f^{1/p}(t)]. \end{aligned}$$

Hence, we can write

$$\begin{aligned} & I_{0r}^\alpha f(t) \\ & \leq \left(\frac{M}{m}\right)^{1/pq} \left(\frac{1}{1-\alpha} \left[1 - e^{-\frac{1-\alpha}{\alpha}t}\right]\right)^{-p/q} \left(I_{0r}^\alpha[f^{1/p}(t)]\right)^p. \end{aligned} \tag{47}$$

Combining (46) with (47), inequality (45) follows.

Acknowledgement

This work was supported by Universidad Nacional de Guinea Ecuatorial (UNGE) and Havana University. The authors would like to thank the anonymous reviewers for their valuable suggestions that helped improve the clarity of this paper.

4 Conclusion

In this paper, we presented some interesting fractional inequalities using the Caputo-Fabrizio fractional integral. The results are concerned with some inequalities using functions with the same sense of variation. As a future work, authors are planning to use these inequalities to prove existence and uniqueness of some Caputo-Fabrizio ordinary fractional differential equations.

References

- [1] Z. Dahmani. The Riemann-Liouville Operator to Generate Some New Inequalities. International Journal of Nonlinear Science (2011), Vol.12, pp. 452-453.
- [2] Mohamed Houas, Mohamed Bezziou. On Some Fractional Integral Inequalities Involving Generalized Riemann-Liouville Fractional Integral, Med. J. Model. Simul.(2016), Vol. 06, pp. 59-60.
- [3] Guotao Wang et. al., A Note on Saigo's Fractional Integral Inequalities. Turkish Journal of Analysis and Number Theory (2014), Vol. 2, No. 3, pp. 65-69.
- [4] S. D. Purohit and R. K. Raina, Chebyshev Type Inequalities for the Saigo Fractional integrals and their q-analogues. Journal of Mathematical Inequalities (2013), Vol. 7, pp. 239-248
- [5] Zoubir Dahmani and Hanane Metakkel El Ard, Generalizations of Some Integral Inequalities Using Riemann-Liouville Operator, Int. J. Open Problems Compt. Math (2011), Vol.4, pp. 41-45.
- [6] Nasser Al-Salti, Erkinjon Karimov and Sebti Kerbal, Boundary-value problems for fractional heat equation involving Caputo-Fabrizio derivative. New Trends in Mathematical Sciences (2016), Vol.4, pp. 79-80.
- [7] Jorge Losada and Juan J. Nieto, Properties of a New Fractional Derivative without Singular Kernel Progr. Fract. Differ. Appl (2015), No.2, pp. 87-89
- [8] S. Taf, K. Brahim, Some new results using Hadamard fractional integral. Int. J. Nonlinear Anal. Appl.(2016), vol.7, No. 1, pp. 103-109.
- [9] Ahmed Anber and et al., New integral inequalities of Feng Qi type via Riemann-Liouville fractional integration, Ser. Math. Inform (2012), Vol. 27, pp. 163-164.
- [10] Michele Caputo and Mauro Fabrizio, Applications of New Time and Spatial Fractional Derivatives with Exponential Kernels, Progr. Fract. Differ. Appl.(2016), Vol.2, pp. 7-8.
- [11] Zhanbing Bai, Haishen Lu, Positive solutions for boundary value problem of nonlinear fractional differential equation, J. Math. Anal. Appl. (2005), Vol. 311, pp. 496-497.
- [12] Vajjanath L. Chinchane, and Deepak B. Pachpatte, New Fractional Inequalities Involving Saigo Fractional Integral Operator, Mathematical Sciences Letters (2014), Vol.3, pp. 134-138.
- [13] M. Caputo and M. Fabrizio, A new Definition of Fractional Derivative without Singular Kernel. Progr. Fract. Differ. Appl. (2015), vol. 1, No. 2, pp. 73-85.
- [14] Feng Gao and Xiao-Jun Yang, Fractional Maxwell Fluid with Fractional Derivative without Singular Kernel, Thermal Science (2016), Vol. 20, pp. 872-873.

- [15] Abdon Atangana and Ilknur Koca, On the new fractional derivative and application to nonlinear Baggs and Freedman model. *J. Nonlinear Sci. Appl.* (2016), vol. 9, pp. 2467-2480.
- [16] J. Choi and P. Agarwal, Some new Saigo type fractional integral inequalities and their q -Analogues. *Abstract and Applied Analysis* (2014), Volume 2014, Article ID 579260, 11 pages, <http://dx.doi.org/10.1155/2014/579260>.
- [17] Vaijanath L. Chinchane, and Deepak B. Pachpatte, A Note on Some Fractional Integral Inequalities Via Hadamard Integral, *Journal of Fractional Calculus and Applications* (2013), Vol.4, pp. 125-128.
- [18] J.F. Gómez-Aguilar et. al, Electrical circuits described by a fractional derivative with regular Kernel, *Revista Mexicana de Física* (2016), vol. 62, pp. 144-154.
- [19] J. F. Gómez-Aguilar et. al, Modeling of a Mass-Spring-Damper System by Fractional Derivatives with and without a Singular Kernel. *Entropy* (2015), vol. 17, pp. 6289-6303.
- [20] W. Sudsutad, S. K. Ntouyas, J. Tariboon, Fractional Integral Inequalities via Hadamard's Fractional Integral. *Abstract and Applied Analysis* (2014), Volume 2014, Article ID 563096, 11 pages, <http://dx.doi.org/10.1155/2014/563096>.
- [21] D. Baleanu, A. Jajarmi, S.S. Sajjadi and D. Mozyrska, A new fractional model and optimal control of a tumor-immune surveillance with non-singular derivative operator, *Chaos* (2019), vol. 29, No. 8, 083127.
- [22] A. Jajarmi, S. Arshad and D. Baleanu, A new fractional modelling and control strategy for the outbreak of dengue fever, *Physica A* (2019) vol. 535, 122524.
- [23] A. Jajarmi, D. Baleanu, S.S. Sajjadi, J.H. Asad, A new feature of the fractional Euler-Lagrange equations for a coupled oscillator using a nonsingular operator approach, *Frontiers in Physics* (2019), vol.7, No. 196, 00196.
- [24] A. Jajarmi, B. Ghanbari and Dumitru Baleanu, A new and efficient numerical method for the fractional modelling and optimal control of diabetes and tuberculosis co-existence, *Chaos* (2019), vol. 29, No.9, 093111.
- [25] Anselmi C., Desantis P. and Scipioni A. , Nanoscale mechanical and dynamical properties of DNA single molecules, *Biophys Chem.*(2005), Vol. 113, pp. 209-221.



Gustavo Asumu received the B.S. and M.S. degrees from Kharkov University, Ukraine in 2011 and 2012, respectively. Now he is Pursuing Ph. D in Mathematics at Havana University, Cuba. He is Professor in Mathematics at Universidad Nacional de Guinea Ecuatorial(UNGE). His research interest is image processing with fractional partial differential equation.



Ángela León Mecías received her Ph.D in Mathematics from the University of Havana, Cuba in 2007. She is a Full Professor at the Applied Mathematics Department, Faculty of Mathematics and Computer Science, University of Havana. Dr. León is the general coordinator of the Master Program in Mathematics of the University of Havana. Her research interests include: numerical solution of differential equations and applications, mathematical methods in image processing and wavelet transform applications.



Mariano Rodríguez Ricard, Full Professor and Consultant at University of Havana. Areas of interest: bifurcations in ODE, pattern formation in reaction diffusion PDE. Hi is member of the Academic Committee of the Doctor Degree Program in Mathematics.