

April 2021

FINDINGS OF VIDEO HEAD IMPULSE TEST IN VESTIBULAR NEURITIS

Sara Omar Elfarouk Zaitoun

Audiovestibular Medical Unit, ENT department, faculty of medicine , zagazig university,
ziton8080@gmail.com

Soha Abd-el Raouf Mekki

Audiovestibular Medical Unit, ENT department, faculty of medicine , zagazig university,
soha.mekki@hotmail.com

Walid Mohamed Nooh

Audiovestibular Medical Unit, ENT department, faculty of medicine , zagazig university,
walidnooh@yahoo.com

Nahla Hassan Gad

Audiovestibular Medical Unit, ENT department, faculty of medicine , zagazig university,
nahla_audiology@yahoo.com

Follow this and additional works at: <https://digitalcommons.aaru.edu.jo/zumj>

Recommended Citation

Zaitoun, Sara Omar Elfarouk; Mekki, Soha Abd-el Raouf; Nooh, Walid Mohamed; and Gad, Nahla Hassan (2021) "FINDINGS OF VIDEO HEAD IMPULSE TEST IN VESTIBULAR NEURITIS," *Zagazig University Medical Journal*: Vol. 27 : Iss. 2 , Article 17.

Available at: <https://digitalcommons.aaru.edu.jo/zumj/vol27/iss2/17>

This Original Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in Zagazig University Medical Journal by an authorized editor. The journal is hosted on [Digital Commons](#), an Elsevier platform. For more information, please contact rakan@aar.edu.jo, marah@aar.edu.jo, dr_ahmad@aar.edu.jo.



Manuscript ID ZUMJ-1909-1508 (R1)

DOI 10.21608/zumj.2019.16812.1508

ORIGINAL ARTICLE**Findings of Video Head Impulse Test in Vestibular Neuritis**Sara Omar El farouk Zaiton¹, Soha Abdelraouf Mekki¹, Walid Mohamed Ibraheem¹ and Nahla Hassan Gad¹¹Audio-vestibular unit, Department of ENT, Faculty of Medicine, Zagazig University, Zagazig, Egypt**Corresponding author**Dr. Sara Omar El farouk Zaitoun
Faculty of Medicine,
Department of ENT,
Zagazig University, Zagazig,
Egypt
EMail: Ziton8080@gmail.com

Submit Date 2019-09-10

Revise Date 2019-09-24

Accept Date 2019-09-30

ABSTRACT**Background:** Vestibular neuritis is an acute unilateral peripheral vestibular disease without other signs of brain stem involvement. A comprehensive multidisciplinary assessment of VN patients including history and vestibular function tests holds great promise for more effective diagnosis. The video head impulse test (vHIT) assesses the vestibulo-ocular reflex (VOR) by using a video-assisted examination of the impulsive maneuver. We studied the result of vHIT in normal population versus vestibular neuritis patients..**Subjects and methods:** A case control study was conducted in Audio-vestibular unit, Department of ENT, Zagazig University Hospitals. 24 patients with vestibular neuritis were included. vHIT was done.**Results:** We calculated gain, gain asymmetry in the study group and compare the results with control group.**Conclusion:** vHIT is the only test that assesses all six semicircular canals independently and with a physiological stimulus, similar to how the patient uses the vestibular ocular reflex system in daily life.**Keywords:** vestibular neuritis, VOR, vHIT.**INTRODUCTION**

Vestibular neuritis represents 5% of all dizziness causes (15% of vestibular causes of dizziness). [1] It diagnosed clinically by history of: acute onset of persistent rotational vertigo, postural imbalance associated with nausea and vomiting without hearing loss lasting approximately 24-72 hours. After several days, the symptoms subsides gradually with varying degrees of dizziness or imbalance with head movement.[2]

Video head impulse test (vHIT) is a relatively new objective quantitative test of vestibular function. It was described by **Halmagi and Curthoys in 1988**. It is a video based equipment test that enables to detect the eye response (vestibulo-ocular reflex) to sudden head impulse stimulations. It used to measure the gain of the vestibulo-ocular reflex and to detect refixation saccades which are detected by using head impulse

stimulations. It assesses all semicircular canals individually. [3]

We thought that vHIT has an additional role in diagnosis of vestibular neuritis.

SUBJECTS AND METHODS

Written informed consent was obtained from all participants and the study was approved by the research ethical committee of Faculty of Medicine, Zagazig University. The work has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for studies involving humans.

All participants in the current study were subjected to the following:

Full history taking: including personal, general medical, otological and detailed dizziness history.

Otological examination.

Basic audiological evaluation.(audiometry and immittance)

Video head impulse test :The vHIT was performed using an Otometrics ICS impulse

system. Recordings were obtained for each of the six semicircular canals in all patients (horizontal, LARP, RALP).

STATISTICAL ANALYSIS

Continuous variables were presented as the mean±SD (normally distributed) or median(range) if not normally distributed. Normality was checked by Shapiro-Wilk test. Homogeneity of variances was checked by Leven's test. Categorical variables were presented by the count and percentage.

Fisher's Exact Test: for (2X2) (RXC) table is used to discover if there is a relationship between two categorical variables. It is an alternative to chi-squared test when the expected cell count is less than five.

Independent-samples t-test: is used to determine if a difference exists between the means of two independent groups on a continuous dependent variable.

Mann-Whitney U test: (nonparametric alternative to independent-samples t-test).

Cohen's Kappa: is a measure of inter-rater agreement for categorical variables when there are two raters.

Threshold for significance:

P-value<.05 indicates a statistically significant difference, P≤.01 indicates a highly a statistically significant difference, P≤.001 indicates a very highly statistically significant difference while, P≥.05 indicates a non-statistically significant difference.

Two-sided tests were used throughout.

All statistical analyses were performed using IBM SPSS Statistics, version 24 (IBM; Armonk, New York, USA).

RESULTS

In the present study, Patients with vestibular neuritis had highly significantly lower lateral right gain compared to left lateral gain Other differences between right and left gain were non-significant (P>0.05), as shown in Table(1).

Compared to controls, patients with vestibular neuritis had statistically significantly higher lateral gain asymmetry (P=0.02), but no statistically significant difference in gain asymmetry (RALP and LALP) were found (P=0.83 vs P=0.24), as shown in Table(2)

In the present study, not all the patients studied showed covert and overt corrective saccades in the vHIT tests. Almost 8 patients (33%) presented overt saccades; 5 patients (20%) presented covert saccades as showed in table (3).

Also we found that the most affected canals in the study group were the lateral canals (29.1%) and the number of uni-canal affection (45%) greater than multi-canal affection (4.2%) Table (4)

In the present study, there was no-statistically significant effect of age on vHIT parameters in vestibular neuritis patients as shown inTable (5)

There was statistical significant +ve correlation between disease duration and gain of right lateral canal among the study group as shown inTable (6)

Table (1): Gain in video-head impulse (vHIT) in patients with vestibular neuritis:

Gain %	Rt	Lt	Independent samples-t test	P-value
	n=24	n=24		
Gain				
Lateral %			3.09	0.003 (S)
mean±SD	0.77±0.31	1.11±0.44		
Anterior %			1.90	0.06 (NS)
mean±SD	1.08±0.29	1.25 ±0.33		
Posterior %			0.75	0.45 (NS)
mean±SD	1.14±0.49	1.06±0.19		

Table (2): Gain asymmetry in video-head impulse (vHIT) in patients with vestibular neuritis and control group

Gain asymmetry (%)	Study group n=24 Median(Range)	Control group n=24 Median(Range)	Mann-Whitney U test	P-value
Lateral	8(0-47.5)	5(0-6.9)	312	0.02 (S)
RALP	6(1-22)	5.5(1-6.5)	135.6	0.83(NS)
LARP	4(1-20)	4(1-7)	213.4	0.24(NS)

Table (3): Saccades in video-head impulse (vHIT) in patients with vestibular neuritis

Saccades	Cases=24
LAT	
COVERT, n (%)	
Negative	21(88)
Right positive	3(12)
OVERT, n (%)	
Negative	18(75)
Right positive	6(25)
LARP	
COVERT, n (%)	
Negative	24(100)
OVERT, n (%)	
Negative	24(100)
RALP	
COVERT, n (%)	
Negative	22(92)
Right anterior positive	2(8)
OVERT, n (%)	
Negative	22(92)
Left posterior positive	2(8)

Table (4): Semicircular canal affection in patients with vestibular neuritis by vHIT.

Affected canal, n (%)	Cases=24
Unilateral canal affection	
Lateral canal	7(29.1)
Posterior canal	3(12.5)
Anterior canal	1(4.2)
Multi- canal affection	1(4.2)
No affection	12(50)

Table (5): Correlation between Age and vHIT measures in study group:

Variable	Age	
	R	P
Gain of lateral canal (right):	-0.708	0.105
Gain of lateral canal (left):	-0.695	0.111
Gain of anterior canal (right):	-0.621	0.604

Gain of anterior canal (left):	-0.007	0.660
Gain of posterior canal (right):	-0.045	0.796
Gain of posterior canal (left):	-0.035	0.811
Gain of asymmetry of lateral canal:	-0.714	0.103
Gain of asymmetry of RALP canal:	-0.033	0.552
Gain of asymmetry of LARP canal:	-0.467	0.204

Table (6):Correlation between disease duration and both caloric and vHIT measures in study group

Variable	Disease duration (months)	
	R	P
Gain of lateral canal (right):	0.45	0.03*
Gain of lateral canal (left):	-0.18	0.41 NS
Gain of anterior canal (right):	-0.11	0.60 NS
Gain of anterior canal (left):	-0.11	0.60 NS
Gain of posterior canal (right):	-0.14	0.50 NS
Gain of posterior canal (left):	-0.14	0.50 NS
Gain of asymmetry of lateral canal:	-0.07	0.74 NS
Gain of asymmetry of RALP canal:	0.26	0.26 NS
Gain of asymmetry of LARP canal:	0.13	0.56 NS

DISCUSSION

Vestibular neuritis is an acute unilateral peripheral vestibular disease without other signs of brain stem involvement. A comprehensive multidisciplinary assessment of VN patients including history and vestibular function tests holds great promise for more effectively diagnosis.

We assess the results of the vHIT in normal population versus vestibular neuritis patients.

The assessment of the vHIT results relies on the analysis of both the gain and gain asymmetry of the VOR and refixation saccades. Yang et al. [4] hypothesized that the normative values of vHIT gain are >0.8 without saccades. This was consistent with the results of the control group but disagreed with Halmagyi et al. [5] who found that values of vHIT gain more than 0.6 is considered normal.

In this study, patients with vestibular neuritis had highly significantly lower lateral

right gain ($P=0.005$). Other differences in right and left gain in both groups were non-significant.

The cause of high affection of lateral canals may be related to the anatomic basis of superior vestibular nerve which supplies the horizontal semicircular canal (along with reduced diameter and increased bony trabeculae of the bony canal housing the superior vestibular nerve and its divisions) make it more likely to be involved in cases of vestibular neuritis opposite to double supply of the IVN in two separate bony canals Taylor et al. [6]. This finding disagreed with Milonski et al. [7] and Mangabeira et al. [8] who found that up to 42% of VN patients present affection in the posterior semicircular canal

In the present study, it was found that not all studied patients showed covert and overt corrective saccades in the vHIT results. Zellhuber et al. [9] focused on CSs as well as vHIT gain values to evaluate hypo function in patients with vestibular disorders as using

gain and CS improved diagnostic accuracy and the sensitivity of vHIT for identifying vestibular loss. This was disagreed with Yang et al. [4] who found that the studies of diagnosis of VN using vHIT relied on gain parameters only but not CS measurements to detect the presence of vestibular hypo function.

Also we found that the most affected canals in the study group were the lateral canals (29.1%) and the number of uni-canal affection (45%) greater than multi-canal affection (4.2%) .

In this study, there was no statistically significant effect of age on vHIT parameters. This finding may be related that all subjects in this study were on the middle age. Regarding the individual function of the vestibular end organs, research has revealed that the functions of all six semicircular canals decline with age starting at age of 40 years Mossman et al. [10] Therefore, the age effect in the present study did not reach significance.

In our study, there were statistical +ve correlation between disease duration and gain of right lateral canal among the study group as in the sequela phase of vestibular neuritis (after 1 to 3 monthes), the vHIT gain recovers at greater velocity until normality returns. Rambold [11] This result also was in agreement with Magliulo, et al. [12] who found that 85.7% of patients had resumed normal vHIT within 1 year after onset.

CONCLUSION

Video head Impulse test is the only test that assesses all six semicircular canals independently and with a physiological stimulus, similar to how the patient uses the vestibular ocular reflex system in daily life.

Conflict of interest:no

Financial Disclosures:no

REFERENCES

- [1] Brodsky JR, Cusick BA and Zhou G.(2016)Vestibular neuritis in children and adolescents: Clinical features and recovery. *Int J Pediatr Otorhinolaryngol* ;83(4):104-8. doi: 10.1016.
- [2] Strupp M and Brandt T. (2009): Vestibular neuritis *Semin Neurol.* ;29(5):509-519.
- [3] Bartl K, Lahnen N, Kohlbecher S and Schneider, E. (2009): Basics and clinical aspects of vertigo and dizziness: Head impulse test using video-oculography *Annals of the New York Academy of Sciences*; 1164:331-333.
- [4] Yang, C., Lee, J., Kang, B. , Lee, H. , Yoo, M. and Park, H. (2016): Quantitative analysis of gains and catch-up saccades of video-head-impulse testing by age in normal subjects. *Clin Otolaryngol.*; 41:532-538
- [5] Halmagyi G., Karlberg M., Curthoys I. and Todd M.(2002): Inferior vestibular neuritis. *Ann N Y Acad Sci. Apr*;956:306–13.
- [6] Taylor, R., McGarvie, L., Reid, N., Young, A., Halmagyi, G. and Welgampola, M. (2016): Vestibular neuritis affects both superior and inferior vestibular nerves. *Neurology* 87:1704–1712.
- [7] Mangabeira Albernaz P. and Zuma EMFC (2014): The video head impulse test. *Acta Otolaryngol.*; 134: 1245-50.
- [8] Miłonski, J., Pietkiewicz, P., Bielinska, M., Kuśmierczyk, K. and Olszewski J. (2014): The use of videonystagmography head impulse test (VHIT) in the diagnostics of semicircular canal injuries in patients with vertigo. *Int J Occup Med Environ Health*; 27: 583-90.
- [9] Zellhuber S, Mahringer A and Rambold H.(2014): Relation of videohead-impulse test and caloric irrigation: a study on the recovery in unilateral vestibular neuritis. *Eur Arch Otorhinolaryngol.* 2014;271:2375-2383
- [10] Mossman B, Mossman S, Purdie G and Schneider E(2015): Age dependent normal horizontal VOR gain of head impulse test as measured with videooculography. *J Otolaryngol Head Neck Surg*;44(1):29
- [11] Rambold, H. (2014): Economic management of vertigo/dizziness disease in a county hospital: video-head-impulse test vs caloric irrigation. *Eur Arch Otorhinolaryngol.*; 272:2621---8.
- [12] Magliulo, G., Iannella, G., Gagliardi S. and Re M (2015): A 1-year follow-up study with C-VEMPs, O-VEMPs and video head impulse testing in vestibular neuritis. *Eur Arch Otorhinolaryngol*; 272 (11): 3277-81

HOW TO CITE

zaitoun, S., Mekki, S., nooh, W., gad, N. FINDINGS OF VIDEO HEAD IMPULSE TEST IN VESTIBULAR NEURITIS. *Zagazig University Medical Journal*, 2021; 2(359-363): -. doi: 10.21608/zumj.2019.16812.1508