

2018

Recent advances in understanding theories of eruption (evidence based review article)

Amany A. Rabea
Amany.ahmed@fue.edu.eg

Follow this and additional works at: <https://digitalcommons.aaru.edu.jo/fdj>



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

A. Rabea, Amany (2018) "Recent advances in understanding theories of eruption (evidence based review article)," *Future Dental Journal*: Vol. 4 : Iss. 2 , PP 189-196.

Available at: <https://digitalcommons.aaru.edu.jo/fdj/vol4/iss2/16>

This Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in Future Dental Journal by an authorized editor. The journal is hosted on [Digital Commons](#), an Elsevier platform. For more information, please contact rakan@aarj.edu.jo, marah@aarj.edu.jo, u.murad@aarj.edu.jo.

Recent advances in understanding theories of eruption (evidence based review article)


Cover Page Footnote

Peer review under responsibility of Faculty of Oral & Dental Medicine, Future University. E-mail address: Amany.Ahmed@fue.edu.eg.



Contents lists available at ScienceDirect

Future Dental Journal

journal homepage: www.elsevier.com/locate/fdj

Recent advances in understanding theories of eruption (evidence based review article)

Amany A. Rabea

Associate Professor of Oral Biology, Faculty of Oral and Dental Medicine, Future University in Egypt, Egypt

ARTICLE INFO

Keywords:

Eruption
Theories
Recent
Molecular determinants

ABSTRACT

Eruption is; movement of the tooth from its developmental site in alveolar bone to its functional position in the oral cavity. Eruption has been a matter of long historical debate. Each of the eruption theories has a say in tooth eruption. So, the aim of this research is to explore the integration of eruption theories to understand the aetiology of the eruption process.

1. Introduction

Eruption is necessary for survival of diverse species [1]. Eruption is continuous process. It does not stop by reaching the occlusal plane, but continues throughout life [2].

2. Types of tooth eruption

2.1. Active eruption

Active eruption is; the movement of the tooth from its developmental site in alveolar bone to its functional position in the oral cavity [3]. Eruption can generally be divided into different phases [4].

2.2. Pre-eruptive movement

It is made by the teeth germs. It is best thought of as the mean by which the teeth are positioned within the jaw for eruptive movement. The Pre-eruptive phase starts from the end of early bell stage till the beginning of root formation [4].

2.3. Eruptive movement

Tooth movements during eruptive phase are subdivided into; intraosseous and supraosseous stages. The eruptive phase begins with the onset of root formation and terminates by tooth appearance in the oral cavity, just before function (pre-functional phase) [4].

2.4. Post-eruptive movement

This movement maintains the tooth position in occlusion by compensation for occlusal and proximal tooth wear. The post-eruptive phase starts when the teeth attain occlusion and continues for as long as each tooth remains in the oral cavity (functional phase) [4].

2.5. Passive eruption

Passive eruption is characterized by the apical shift of the dento-gingival junction. As this occurs, the length of the clinical crown increases as the epithelial attachment migrates apically [5].

Although the movement of teeth into function has been the subject of extensive research, there is no consensus regarding the mechanism involved [1].

2.6. Normal eruption of human teeth

Tooth eruption in humans has numerous aspects [6]. The eruption process starts with onset of root formation [7]. From this time till tooth appearance in the oral cavity is called eruption time [6].

3. Theories of eruption

There is no consistent understanding of the cause behind tooth eruption. The aetiology behind eruption and explanation of the eruption mechanism seem to be essential to perform aetiology based treatment [6].

Peer review under responsibility of Faculty of Oral & Dental Medicine, Future University.
E-mail address: Amany.Ahmed@fue.edu.eg.

<https://doi.org/10.1016/j.fdj.2018.05.001>

Received 18 January 2018; Received in revised form 14 April 2018; Accepted 6 May 2018
Available online 09 May 2018

2314-7180/ © 2018 Published by Elsevier B.V. on behalf of Faculty of Oral & Dental Medicine, Future University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1. Diagram of basal end of rat incisor showing cushioned hammock ligament [8].

3.1. Eruption theories with recent points of view

3.1.1. Cushioned hammock theory

It was proposed by Harry Sicher [8]. This theory assumed that ligament (cushioned hammock ligament) below a tooth is responsible for eruption (Fig. 1).

However the ligament described by Sicher [8] was an artifact in slide preparation [9].

3.1.2. Root formation theory

The root formation theory assumes that the proliferating root encounters a fixed structure; and the apically directed force is converted into a reactive occlusal force that causes coronal movement of the erupting tooth (Fig. 2) [4].

However, there are facts refuted this hypothesis such as; rootless teeth can erupt, some teeth erupt greater distance than the total root length; and the teeth erupt after completion of root formation or when the tissue forming the root is removed [4]. Also, the onset of root formation does not coincide with the eruptive movement [4]. Moreover, newly formed dentin at root apex is unmineralized and can be deformed by trauma [10].

Recently it is postulated that since tension results in bone




Fig. 2. Photomicrograph showing root formation [4].

deposition, the tensile responses that occur during teeth eruption are of great concern.

In the (intra-osseous stage of eruptive phase), root formation and jaw growth lead to compressive coronal hydrostatic stress. This induces the dental follicle and stellate reticulum cells to secrete mediators for bone resorption. Furthermore, the root formation would produce tensile apical hydrostatic stress in the teeth germs that leads to bone deposition [11,12]. Moreover, pre-occlusal eruption (supra-osseous stage of eruptive phase) is completed by root growth and bone formation at crypt base [3].

3.1.3. Vascular pressure/blood vessel thrust or hydrostatic pressure theory

This theory suggests that a local increase in tissue fluid pressure in the periapical region is sufficient to move the tooth [13].

However, this is debatable because root and local vasculature excision, does not prevent tooth eruption [13].

Recently it is reported that the hydrostatic pressure theory occurs during postemergent eruption [12]. This is due to that dental follicle secrete mediators, such as vascular endothelial growth factor (VEGF), that cause angiogenesis and so increase in the apical tissue pressure that lead to tooth eruption [12]. Moreover, hydrostatic pressure theory was supported by several studies that confirm tooth eruption after a local injection of vasodilators [12]. Whereas injection of vasoconstrictors caused decrease in the rate of eruption [14].

3.1.4. Bone remodeling theory

This theory based on; bone resorption occurs coronally and bone apposition occurs apically. The dental follicle is the source for osteoblasts and osteoclasts [4,13].

Whether bone remodeling that occurs around teeth causes or is the effect of tooth movement is not known, and both circumstances may apply [13]. The strongest evidence in support of bone remodeling as a cause of tooth movement comes from a series of experiments in dogs. When the developing premolar is removed without disturbing the dental follicle, or if eruption is prevented by wiring the tooth germ down to the lower border of the mandible, an eruptive pathway still forms within the bone overlying the enucleated tooth as osteoclasts widen the gubernacular canal (Fig. 3). If the dental follicle is removed, however, no eruptive pathway forms. Furthermore, if a metal or silicone replica replaces tooth germ and so as long as the dental follicle is




Fig. 3. Photomicrograph showing bone remodeling [4].



Fig. 4. Clinical photograph of patient with mutation in the PTH1R gene [12].

retained, the replica will erupt, with formation of an eruptive pathway. It is concluded that programmed bone remodeling can and does occur (i.e., an eruptive pathway forms in bone without a developing and growing tooth). Second, the dental follicle is involved. However, the conclusion cannot be drawn that the demonstration of an eruptive pathway forming within bone means that bone remodeling is responsible for tooth movement unless coincident bone deposition also can be demonstrated at the base of the crypt where its prevention can interfere with tooth eruption [4].

Recently the molecular basis of tooth eruption supports bone remodeling theory as it confirms that mutation of the parathyroid hormone receptor 1 (PTH1R) gene is correlated with disturbances in bone remodeling and leads to primary failure of eruption (PFE) [12]. PFE is nonsyndromic eruption disturbance that is not associated with defective osteoclasts [12]. The affected teeth had supra-crestal presentation and progressive open bite which is hallmark criteria of PFE (Fig. 4) [12,15].

3.1.5. Dental follicle theory

The follicular theory postulates that the dental follicle is capable of inducing, bone resorption above the developing crown and bone apposition below it (Fig. 5). This enables the formation of an eruptive path to occur through which the tooth will be passively conducted [16]. In osteopetrotic animal, which lack a factor that stimulates differentiation of osteoclasts, eruption is prevented, because no mechanism for bone removal exists. However, local administration of this factor, colony-stimulating factor 1 (CSF-1), permits the differentiation of osteoclasts and eruption occurs [4].

Recently molecular studies have revealed that eruption is regulated by inductive signals between the dental follicle, reduced enamel epithelium (REE), stellate reticulum and alveolar bone [17]. Regional differences in the dental follicle were described [17]. It is suggested that the coronal aspect of the dental follicle regulates osteoclastogenesis (bone resorption) and the basal aspect of the dental follicle regulates osteogenesis (bone formation) [17]. This was assessed using laser capture microdissection (Fig. 6). Real time reverse transcription-




Fig. 5. Photomicrograph of a developing dog tooth indicating the position of the dental follicle (F) around the developing tooth, developing mandibular alveolar bone (M) and overlying oral epithelium (E) [17].




Fig. 6. A diagram showing: Laser capture microdissection (LCM) [18].

polymerase chain reaction (RT-PCR) was used to assess the expression of bone resorption and bone formation marker genes. The receptor activator of nuclear factor kappa B ligand (RANKL) gene is a marker gene for bone resorption. Bone morphogenetic protein-2 (BMP-2) gene is a marker for bone formation. The results showed a higher expression of RANKL genes in the coronal half of the dental follicle, and higher expression of BMP-2 genes in the basal half of the follicle [17].

3.1.5.1. Current concepts concerning the paracrine signaling function of the dental follicle in tooth eruption. Tooth development is regulated by a cascade of mutual interactions between the dental epithelium and the dental mesenchyme [19].

Correspondingly, the process of tooth eruption is regulated by cellular events leading to the recruitment of monocytes to the dental follicle followed by bone resorption [20]. This molecular events are initiated by interactions between the dental follicle, the REE and the stellate reticulum [21].

Apoptosis of epithelial cells during the advanced stages of enamel secretion have been reported [22]. This apoptotic process has a direct influence on osteoclastogenesis through the release of interleukin-1 α (IL-1 α) by stellate reticulum cells, which its receptors are located in the dental follicle [20]. IL-1 α consequently stimulates the expression of CSF-1 and monocyte chemoattractant protein-1 (MCP-1) in the dental follicle, allowing the dental follicle to act as a chemoattractant for monocytes [20]. The stellate reticulum cells also participate in bone resorption by releasing parathyroid hormone-related protein (PTHrP) which further increases the expression of both MCP-1 and CSF-1 [23]. CSF-1 down-regulates the expression of osteoprotegerin (OPG), a well-known snare receptor for RANKL which inhibit osteoclast differentiation [24]. The cells of the REE and stellate reticulum therefore exert a paracrine effect on the dental follicle, enhancing the expression of chemoattractant molecules [25]. Additionally, the REE also secretes proteases that aid in creating an eruption pathway through enzymatic digestion of collagens [26]. Other molecules such as epidermal growth factor (EGF) and transforming growth factor β 1 (TGF- β 1) released by the cells of the dental follicle enhance also the expression of CSF-1 in the dental follicle (Figs. 7 and 8) [27].

Bone resorption is however not sufficient for the displacement of the tooth. Thus, coronal bone resorption must be coupled with apical bone formation [28]. The expression of these BMPs is greatly enhanced by tumor necrosis factor- α (TNF- α) [28]. The cascade of signaling events at the apical aspect of the developing tooth, are not completely elucidated [29].

Matrix metalloproteinases (MMPs) are capable of degrading extracellular matrix (ECM) structural. They play important roles in tissue development [31].

Membrane-type 1 matrix metalloproteinase (MT1-MMP) is specific for collagens (I, II, and III), gelatin, fibronectin, and other matrix molecules [32]. MT1-MMP is essential during development and expressed




Fig. 7. Paracrine signaling at the coronal half of the erupting tooth [17].




Fig. 8. Paracrine signaling between the stellate reticulum and dental follicle, as well as within the dental follicle only [30].

in the tooth and the surrounding connective tissue [33,34].

Migratory capabilities of follicle cells are disturbed in the absence of MT1-MMP, as collagen content increases in the dental follicle. Also MMPs have role in cell migration [35].

3.1.6. Periodontal ligament traction theory

This theory is based on the postulation that periodontal ligament (PDL)-dental follicle complex possesses eruptive force due to the traction power that fibroblasts have [4]. In continuously growing rodent and rabbit incisors teeth, surgical procedures performed and effectively eliminated dental pulpal pressure, dentin formation, and the cervical loop from contributing to incisor eruption, and PDL is the only tissue responsible for eruption of this tooth [36]. Moreover, fibroblasts move incisally along the erupting tooth, and their contraction generates significant force for tooth eruption [37].

However, experiments in a rat model using lathyrogens (amino acid derivatives that cause defective fibril formation when applied to the

PDL) did not cause different rate of eruption when compared to untreated rats [38]. Moreover, the rate of collagen turnover is much higher than that of eruption [39]. Additionally, there is no difference in metabolic structures within fibroblasts in the PDL of rapidly erupting teeth and fully erupted teeth [40]. Also, periodontal fibroblasts exhibit characteristics of cells actively synthesizing and secreting [41]. Moreover, the ability of rootless teeth to erupt on schedule indicates that the PDL is not essential for eruption [42].

Recently it is hypothesized that functional stress results in greater strain in dental follicle and PDL than in bone, these soft tissues act as stress sensors [43]. Bone resorption and deposition involved with tooth movement, is critical surface phenomena at the interface between soft and bony tissues surrounding the developing tooth [44]. Moreover, teeth normally drift forward and upwards in the jaws, which is not the case in either osseointegrated implants or when there is pathological fusion of the bone to teeth [45]. Furthermore, The PDL has mechanosensor activity during orthodontic treatment [46].

Lately, it is confirmed that PDL fibroblast orientation increased significantly during eruption (Fig. 9) [47]. It is proposed that PDL fibroblasts connect to collagen fibers inserting into the tooth's cementum so their occlusal migration pulls the tooth toward the oral cavity [47].

PDL fibroblasts are spindle-shaped with multiple processes that join with other cells during ligament development [48]. The fibroblasts long axis is aligned parallel with the orientation of collagen fibers [49]. They both create and are responsive to mechanical forces through focal adhesion complexes [50]. In PDL fibroblasts, changes in the strain environment induce changes in gene expression, such as the upregulation of collagen 1 [51]. Also, the importance of the CD44 receptor in cell-cell and cell-matrix interactions is demonstrated [52]. The proliferation and migration of PDL cells has been linked with CD44/HA interactions [53].

3.1.7. Molecular determinants in tooth eruption

Eruption molecules control the timing of the cellular events of eruption [54].

The RANKL needed for alveolar bone resorption come from the dental follicle. Also CSF-1, down-regulates the expression of OPG to enable osteoclastogenesis to occur. Also secreted frizzled-related protein-1 (SFRP-1) that inhibits osteoclastogenesis, also has its gene expression down-regulated in the dental follicle [55].

CSF-1 and MCP-1 were maximally expressed in the dental follicle. Endothelial monocyte-activating polypeptide (EMAP-II) has been shown to have a chemotactic effect on mononuclear cells. It up regulates the gene expression of both CSF-1 and MCP-1 [56].

The transcription factor gene c-fos and the transcription factor genes nuclear factor kappa-light-chain-enhancer of activated B cells (NFκB 1) and (NFκB 2) are needed for osteoclast differentiation [57]. Type I receptor of interleukin-1α (IL-1R), is present in the dental follicle, and its gene is enhanced by a ligand in the stellate reticulum [58].

Prior to eruption, CSF-1 expression is reduced and appears to be replaced by VEGF which is highly expressed in the dental sac. VEGF up regulates the expression of RANK [59].

BMP-2 is highly expressed in the basal half of the dental follicle. BMPs regulate the expression of Cbfa1 (core binding factor a1). It acts as a key transcriptional regulator of osteoblast differentiation during bone formation [60].

At the onset of eruption, the sialoprotein of 95,000 relative molecular weight (DF-95), is reduced by exactly the amount of three new sialoproteins with MW 20–25,000. Fragmentation of DF-95 is a biochemical marker of the beginning of tooth eruption. Immunolocalization of DF-95 in the REE is a biochemical evidence in initiation of eruption. Proteases of the enamel organ cause fragmentation of DF-95 and release metalloproteinase thus they initiate eruption [61].

Wnt/β-catenin signaling plays an important role in bone formation and regeneration [62].




Fig. 9. Photomicrographs showing: PDL fibroblast orientation within the cervical region during mandibular first molar (M1) eruption and occlusion. Intraosseous eruption, (A–C), mucosal penetration, (D–F), and preocclusal eruption (G–I). Enamel (e), dentin (d), bone (b) and the PDL (p). KO, knockout; WT, wild type. Scale bar = 100 μ m [47].




Fig. 10. Photomicrograph showing gubernacular canal and cord [4].

The gubernacular cord formation starts from the remnants of dental lamina cells [63]. This structure is located behind the deciduous tooth [64]. In this cord, there is EGF (epithelial growth factor), which has the capacity to induce bone resorption leaving a space surrounding this cord, so-called the gubernacular canal (Fig. 10) [65].

3.2. Recent theories of eruption

3.2.1. Bite forces sensed by soft tissue dental follicles theory

This theory postulates that follicular soft tissues detect bite-forces and so direct bone remodeling with the effect of enabling tooth eruption [11].

The authors reported the highest equivalent strains induced by bite

forces in the dental follicle and PDL in both erupted and unerupted teeth irrespective of incisive or unilateral molar bite forces (Fig. 11). Examination of the soft tissue dental follicles, suggested broad areas of compression in overlying crowns, and wide zones of tension in follicle below root apices (Fig. 12) [11]. So, these soft act as relevant stress sensors [11].

3.2.2. Innervation-provoked pressure theory

This theory postulates that the root membrane acts as a glandular membrane. So, the innervation in this membrane causes pressure in the apical part of the tooth which results in tooth eruption [6].

It is hypothesized that tooth eruption depends on [6].

- (1) Space in the pathway of eruption,
- (2) Pressure from below,
- (3) Adaptation of the periodontal membrane.

1. The crown follicle creates the necessary space in the eruption path [6].

2. The root membrane functions as a glandular membrane. So, the innervation causes overpressure that causes the tooth to elevate in the eruption direction [6,66,67].

3. The adaptability of the periodontal membrane is essential for eruption [6,68].

3.2.3. The equilibrium theory

After the functional plane is reached, the eruption of the tooth is balanced in response to the growth of the vertical growth of the mandible [69].

As the mandible grows vertically away from the maxilla, the teeth have more room to erupt occlusally in order to maintain occlusal contact with the opposing arch. This model of tooth eruption reinforces the idea that postemergent tooth eruption, after reaching functional occlusion, is controlled by forces impeding eruption, as opposed to encouraging forces. These balancing forces of masticatory function and the soft tissue pressures from the lips, cheeks, and tongue are the rate-limiting factors of postfunctional occlusal eruption [12,69]




Fig. 11. Diagrams showing patterns of equivalent strain. Irrespective the loading applied, equivalent strain was high in soft tissues of the PDL (red arrows) and dental follicles (green arrows), and lower levels of strain in the hard tissues [11].




Fig. 12. Dental follicle compression (red) and tension (green) during bite forces. The upper surfaces of dental follicles are subject to greater compression, as compared with the lower surfaces which were subject to greater tension [11].

The lasting eruptive movement that occurs while the teeth are free of contact, supports the idea that eruptive control is based on the continuous force of the surrounding soft tissues [12].

3.2.4. Neuromuscular theory or unification theory

The neuromuscular theory or unification theory of tooth eruption states that the synchronized forces of the orofacial muscles, under the control of the central nervous system, are responsible for the active movements of a tooth and the molecular events prepared a pathway under the control of these forces [68,70–74].

The coordinated neuromuscular forces are converted into electrical, electrochemical and biomechanical energies for the stimulation of cellular and molecular activities within and around the dental follicle and enamel organ to prepare a pathway as well as other cellular functions for eruption of a developing tooth [55,70].

4. Evidence based on current systematic reviews

- [1] A detailed analysis of the findings from scientific studies on mechanisms of tooth eruption was conducted [70].
- [2] Systematic analysis of epidemiologic studies as well as many studies on animal tissues was carried out to understand the mechanisms behind tooth eruption [6].
- [3] Other authors conducted a systematic review to outline the possible mechanism of tooth eruption right from its development in the bony crypt to its eruption till the occlusal level [3].

5. Conclusions and future considerations

- To sum up, tooth eruption must be considered as a stage of tooth development.
- Root follicle, periodontal membrane, and crown follicle are involved in the eruption process.
- Dental follicle and PDL play mechanosensor role in tooth eruption.
- The orientation of PDL fibroblasts determines the tooth directional movement.
- Root formation produces compressive coronal and tensile apical hydrostatic stress resulting in tooth eruption.
- The molecular and enzymic activities, are controlled by neuromuscular forces.
- Each of the eruption theories has a say to some portion of the eruption process.

The presence of stem cells in the dental follicle [75], raises around their potential role in tooth eruption. Identification of role of stem cells during tooth eruption still needs further research.

References

- [1] Rajic Z, Rajic-Mestrovic S, Verzak Z. Chronology, dynamics and period of permanent tooth eruption in zagreb children (Part II). *Coll Antropol* 2000;24(1):137–43.
- [2] Newman HN. On passive eruption. *J West Soc Periodontol Periodontol Abstr* 1994;42(2):41–4.
- [3] Srinath SK, Sahana S, Vishwanath SK, Ritu S. Mechanism of tooth eruption & its clinical significance - a systematic review of literature. *Elixir Dentistry* 2013;65:19676–80.
- [4] Nanci A. *Ten Cate's oral histology: development, structure and function*. eighth ed. Elsevier; 2013.
- [5] Young Yim J, Il Kim T, Jo Seol Y, Moo Lee Y, Ku Y, Chul Rhyu I, Pyoung Chung C, Boo Han S. Crown lengthening for altered passive eruption. *J Korean Acad Periodontol*. 2008;38(2):247–52.
- [6] Kjær I. Mechanism of human tooth eruption: review article including a new theory for future studies on the eruption process. *Sci Tech Rep* 2014;2014:1–13.
- [7] Björk A, Skieller V. Normal and abnormal growth of the mandible. A synthesis of longitudinal cephalometric implant studies over a period of 25 years. *EJO (Eur J Orthod)* 1983;5(1):1–46.
- [8] Sicher H. Tooth eruption: the axial movement of continuously growing teeth. *J Drug Res (Cairo)* 1942;21:395–402.
- [9] Edward HF. *Craniofacial development, growth and evolution*. first ed. Bateson; 2002.
- [10] Topouzelis N, Tsaousoglou P, Pisoka V, Zouloumis L. Dilaceration of maxillary

- central incisor: a literature review. *Dent Traumatol* 2010;26:427–33.
- [11] Sarrafpour B, Swain M, Li Q, Zoellner H. Tooth eruption results from bone remodeling driven by bite forces sensed by soft tissue dental follicles: a finite element analysis. *PLoS One* 2013;8(3):1–18.
- [12] Frazier-Bowers SA, Hendricks HM. Failure of tooth eruption: diagnosis and management. In: Wright JT, editor. *Craniofacial and dental developmental defects: diagnosis and management*. Springer; 2015.
- [13] Bhaskar SN. *Orban's oral histology and embryology*. eleventh ed. Mosby; 1991.
- [14] Shimada A, Shibata T, Komatsu K. Relationship between the tooth eruption and regional blood flow in angiotensin II-induced hypertensive rats. *Arch Oral Biol* 2004;49:427–33.
- [15] Frazier-Bowers SA, Puranik CP, Mahaney MC. The etiology of eruption disorders – further evidence of a ‘genetic paradigm.’ *Semin. Orthod*. 2010;16(3):180–5.
- [16] Marks Jr. SC, Cahill DR. Experimental study in the dog of the non-active role of the tooth in the eruptive process. *Arch Oral Biol* 1984;29:311–22.
- [17] Nel S, Hendrik HD, Boy SC, Raubenheimer EJ. Recent perspectives vis-à-vis the biological basis of tooth eruption. *SADJ* 2015;70(6):238–41.
- [18] Marko-Varga GA, Fehniger TE. Microscale protein expression profiling during disease evolution. *J Chromatogr* 2004;1053:279–90.
- [19] Bector KB, Nolting D, Bector JP, Kjaer I. Immunohistochemical localization of epithelial rests of Malassez in human periodontal membrane. *Eur J Orthod* 2007;29(4):350–3.
- [20] Wise GE, Zhao L, Grier RL. Localization and expression of CSF-1 receptor in rat dental follicle cells. *J Dent Res* 1997;76(6):1244–9.
- [21] Vahtokari A, Vainio S, Thesleff I. Associations between transforming growth factor beta 1 RNA expression and epithelial-mesenchymal interactions during tooth morphogenesis. *Development* 1991;113(3):985–94.
- [22] Vahtokari A, Aberg T, Thesleff I. Apoptosis in the developing tooth: association with an embryonic signaling center and suppression by EGF and FGF-4. *Development* 1996;122(1):121–9.
- [23] Wise GE, Que BG, Huang H, Lumpkin SJ. Enhancement of gene expression in rat dental follicle cells by parathyroid hormone-related protein. *Arch Oral Biol* 2000;45(10):903–9.
- [24] Wise GE, Ren Y, Yao S. Regulation of osteoprotegerin gene expression in dental follicle cells. *J Dent Res* 2003;82(4):298–302.
- [25] Wise GE, Marks SC, Zhao L. Effect of CSF-1 on *in vivo* expression of c-fos in the dental follicle during tooth eruption. *Eur J Oral Sci* 1998;106(1):397–400.
- [26] Park SJ, Bae HS, Cho YS, Lim SR, Kang SA, Park JC. Apoptosis of the reduced enamel epithelium and its implications for bone resorption during tooth eruption. *J Mol Histol* 2013;44(1):65–73.
- [27] Fox SW, Lovibond AC. Current insights into the role of transforming growth factor-beta in bone resorption. *Mol Cell Endocrinol* 2005;243(1–2):19–26.
- [28] Yao S, Prpic V, Pan F, Wise GE. TNF-alpha upregulates expression of BMP-2 and BMP-3 genes in the rat dental follicle—implications for tooth eruption. *Connect Tissue Res* 2010;51(1):59–66.
- [29] Bradaschia-Correa V, Casado-Gomez I, Moreira MM, Ferreira LB, Arana-Chavez VE. Immunolocalization of Smad-4 in developing molar roots of alendronate-treated rats. *Arch Oral Biol* 2013;58(11):1744–50.
- [30] Wise GE. Cell and molecular biology of tooth eruption. In: Davidovitch Z, editor. *Biological mechanisms of tooth eruption, resorption and replacements by implants*. AL: EBSCO Media; 1998.
- [31] Klein T, Bischoff R. Physiology and pathophysiology of matrix metalloproteinases. *Amino Acids* 2011;41:271–90.
- [32] Itoh Y. Membrane-type matrix metalloproteinases: their functions and regulations. *Matrix Biol* 2015;44–46:207–23.
- [33] Evans BR, Mosig RA, Lobl M, Martignetti CR, Camacho C, Grum-Tokars V, Glucksman MJ, Martignetti JA. Mutation of membrane type-1 metalloproteinase, MT1-MMP, causes the multicentric osteolysis and arthritis disease Winchester syndrome. *Am J Hum Genet* 2012;91:572–6.
- [34] Xu H, Snider TN, Wimer HF, Yamada SS, Yang T, Holmbeck K, Foster BL. Multiple essential MT1-MMP functions in tooth root formation, dentinogenesis, and tooth eruption. *Matrix Biol* 2016;52–54:266–83.
- [35] Fraley SI, Wu PH, He L, Feng Y, Krisnamurthy R, Longmore GD, Wirtz D. Three-dimensional matrix fiber alignment modulates cell migration and MT1-MMP utility by spatially and temporally directing protrusions. *Sci Rep* 2015;5:14580.
- [36] Moxham BJ, Berkovitz BK. The effects of root transection on the unimpeded eruption rate of the rabbit mandibular incisor. *Arch Oral Biol* 1974;19:903–9.
- [37] Kasugai S, Suzuki S, Shibata S, Yasui S, Amano H, Ogura H. Measurements of the isometric contractile forces generated by dog periodontal ligament fibroblasts *in vitro*. *Arch Oral Biol* 1990;35:597–601.
- [38] Berkovitz BK, Migdalski A, Solomon M. The effect of the lathyritic agent aminocetonitrile on the unimpeded eruption rate in normal and root-resected rat lower incisors. *Arch Oral Biol* 1972;17(12):1755–63.
- [39] Berkovitz BK. How teeth erupt. *Dent Update* 1990;17:206–10.
- [40] Ten Cate AR, Deporter DA, Freeman E. The role of fibroblasts in the remodeling of periodontal ligament during physiological tooth movement. *Am J Orthod* 1976;69:155–68.
- [41] Taverne AA. Collagen responsible for tooth eruption? A study of the eruption of rat incisors. *Aust Orthod J* 1993;12(4):199–206.
- [42] Brin I, Zilberman Y, Galili D, Fuks A. Eruption of rootless teeth in congenital renal disease. *Oral Surg Oral Med Oral Pathol* 1985;60:61–4.
- [43] Yao S, Pan F, Wise GE. Chronological gene expression of parathyroid hormone-related protein (PTHrP) in the stellate reticulum of the rat: implications for tooth eruption. *Arch Oral Biol* 2007;52:228–32.
- [44] Sarrafpour B, Rungsiyakull C, Swain M, Li Q, Zoellner H. Finite element analysis suggests functional bone strain accounts for continuous post-eruptive emergence of

- teeth. *Arch Oral Biol* 2012;57:1070–8.
- [45] Blanco Carrion J, Ramos Barbosa I, Perez Lopez J. Osseointegrated implants as orthodontic anchorage and restorative abutments in the treatment of partially edentulous adult patients. *Int. J. Periodontics Restorative Dent.* 2009;29:333–40.
- [46] Wu J, Li Y, Fan X, Zhang C, Wang Y, et al. Analysis of gene expression profile of periodontal ligament cells subjected to cyclic compressive force. *DNA Cell Biol* 2011;30:865–73.
- [47] Popowics T, Boyd T, Hinderberger H. Eruptive and functional changes in periodontal ligament fibroblast orientation in CD44 wild-type vs. knockout mice. *J Periodontal Res* 2014;49:355–62.
- [48] Yamamoto T, Wakita M. Bundle formation of principal fibers in rat molars. *J Periodontal Res* 1992;27:20–7.
- [49] McCulloch CAG, Lekic P, McKee MD. Role of physical forces in regulating the form and function of the periodontal ligament. *Periodontol* 2000;2000(24):56–72.
- [50] Chiquet M, Gelman L, Lutz R, Maier S. From mechanotransduction to extracellular matrix gene expression in fibroblasts. *Biochim Biophys Acta* 2009;1793:911–20.
- [51] Kook SH, Hwang JM, Park JS, et al. Mechanical force induces type I collagen expression in human periodontal ligament fibroblasts through activation of ERK/JNK and AP-1. *J Cell Biochem* 2009;106:1060–7.
- [52] Naor D, Nedvetzki S, Walmsley M, et al. CD44 involvement in autoimmune inflammations. *Ann N Y Acad Sci* 2007;1110:233–47.
- [53] Shimabukuro Y, Terashima H, Takedachi M, et al. Fibroblast growth factor-2 stimulates directed migration of periodontal ligament cells via P13K/AKT signaling and CD44/Hyaluronan interaction. *J. Cell Physiol.* 2011;226:809–21.
- [54] Wise GE, Frazier-Bowers S, D'Souza RN. Cellular, molecular, and genetic determinants of tooth eruption. *Crit Rev Oral Biol Med* 2002;13:323–34.
- [55] Wise GE. Cellular and molecular basis of tooth eruption. *Orthod Craniofac Res* 2009;12(2):67–73.
- [56] Liu D, Wise GE. Expression of endothelial monocyte activating polypeptide II in the rat dental follicle and its potential role in tooth eruption. *Eur J Oral Sci* 2008;116:334–40.
- [57] Iotsova V, Caamaño J, Loy J, Yang Y, Lewin A, Bravo R. Osteopetrosis in mice lacking NF- κ B1 and NF- κ B2. *Nat Med* 1997;3:1285–9.
- [58] Que BG, Lumpkin SJ, Wise GE. Effect of IL-1 α on NF κ B gene expression in the dental follicle—implications for tooth eruption. *Arch Oral Biol* 1999;44:961–7.
- [59] Yao S, Liu D, Pan F, Wise GE. Effect of vascular endothelial growth factor on RANK gene expression in osteoclast precursors and on osteoclastogenesis. *Arch Oral Biol* 2006;51:596–602.
- [60] Wise GE, Yao S. Regional differences of expression of BMP-2 and RANKL in the rat dental follicle. *Eur. J. Oral Sci* 2006;114:512–6.
- [61] Marks SC, Schroeder HE. Tooth eruption: theories and facts. *Anat Rec* 1996;245:374–93.
- [62] Tak-Heun K, Cheol-Hyeon B, Eun-Ha J, Chi-Young Y, Young B, Seung O, et al. Col1a1-cre mediated activation of b-catenin leads to aberrant dento-alveolar complex formation. *Anat. Cell Biol.* 2012;45:193–202.
- [63] Consolaro A. Tracionamento ortodôntico: possíveis consequências nos caninos superiores e dentes adjacentes (parte 1). *Dental Press J. Orthod.* 2010;15(4):15–23.
- [64] Ide F, Mishima K, Kikuchi K, Horie N, Yamachika S, Satomura K, et al. Development and growth of adenomatoid odontogenic tumor related to formation and eruption of teeth. *Head and Neck Pathol.* 2011;5:123–32.
- [65] Consolaro A. Reabsorções dentárias nas especialidades clínicas. third ed. Dental Press; 2012.
- [66] Kjær I, Kocsis G, Nodal M, Christensen LR. Aetiological aspects of mandibular tooth agenesis-focusing on the role of nerve, oral mucosa, and supporting tissues. *EJO (Eur J Orthod)* 1994;16(5):371–5.
- [67] Becktor KB, Hansen BF, Nolting D, Kjaer I. Spatiotemporal expression of NGFR during pre-natal human tooth development. *Orthod Craniofac Res* 2002;5(2):85–9.
- [68] Bille ML, Thomsen B, Kjær I. Apoptosis in the human periodontal membrane evaluated in primary and permanent teeth. *Acta Odontol Scand* 2011;69(6):385–8.
- [69] Proffit WR. Contemporary orthodontics. fifth ed. Elsevier; 2013.
- [70] Loto AO. Tooth eruption: a neuromuscular theory, part one. *J. Craniomax. Res.* 2017;4(1):278–83.
- [71] Wise GE, King GJ. Tooth movement. *J Dent Res* 2008;87:414–34.
- [72] Ruta A, Irena B, Janina T. Factors influencing permanent teeth eruption. Part one – general factors. *Stomatologija. Baltic Dental and Maxillofacial Journal* 2010;12:67–72.
- [73] Ash MM, Nelson SJ. Wheeler's dental anatomy, physiology, and occlusion. ninth ed. Elsevier; 2003.
- [74] Kiliaridis S. The importance of masticatory muscle function in dentofacial growth. *Semin Orthod* 2006;12(2):110–9.
- [75] Yao S, Pan F, Prpic V, Wise GE. Differentiation of stem cells in the dental follicle. *J Dent Res* 2008;87:767–71.