

2015

Characterizations of b-Soft Separation Axioms in Soft Topological Spaces

S. A. El-Sheikh

*Mathematics Department, Faculty of Education, Ain Shams University, Cairo, Egypt.,
alaa_8560@yahoo.com*

Rodyna A. Hosny

Mathematics Department, Faculty of Science, Zagazig University, Zagazig, Egypt., alaa_8560@yahoo.com

A. M. Abd El-latif

*Mathematics Department, Faculty of Education, Ain Shams University, Cairo, Egypt.,
alaa_8560@yahoo.com*

Follow this and additional works at: <https://digitalcommons.aaru.edu.jo/isl>

Recommended Citation

A. El-Sheikh, S.; A. Hosny, Rodyna; and M. Abd El-latif, A. (2015) "Characterizations of b-Soft Separation Axioms in Soft Topological Spaces," *Information Sciences Letters*: Vol. 4 : Iss. 3 , Article 5.
Available at: <https://digitalcommons.aaru.edu.jo/isl/vol4/iss3/5>

This Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in *Information Sciences Letters* by an authorized editor. The journal is hosted on [Digital Commons](#), an Elsevier platform. For more information, please contact rakan@aarj.edu.jo, marah@aarj.edu.jo, u.murad@aarj.edu.jo.

Characterizations of *b*-Soft Separation Axioms in Soft Topological Spaces

S. A. El-Sheikh¹, Rodyna A. Hosny² and A. M. Abd El-latif^{1,*}

¹ Mathematics Department, Faculty of Education, Ain Shams University, Cairo, Egypt.

² Mathematics Department, Faculty of Science, Zagazig University, Zagazig, Egypt.

Received: 2 Jul. 2015, Revised: 3 Aug. 2015, Accepted: 4 Aug. 2015

Published online: 1 Sep. 2015

Abstract: Many scientists have studied and improved the soft set theory, which is initiated by Molodtsov [33] and easily applied to many problems having uncertainties from social life. The main purpose of our paper, is to introduce new soft separation axioms based on the *b*-open soft sets which are more general than of the open soft sets. We show that, the properties of soft *b-T_i*-spaces ($i = 1, 2$) are soft topological properties under the bijection and irresolute open soft mapping. Also, the property of being soft *b*-regular and soft *b*-normal are soft topological properties under bijection, irresolute soft and irresolute open soft functions. Further, we show that the properties of being soft *b-T_i*-spaces ($i = 1, 2, 3, 4$) are hereditary properties.

Keywords: Soft set, Soft topological space, Soft interior, Soft closure, Open soft, Closed soft, Soft *b*- separation axioms, Soft *b-T_i*-spaces ($i = 1, 2, 3, 4$), Soft *b*-regular, Soft *b*-normal, *b*-irresolute soft functions, Irresolute *b*-open soft functions.

1 Introduction

In real life situation, the problems in economics, engineering, social sciences, medical science etc. do not always involve crisp data. So, we cannot successfully use the traditional classical methods because of various types of uncertainties presented in these problems. To exceed these uncertainties, some kinds of theories were given like theory of fuzzy set, intuitionistic fuzzy set, rough set, bipolar fuzzy set, i.e. which we can use as mathematical tools for dealings with uncertainties. But, all these theories have their inherent difficulties. The reason for these difficulties Molodtsov [33] initiated the concept of soft set theory as a new mathematical tool for dealing with uncertainties which is free from the above difficulties. In [33,34], Molodtsov successfully applied the soft theory in several directions, such as smoothness of functions, game theory, operations research, Riemann integration, Perron integration, probability, theory of measurement, and so on. After presentation of the operations of soft sets [31], the properties and applications of soft set theory have been studied increasingly [6,27,34]. Xiao et al.[44] and Pei and Miao [37] discussed the relationship between soft sets and information systems. They showed that soft sets are a

class of special information systems. In recent years, many interesting applications of soft set theory have been expanded by embedding the ideas of fuzzy sets [4,5,9,16,25,29,30,31,32,34,35,47]. To develop soft set theory, the operations of the soft sets are redefined and a uni-int decision making method was constructed by using these new operations [10].

Recently, in 2011, Shabir and Naz [40] initiated the study of soft topological spaces. They defined soft topology as a collection τ of soft sets over X . Consequently, they defined basic notions of soft topological spaces such as open soft and closed soft sets, soft subspace, soft closure, soft nbd of a point, soft separation axioms, soft regular spaces and soft normal spaces and established their several properties. Min in [43] investigate some properties of these soft separation axioms. In [17], Kandil et al. introduced some soft operations such as semi open soft, pre open soft, α -open soft and β -open soft and investigated their properties in detail. Kandil et al. [24] introduced the notion of soft semi separation axioms. In particular they study the properties of the soft semi regular spaces and soft semi normal spaces. The notion of soft ideal was initiated for the first time by Kandil et al.[20]. They also introduced the concept of soft local function. These concepts are discussed with a view to find

* Corresponding author e-mail: Alaa_8560@yahoo.com, Alaa8560@hotmail.com

new soft topologies from the original one, called soft topological spaces with soft ideal (X, τ, E, \tilde{I}) . Applications to various fields were further investigated by Kandil et al. [18, 19, 21, 22, 23, 26]. The notion of supra soft topological spaces was initiated for the first time by El-sheikh and Abd El-latif [13]. They also introduced new different types of subsets of supra soft topological spaces and study the relations between them in detail. The notion of b-open soft sets was initiated for the first time by El-sheikh and Abd El-latif [12], which is extended by Abd El-latif et al. in [1]. Maji et al. [29] initiated the study involving both fuzzy sets and soft sets. In [8] the notion of fuzzy soft set was introduced as a fuzzy generalization of soft sets and some basic properties of fuzzy soft sets are discussed in detail. Then, many scientists such as X. Yang et al. [45], improved the concept of fuzziness of soft sets. In [2], Karal and Ahmed defined the notion of a mapping on classes of fuzzy soft sets, which is a fundamental important in fuzzy soft set theory, to improve this work and they studied properties of fuzzy soft images and fuzzy soft inverse images of fuzzy soft sets. Chang [11] introduced the concept of fuzzy topology on a set X by axiomatizing a collection \mathfrak{T} of fuzzy subsets of X . Tanay et al. [41] introduced the definition of fuzzy soft topology over a subset of the initial universe set while Roy and Samanta [39] gave the definition of fuzzy soft topology over the initial universe set. Some fuzzy soft topological properties based on fuzzy semi open soft sets namely, fuzzy semi open soft sets, fuzzy semi closed soft sets, fuzzy semi soft interior, fuzzy semi soft closure fuzzy semi separation axioms and fuzzy soft semi connectedness, were introduced by Kandil et al. in [16, 25].

The purpose of this paper, is to introduce the notion of soft b-separation axioms. In particular we study the properties of the soft b-regular spaces and soft b-normal spaces. We show that if x_E is b-closed soft set for all $x \in X$ in a soft topological space (X, τ, E) , then (X, τ, E) is soft b- T_1 -space. Also, we show that if a soft topological space (X, τ, E) is soft b- T_3 -space, then $\forall x \in X$, x_E is b-closed soft set. This paper, not only can form the theoretical basis for further applications of topology on soft sets, but also lead to the development of information systems.

2 Preliminaries

Definition 2.1.[33] Let X be an initial universe and E be a set of parameters. Let $P(X)$ denote the power set of X and A be a non-empty subset of E . A pair (F, A) denoted by F_A is called a soft set over X , where F is a mapping given by $F : A \rightarrow P(X)$. In other words, a soft set over X is a parametrized family of subsets of the universe X . For a particular $e \in A$, $F(e)$ may be considered the set of e -approximate elements of the soft set (F, A) and if $e \notin A$, then $F(e) = \phi$ i.e

$F_A = \{F(e) : e \in A \subseteq E, F : A \rightarrow P(X)\}$. The family of all these soft sets over X denoted by $SS(X)_A$.

Definition 2.2.[31] Let $F_A, G_B \in SS(X)_E$. Then F_A is soft subset of G_B , denoted by $F_A \subseteq G_B$, if

- (1) $A \subseteq B$, and
- (2) $F(e) \subseteq G(e), \forall e \in A$.

In this case, F_A is said to be a soft subset of G_B and G_B is said to be a soft superset of F_A , $G_B \supseteq F_A$.

Definition 2.3.[31] Two soft subset F_A and G_B over a common universe set X are said to be soft equal if F_A is a soft subset of G_B and G_B is a soft subset of F_A .

Definition 2.4.[6] The complement of a soft set (F, A) , denoted by $(F, A)^c$, is defined by $(F, A)^c = (F^c, A)$, $F^c : A \rightarrow P(X)$ is a mapping given by $F^c(e) = X - F(e)$, $\forall e \in A$ and F^c is called the soft complement function of F . Clearly, $(F^c)^c$ is the same as F and $((F, A)^c)^c = (F, A)$.

Definition 2.5.[40] The difference between two soft sets (F, E) and (G, E) over the common universe X , denoted by $(F, E) - (G, E)$ is the soft set (H, E) where for all $e \in E$, $H(e) = F(e) - G(e)$.

Definition 2.6.[40] Let (F, E) be a soft set over X and $x \in X$. We say that $x \in (F, E)$ read as x belongs to the soft set (F, E) whenever $x \in F(e)$ for all $e \in E$. The soft set (F, E) over X such that $F(e) = \{x\} \forall e \in E$ is called singleton soft point and denoted by x_E or (x, E) .

Definition 2.7.[31] A soft set (F, A) over X is said to be a NULL soft set denoted by $\tilde{\phi}$ or ϕ_A if for all $e \in A$, $F(e) = \phi$ (null set).

Definition 2.8.[31] A soft set (F, A) over X is said to be an absolute soft set denoted by \tilde{X} or X_A if for all $e \in A$, $F(e) = X$. Clearly, we have $X_A^c = \phi_A$ and $\phi_A^c = X_A$.

Definition 2.9.[40] Let (F, E) be a soft set over X and $x \in X$. We say that $x \in (F, E)$ read as x belongs to the soft set (F, E) whenever $x \in F(e)$ for all $e \in E$.

Definition 2.10.[31] The union of two soft sets (F, A) and (G, B) over the common universe X is the soft set (H, C) , where $C = A \cup B$ and for all $e \in C$,

$$H(e) = \begin{cases} F(e), & e \in A - B, \\ G(e), & e \in B - A, \\ F(e) \cup G(e), & e \in A \cap B. \end{cases}$$

Definition 2.11.[31] The intersection of two soft sets (F, A) and (G, B) over the common universe X is the soft set (H, C) , where $C = A \cap B$ and for all $e \in C$, $H(e) = F(e) \cap G(e)$. Note that, in order to efficiently discuss, we consider only soft sets (F, E) over a universe X in which all the parameter set E are same. We denote the family of these soft sets by $SS(X)_E$.

Definition 2.12.[48] Let I be an arbitrary indexed set and $L = \{(F_i, E), i \in I\}$ be a subfamily of $SS(X)_E$.

- (1) The union of L is the soft set (H, E) , where $H(e) = \bigcup_{i \in I} F_i(e)$ for each $e \in E$. We write $\bigcup_{i \in I} (F_i, E) = (H, E)$.

(2)The intersection of L is the soft set (M, E) , where $M(e) = \bigcap_{i \in I} F_i(e)$ for each $e \in E$. We write $\tilde{\bigcap}_{i \in I} (F_i, E) = (M, E)$.

Definition 2.13.[40] Let τ be a collection of soft sets over a universe X with a fixed set of parameters E , then $\tau \subseteq SS(X)_E$ is called a soft topology on X if

- (1) $\tilde{X}, \tilde{\phi} \in \tau$, where $\tilde{\phi}(e) = \phi$ and $\tilde{X}(e) = X, \forall e \in E$,
- (2)the union of any number of soft sets in τ belongs to τ ,
- (3)the intersection of any two soft sets in τ belongs to τ .

The triplet (X, τ, E) is called a soft topological space over X . A soft set (F, A) over X is said to be closed soft set in X , if its relative complement $(F, A)^c$ is an open soft set.

Definition 2.14.[14] Let (X, τ, E) be a soft topological space. The members of τ are said to be open soft sets in X . We denote the set of all open soft sets over X by $OS(X, \tau, E)$, or when there can be no confusion by $OS(X)$ and the set of all closed soft sets by $CS(X, \tau, E)$, or $CS(X)$.

Definition 2.15.[40] Let (X, τ, E) be a soft topological space and $(F, E) \in SS(X)_E$. The soft closure of (F, E) , denoted by $cl(F, E)$ is the intersection of all closed soft super sets of (F, E) i.e

$$cl(F, E) = \tilde{\bigcap} \{ (H, E) : (H, E) \text{ is closed soft set and } (F, E) \tilde{\subseteq} (H, E) \}.$$

Definition 2.16.[48] Let (X, τ, E) be a soft topological space and $(F, E) \in SS(X)_E$. The soft interior of (G, E) , denoted by $int(G, E)$ is the union of all open soft subsets of (G, E) i.e

$$int(G, E) = \tilde{\bigcup} \{ (H, E) : (H, E) \text{ is an open soft set and } (H, E) \tilde{\subseteq} (G, E) \}.$$

Definition 2.17.[48] The soft set $(F, E) \in SS(X)_E$ is called a soft point in X_E if there exist $x \in X$ and $e \in E$ such that $F(e) = \{x\}$ and $F(e^c) = \phi$ for each $e^c \in E - \{e\}$, and the soft point (F, E) is denoted by x_e .

Definition 2.18.[48] The soft point x_e is said to be belonging to the soft set (G, A) , denoted by $x_e \tilde{\in} (G, A)$, if for the element $e \in A, F(e) \subseteq G(e)$.

Theorem 2.1.[42] Let (X, τ, E) be a soft topological space. A soft point $x_e \tilde{\in} cl(F, E)$ if and only if each soft neighborhood of x_e intersects (F, E) .

Definition 2.19.[40] Let (X, τ, E) be a soft topological space and Y be a non null subset of X . Then \tilde{Y} denotes the soft set (Y, E) over X for which $Y(e) = Y \forall e \in E$.

Definition 2.20.[40] Let (X, τ, E) be a soft topological space, $(F, E) \in SS(X)_E$ and Y be a non null subset of X . Then the sub soft set of (F, E) over Y denoted by (F_Y, E) , is defined as follows:

$$F_Y(e) = Y \cap F(e) \forall e \in E.$$

In other words $(F_Y, E) = \tilde{Y} \tilde{\cap} (F, E)$.

Definition 2.21.[40] Let (X, τ, E) be a soft topological space and Y be a non null subset of X . Then

$$\tau_Y = \{ (F_Y, E) : (F, E) \in \tau \}$$

is said to be the soft relative topology on Y and (Y, τ_Y, E) is called a soft subspace of (X, τ, E) .

Theorem 2.2.[40] Let (Y, τ_Y, E) be a soft subspace of a soft topological space (X, τ, E) and $(F, E) \in SS(X)_E$. Then

- (1)If (F, E) is an open soft set in Y and $\tilde{Y} \in \tau$, then $(F, E) \in \tau$.
- (2) (F, E) is an open soft set in Y if and only if $(F, E) = \tilde{Y} \tilde{\cap} (G, E)$ for some $(G, E) \in \tau$.
- (3) (F, E) is a closed soft set in Y if and only if $(F, E) = \tilde{Y} \tilde{\cap} (H, E)$ for some (H, E) is τ -closed soft set.

Definition 2.22.[12] Let (X, τ, E) be a soft topological space and $(F, E) \in SS(X)_E$. Then (F, E) is called a b-open soft set if $(F, E) \tilde{\subseteq} cl(int(F, E)) \tilde{\cup} int(cl(F, E))$. The set of all b-open soft sets is denoted by $BOS(X, \tau, E)$, or $BOS(X)$ and the set of all b-closed soft sets is denoted by $BCS(X, \tau, E)$, or $BCS(X)$.

Definition 2.23.[12] Let (X, τ, E) be a soft topological space and $(F, E) \in SS(X)_E$. Then, the b-soft interior of (F, E) is denoted by $bSint(F, E)$, where $bSint(F, E) = \tilde{\bigcup} \{ (G, E) : (G, E) \tilde{\subseteq} (F, E), (G, E) \in BOS(X) \}$.

Also, the b-soft closure of (F, E) is denoted by $bScl(F, E)$, where $bScl(F, E) = \tilde{\bigcap} \{ (H, E) : (H, E) \in BCS(X), (F, E) \tilde{\subseteq} (H, E) \}$.

Definition 2.24.[3] Let $SS(X)_A$ and $SS(Y)_B$ be families of soft sets, $u : X \rightarrow Y$ and $p : A \rightarrow B$ be mappings. Let $f_{pu} : SS(X)_A \rightarrow SS(Y)_B$ be a mapping. Then;

- (1)If $(F, A) \in SS(X)_A$. Then the image of (F, A) under f_{pu} , written as $f_{pu}(F, A) = (f_{pu}(F), p(A))$, is a soft set in $SS(Y)_B$ such that

$$f_{pu}(F)(b) = \begin{cases} \bigcup_{x \in p^{-1}(b) \cap A} u(F(x)), & p^{-1}(b) \cap A \neq \phi, \\ \phi, & \text{otherwise.} \end{cases}$$

for all $b \in B$.

- (2)If $(G, B) \in SS(Y)_B$. Then the inverse image of (G, B) under f_{pu} , written as $f_{pu}^{-1}(G, B) = (f_{pu}^{-1}(G), p^{-1}(B))$, is a soft set in $SS(X)_A$ such that

$$f_{pu}^{-1}(G)(a) = \begin{cases} u^{-1}(G(p(a))), & p(a) \in B, \\ \phi, & \text{otherwise.} \end{cases}$$

for all $a \in A$.

The soft function f_{pu} is called surjective if p and u are surjective, also is said to be injective if p and u are injective.

Definition 2.25.[17, 28, 48]

Let (X, τ_1, A) and (Y, τ_2, B) be soft topological spaces and $f_{pu} : SS(X)_A \rightarrow SS(Y)_B$ be a function. Then, the function f_{pu} is said to be

- (1)The function f_{pu} is said to be continuous soft (cts-soft) if $f_{pu}^{-1}(G, B) \in \tau_1 \forall (G, B) \in \tau_2$.
- (2)The function f_{pu} is said to be open soft if $f_{pu}(G, A) \in \tau_2 \forall (G, A) \in \tau_1$.
- (3)The function f_{pu} is said to be b-irresolute soft if $f_{pu}^{-1}(G, B) \in BOS(X) [f_{pu}^{-1}(F, B) \in BCS(X)] \forall (G, B) \in BOS(Y) [(F, B) \in BCS(Y)]$.

(6) The function f_{pu} is said to be irresolute b-open (closed) soft if $f_{pu}(G, A) \in BOS(Y)[f_{pu}(F, A) \in BCS(Y)] \forall (G, A) \in BOS(X)[(F, A) \in BCS(Y)]$.

Theorem 2.3.[3] Let $SS(X)_A$ and $SS(Y)_B$ be families of soft sets. For the soft function $f_{pu} : SS(X)_A \rightarrow SS(Y)_B$, the following statements hold,

- $f_{pu}^{-1}((G, B)^c) = (f_{pu}^{-1}(G, B))^c \forall (G, B) \in SS(Y)_B$.
- $f_{pu}(f_{pu}^{-1}((G, B))) \subseteq (G, B) \forall (G, B) \in SS(Y)_B$. If f_{pu} is surjective, then the equality holds.
- $(F, A) \subseteq f_{pu}^{-1}(f_{pu}((F, A))) \forall (F, A) \in SS(X)_A$. If f_{pu} is injective, then the equality holds.
- $f_{pu}(\tilde{X}) \subseteq \tilde{Y}$. If f_{pu} is surjective, then the equality holds.
- $f_{pu}^{-1}(\tilde{Y}) = \tilde{X}$ and $f_{pu}(\tilde{\phi}_A) = \tilde{\phi}_B$.
- If $(F, A) \subseteq (G, A)$, then $f_{pu}(F, A) \subseteq f_{pu}(G, A)$.
- If $(F, B) \subseteq (G, B)$, then $f_{pu}^{-1}(F, B) \subseteq f_{pu}^{-1}(G, B) \forall (F, B), (G, B) \in SS(Y)_B$.
- $f_{pu}^{-1}[(F, B) \cup (G, B)] = f_{pu}^{-1}(F, B) \cup f_{pu}^{-1}(G, B)$ and $f_{pu}^{-1}[(F, B) \cap (G, B)] = f_{pu}^{-1}(F, B) \cap f_{pu}^{-1}(G, B) \forall (F, B), (G, B) \in SS(Y)_B$.
- $f_{pu}[(F, A) \cup (G, A)] = f_{pu}(F, A) \cup f_{pu}(G, A)$ and $f_{pu}[(F, A) \cap (G, A)] = f_{pu}(F, A) \cap f_{pu}(G, A) \forall (F, A), (G, A) \in SS(X)_A$. If f_{pu} is injective, then the equality holds.

3 Soft b-separation axioms

Definition 3.1. Let (X, τ, E) be a soft topological space and $x, y \in X$ such that $x \neq y$. Then, (X, τ, E) is called a soft b- T_0 -space if there exist b-open soft sets (F, E) and (G, E) such that either $x \in (F, E)$ and $y \notin (F, E)$ or $y \in (G, E)$ and $x \notin (G, E)$.

Proposition 3.1. Let (X, τ, E) be a soft topological space and $x, y \in X$ such that $x \neq y$. If there exist b-open soft sets (F, E) and (G, E) such that either $x \in (F, E)$ and $y \in (F, E)^c$ or $y \in (G, E)$ and $x \in (G, E)^c$. Then, (X, τ, E) is soft b- T_0 -space. **Proof.** Let $x, y \in X$ such that $x \neq y$. Let (F, E) and (G, E) be b-open soft sets such that either $x \in (F, E)$ and $y \in (F, E)^c$ or $y \in (G, E)$ and $x \in (G, E)^c$. If $x \in (F, E)$ and $y \in (F, E)^c$. Then $y \in (F(e))^c$ for all $e \in E$. This implies that, $y \notin F(e)$ for all $e \in E$. Therefore, $y \notin (F, E)$. Similarly, if $y \in (G, E)$ and $x \in (G, E)^c$, then $x \notin (G, E)$. Hence, (X, τ, E) is soft b- T_0 -space.

Theorem 3.1. A soft subspace (Y, τ_Y, E) of a soft b- T_0 -space (X, τ, E) is soft b- T_0 .

Proof. Let $x, y \in Y$ such that $x \neq y$. Then $x, y \in X$ such that $x \neq y$. Hence, there exist b-open soft sets (F, E) and (G, E) in X such that either $x \in (F, E)$ and $y \notin (F, E)$ or $y \in (G, E)$ and $x \notin (G, E)$. Since $x \in Y$. Then $x \in \tilde{Y}$. Hence, $x \in \tilde{Y} \cap (F, E) = (F_Y, E)$, (F, E) is b-open soft set. Consider $y \notin (F, E)$, This implies that, $y \notin F(e)$ for some $e \in E$. Therefore, $y \notin \tilde{Y} \cap (F, E) = (F_Y, E)$. Similarly, if $y \in (G, E)$ and $x \notin (G, E)$, then $y \in (G_Y, E)$ and $x \notin (G_Y, E)$. Thus, (Y, τ_Y, E) is soft b- T_0 .

Definition 3.2. Let (X, τ, E) be a soft topological space and $x, y \in X$ such that $x \neq y$. Then, (X, τ, E) is called a soft b- T_1 -space if there exist b-open soft sets (F, E) and (G, E) such that $x \in (F, E)$ and $y \notin (F, E)$ and $y \in (G, E)$ and $x \notin (G, E)$.

Proposition 3.2. Let (X, τ, E) be a soft topological space and $x, y \in X$ such that $x \neq y$. If there exist b-open soft sets (F, E) and (G, E) such that $x \in (F, E)$ and $y \in (F, E)^c$ and $y \in (G, E)$ and $x \in (G, E)^c$. Then (X, τ, E) is soft b- T_1 -space.

Proof. It is similar to the proof of Proposition 3.1.

Theorem 3.2. A soft subspace (Y, τ_Y, E) of a soft b- T_1 -space (X, τ, E) is soft b- T_1 .

Proof. It is similar to the proof of Theorem 3.

Theorem 3.3 Let (X, τ, E) be a soft topological space. If x_E is b-closed soft set in τ for all $x \in X$, then (X, τ, E) is soft b- T_1 -space.

Proof. Suppose that $x \in X$ and x_E is b-closed soft set in τ . Then x_E^c is b-open soft set in τ . Let $x, y \in X$ such that $x \neq y$. For $x \in X$ and x_E^c is b-open soft set such that $x \notin x_E^c$ and $y \in x_E^c$. Similarly y_E^c is b-open soft set in τ such that $y \notin y_E^c$ and $x \in y_E^c$. Thus, (X, τ, E) is soft b- T_1 -space over X .

Definition 3.3. Let (X, τ, E) be a soft topological space and $x, y \in X$ such that $x \neq y$. Then (X, τ, E) is called a soft b-Hausdorff space or soft b- T_2 -space if there exist b-open soft sets (F, E) and (G, E) such that $x \in (F, E)$, $y \in (G, E)$ and $(F, E) \cap (G, E) = \tilde{\phi}$.

Theorem 3.4. For a soft topological space (X, τ, E) we have:

soft b- T_2 -space \Rightarrow soft b- T_1 -space \Rightarrow soft b- T_0 -space.

Proof.

- Let (X, τ, E) be a soft b- T_2 -space and $x, y \in X$ such that $x \neq y$. Then, there exist b-open soft sets (F, E) and (G, E) such that $x \in (F, E)$, $y \in (G, E)$ and $(F, E) \cap (G, E) = \tilde{\phi}$. Since $(F, E) \cap (G, E) = \tilde{\phi}$. Then, $x \notin (G, E)$, $y \notin (F, E)$. Therefore, there exist b-open soft sets (F, E) and (G, E) such that $x \in (F, E)$ and $y \notin (F, E)$ and $y \in (G, E)$ and $x \notin (G, E)$. Thus, (X, τ, E) is soft b- T_1 -space.
- Let (X, τ, E) be a soft b- T_1 -space and $x, y \in X$ such that $x \neq y$. Then, there exist b-open soft sets (F, E) and (G, E) such that $x \in (F, E)$ and $y \notin (F, E)$ and $y \in (G, E)$ and $x \notin (G, E)$. Obviously then we have, either $x \in (F, E)$ and $y \notin (F, E)$ or $y \in (G, E)$ and $x \notin (G, E)$. Thus, (X, τ, E) is soft b- T_0 -space.

Remark 3.1. The converse of Theorem 3.4 is not true in general, as shown in the following examples.

Examples 3.1.

- Let $X = \{a, b\}$, $E = \{e_1, e_2\}$ and $\tau = \{\tilde{X}, \tilde{\phi}, (F_1, E), (F_2, E), (F_3, E)\}$ where $(F_1, E), (F_2, E), (F_3, E)$ are soft sets over X defined as follows:

$$F_1(e_1) = X, \quad F_1(e_2) = \{b\},$$

$$F_2(e_1) = \{a\}, \quad F_2(e_2) = X,$$

$$F_3(e_1) = \{a\}, \quad F_3(e_2) = \{b\}.$$

Then, τ defines a soft topology on X . Also, (X, τ, E) is soft b- T_1 -space, but it is not a soft b- T_2 -space, for $a, b \in X$ and $a \neq b$, but there is no b-open soft sets (F, E) and (G, E) such that $a \in (F, E)$, $b \in (G, E)$ and $(F, E) \tilde{\cap} (G, E) = \tilde{\phi}$.

(2) Let $X = \{a, b\}$, $E = \{e_1, e_2\}$ and $\tau = \{\tilde{X}, \tilde{\phi}, (F_1, E)\}$ where (F_1, E) is soft set over X defined as follows by $F_1(e_1) = X, \quad F_1(e_2) = \{b\}$.

Then τ defines a soft topology on X . Also (X, τ, E) is soft b- T_0 -space but not a soft b- T_1 -space, since $a, b \in X$, $a \neq b$, but all the b-open soft sets which contains a also contains b .

Theorem 3.5 A soft subspace (Y, τ_Y, E) of a soft b- T_2 -space (X, τ, E) is soft b- T_2 .

Proof. Let $x, y \in Y$ such that $x \neq y$. Then $x, y \in X$ such that $x \neq y$. Hence, there exist b-open soft sets (F, E) and (G, E) in X such that $x \in (F, E)$, $y \in (G, E)$ and $(F, E) \tilde{\cap} (G, E) = \tilde{\phi}$. It follows that, $x \in F(e), y \in G(e)$ and $F(e) \cap G(e) = \phi$ for all $e \in E$. This implies that, $x \in Y \cap F(e), y \in Y \cap G(e)$ and $F(e) \cap G(e) = \phi$ for all $e \in E$. Thus, $x \in \tilde{Y} \tilde{\cap} (F, E) = (F_Y, E)$, $y \in \tilde{Y} \tilde{\cap} (G, E) = (G_Y, E)$ and $(F_Y, E) \tilde{\cap} (G_Y, E) = \tilde{\phi}$, where $(F_Y, E), (G_Y, E)$ are b-open soft sets in Y . Therefore, (Y, τ_Y, E) is soft b- T_2 -space.

Definition 3.4. Let (X, τ, E) be a soft topological space, (G, E) be a b-closed soft set in X and $x \in X$ such that $x \notin (G, E)$. If there exist b-open soft sets (F_1, E) and (F_2, E) such that $x \in (F_1, E)$, $(G, E) \tilde{\subseteq} (F_2, E)$ and $(F_1, E) \tilde{\cap} (F_2, E) = \tilde{\phi}$, then (X, τ, E) is called a soft b-regular space. A soft b-regular T_1 -space is called a soft b- T_3 -space.

Proposition 3.3. Let (X, τ, E) be a soft topological space, (G, E) be a b-closed soft set in X and $x \in X$ such that $x \notin (G, E)$. If (X, τ, E) is soft b-regular space, then there exists a b-open soft set (F, E) such that $x \in (F, E)$ and $(F, E) \tilde{\cap} (G, E) = \tilde{\phi}$.

Proof. Obvious from Definition 3.4.

Proposition 3.4. Let (X, τ, E) be a soft topological space, $(F, E) \in SS(X)_E$ and $x \in X$. Then:

- (1) $x \in (F, E)$ if and only if $x_E \tilde{\subseteq} (F, E)$.
- (2) If $x_E \tilde{\cap} (F, E) = \tilde{\phi}$, then $x \notin (F, E)$.

Proof. Obvious.

Theorem 3.6. Let (X, τ, E) be a soft topological space and $x \in X$. If (X, τ, E) is soft b-regular space, then:

- (1) $x \notin (F, E)$ if and only if $x_E \tilde{\cap} (F, E) = \tilde{\phi}$ for every b-closed soft set (F, E) .
- (2) $x \notin (G, E)$ if and only if $x_E \tilde{\cap} (G, E) = \tilde{\phi}$ for every b-open soft set (G, E) .

Proof.

(1) Let (F, E) be a b-closed soft set such that $x \notin (F, E)$. Since (X, τ, E) is soft b-regular space. Then, by Proposition 3.3, there exists a b-open soft set (G, E) such that $x \in (G, E)$ and $(F, E) \tilde{\cap} (G, E) = \tilde{\phi}$. It follows that, $x_E \tilde{\subseteq} (G, E)$ from Proposition 3.4 (1). Hence, $x_E \tilde{\cap} (F, E) = \tilde{\phi}$. Conversely, if $x_E \tilde{\cap} (F, E) = \tilde{\phi}$, then $x \notin (F, E)$ from Proposition 3.4 (2).

(2) Let (G, E) be a b-open soft set such that $x \notin (G, E)$. If $x \notin G(e)$ for all $e \in E$, then we get the proof. If $x \notin G(e_1)$ and $x \in G(e_2)$ for some $e_1, e_2 \in E$, then $x \in G^c(e_1)$ and $x \notin G^c(e_2)$ for some $e_1, e_2 \in E$. This means that, $x_E \tilde{\cap} (G, E) \neq \tilde{\phi}$. Hence, $(G, E)^c$ is b-closed soft set such that $x \notin (G, E)^c$. It follows by (1) $x_E \tilde{\cap} (G, E)^c = \tilde{\phi}$. This implies that, $x_E \tilde{\subseteq} (G, E)$ and so $x \in (G, E)$, which is contradiction with $x \notin G(e_1)$ for some $e_1 \in E$. Therefore, $x_E \tilde{\cap} (G, E) = \tilde{\phi}$. Conversely, if $x_E \tilde{\cap} (G, E) = \tilde{\phi}$, then it obvious that $x \notin (G, E)$. This completes the proof.

Corollary 3.1. Let (X, τ, E) be a soft topological space and $x \in X$. If (X, τ, E) is soft b-regular space, then the following are equivalent:

- (1) (X, τ, E) is soft b- T_1 -space.
- (2) $\forall x, y \in X$ such that $x \neq y$, there exist b-open soft sets (F, E) and (G, E) such that $x_E \tilde{\subseteq} (F, E)$ and $y_E \tilde{\cap} (F, E) = \tilde{\phi}$ and $y_E \tilde{\subseteq} (G, E)$ and $x_E \tilde{\cap} (G, E) = \tilde{\phi}$.

Proof. Obvious from Theorem 3.6.

Theorem 3.7. Let (X, τ, E) be a soft topological space and $x \in X$. Then the following are equivalent:

- (1) (X, τ, E) is soft b-regular space.
- (2) For every b-closed soft set (G, E) such that $x_E \tilde{\cap} (G, E) = \tilde{\phi}$, there exist b-open soft sets (F_1, E) and (F_2, E) such that $x_E \tilde{\subseteq} (F_1, E)$, $(G, E) \tilde{\subseteq} (F_2, E)$ and $(F_1, E) \tilde{\cap} (F_2, E) = \tilde{\phi}$.

Proof.

- (1) \Rightarrow (2) Let (G, E) be a b-closed soft set such that $x_E \tilde{\cap} (G, E) = \tilde{\phi}$. Then $x \notin (G, E)$ from Theorem 3.6 (1). It follows by (1), there exist b-open soft sets (F_1, E) and (F_2, E) such that $x \in (F_1, E)$, $(G, E) \tilde{\subseteq} (F_2, E)$ and $(F_1, E) \tilde{\cap} (F_2, E) = \tilde{\phi}$. This means that, $x_E \tilde{\subseteq} (F_1, E)$, $(G, E) \tilde{\subseteq} (F_2, E)$ and $(F_1, E) \tilde{\cap} (F_2, E) = \tilde{\phi}$.
- (2) \Rightarrow (1) Let (G, E) be a b-closed soft set such that $x \notin (G, E)$. Then $x_E \tilde{\cap} (G, E) = \tilde{\phi}$ from Theorem 3.6 (1). It follows by (2), there exist b-open soft sets (F_1, E) and (F_2, E) such that $x_E \tilde{\subseteq} (F_1, E)$, $(G, E) \tilde{\subseteq} (F_2, E)$ and $(F_1, E) \tilde{\cap} (F_2, E) = \tilde{\phi}$. Hence, $x \in (F_1, E)$, $(G, E) \tilde{\subseteq} (F_2, E)$ and $(F_1, E) \tilde{\cap} (F_2, E) = \tilde{\phi}$. Thus, (X, τ, E) is soft b-regular space.

Theorem 3.8. Let (X, τ, E) be a soft topological space. If (X, τ, E) is soft b- T_3 -space, then $\forall x \in X$, x_E is b-closed soft set.

Proof. We want to prove that x_E is b-closed soft set, which is sufficient to prove that x_E^c is b-open soft set for

all $y \in \{x\}^c$. Since (X, τ, E) is soft $b-T_3$ -space, then there exist b -open soft sets $(F, E)_y$ and (G, E) such that $y_E \subseteq (F, E)_y$ and $x_E \cap (F, E)_y = \tilde{\phi}$ and $x_E \subseteq (G, E)$ and $y_E \cap (G, E) = \tilde{\phi}$. It follows that, $\bigcup_{y \in \{x\}^c} (F, E)_y \subseteq x_E^c$. Now, we want to prove that $x_E^c \subseteq \bigcup_{y \in \{x\}^c} (F, E)_y$. Let $\bigcup_{y \in \{x\}^c} (F, E)_y = (H, E)$, where $H(e) = \bigcup_{y \in \{x\}^c} F(e)_y$ for all $e \in E$. Since $x_E^c(e) = \{x\}^c$ for all $e \in E$ from Definition 2.6. So, for all $y \in \{x\}^c$ and $e \in E$, $x_E^c(e) = \{x\}^c = \bigcup_{y \in \{x\}^c} \{y\} = \bigcup_{y \in \{x\}^c} y_E(e) \subseteq \bigcup_{y \in \{x\}^c} F(e)_y = H(e)$. Thus, $x_E^c \subseteq \bigcup_{y \in \{x\}^c} (F, E)_y$ from Definition 2.2, and so $x_E^c = \bigcup_{y \in \{x\}^c} (F, E)_y$. This means that, x_E^c is b -open soft set for all $y \in \{x\}^c$. Therefore, x_E is b -closed soft set.

Theorem 3.9. Every soft $b-T_3$ -space is soft $b-T_2$ -space.

Proof. Let (X, τ, E) be a soft $b-T_3$ -space and $x, y \in X$ such that $x \neq y$. By Theorem 3.9, y_E is b -closed soft set and $x \notin y_E$. It follows from the soft b -regularity, there exist b -open soft sets (F_1, E) and (F_2, E) such that $x \in (F_1, E)$, $y_E \subseteq (F_2, E)$ and $(F_1, E) \cap (F_2, E) = \tilde{\phi}$. Thus, $x \in (F_1, E)$, $y \in y_E \subseteq (F_2, E)$ and $(F_1, E) \cap (F_2, E) = \tilde{\phi}$. Therefore, (X, τ, E) is soft $b-T_2$ -space.

Theorem 3.10. A soft subspace (Y, τ_Y, E) of a soft $b-T_3$ -space (X, τ, E) is soft $b-T_3$.

Proof. By Theorem 3.2, (Y, τ_Y, E) is soft $b-T_1$ -space. Now, we want to prove that (Y, τ_Y, E) is soft b -regular space. Let $y \in Y$ and (G, E) be a b -closed soft set in Y such that $y \notin (G, E)$. Then, $(G, E) = (Y, E) \cap (F, E)$ for some b -closed soft set (F, E) in X from Theorem 2.2. Hence, $y \notin (Y, E) \cap (F, E)$. But $y \in (Y, E)$, so $y \notin (F, E)$. Since (X, τ, E) is soft $b-T_3$ -space, so there exist b -open soft sets (F_1, E) and (F_2, E) in X such that $y \in (F_1, E)$, $(F, E) \subseteq (F_2, E)$ and $(F_1, E) \cap (F_2, E) = \tilde{\phi}$. Take $(G_1, E) = (Y, E) \cap (F_1, E)$ and $(G_2, E) = (Y, E) \cap (F_2, E)$, then $(G_1, E), (G_2, E)$ are b -open soft sets in Y such that $y \in (G_1, E)$, $(G, E) \subseteq (Y, E) \cap (F_2, E) = (G_2, E)$ and $(G_1, E) \cap (G_2, E) \subseteq (F_1, E) \cap (F_2, E) = \tilde{\phi}$. Thus, (Y, τ_Y, E) is soft $b-T_3$ -space.

Definition 3.5 Let (X, τ, E) be a soft topological space, $(F, E), (G, E)$ be b -closed soft sets in X such that $(F, E) \cap (G, E) = \tilde{\phi}$. If there exist b -open soft sets (F_1, E) and (F_2, E) such that $(F, E) \subseteq (F_1, E)$, $(G, E) \subseteq (F_2, E)$ and $(F_1, E) \cap (F_2, E) = \tilde{\phi}$, then (X, τ, E) is called a soft b -normal space. A soft b -normal T_1 -space is called a soft $b-T_4$ -space.

Theorem 3.11. Let (X, τ, E) be a soft topological space and $x \in X$. Then the following are equivalent:

- (1) (X, τ, E) is soft b -normal space.
- (2) For every b -closed soft set (F, E) and b -open soft set (G, E) such that $(F, E) \subseteq (G, E)$, there exists a b -open soft set (F_1, E) such that $(F, E) \subseteq (F_1, E)$, $bScl(F_1, E) \subseteq (G, E)$.

Proof.

- (1) \Rightarrow (2) Let (F, E) be a b -closed soft set and (G, E) be a b -open soft set such that $(F, E) \subseteq (G, E)$. Then

$(F, E), (G, E)^c$ are b -closed soft sets such that $(F, E) \cap (G, E)^c = \tilde{\phi}$. It follows by (1), there exist b -open soft sets (F_1, E) and (F_2, E) such that $(F, E) \subseteq (F_1, E)$, $(G, E)^c \subseteq (F_2, E)$ and $(F_1, E) \cap (F_2, E) = \tilde{\phi}$. Now, $(F_1, E) \subseteq (F_2, E)^c$, so $bScl(F_1, E) \subseteq bScl(F_2, E)^c = (F_2, E)^c$, where (G, E) is b -open soft set. Also $(F_2, E)^c \subseteq (G, E)$. Hence, $bScl(F_1, E) \subseteq (F_2, E)^c \subseteq (G, E)$. Thus, $(F, E) \subseteq (F_1, E)$, $bScl(F_1, E) \subseteq (G, E)$.

- (2) \Rightarrow (1) Let $(G_1, E), (G_2, E)$ be b -closed soft sets such that $(G_1, E) \cap (G_2, E) = \tilde{\phi}$. Then $(G_1, E) \subseteq (G_2, E)^c$, then by hypothesis, there exists a b -open soft set (F_1, E) such that $(G_1, E) \subseteq (F_1, E)$, $bScl(F_1, E) \subseteq (G_2, E)^c$. So $(G_2, E) \subseteq [bScl(F_1, E)]^c$, $(G_1, E) \subseteq (F_1, E)$ and $[bScl(F_1, E)]^c \cap (F_1, E) = \tilde{\phi}$, where (F_1, E) and $[bScl(F_1, E)]^c$ are b -open soft sets. Thus, (X, τ, E) is soft b -normal space.

Theorem 3.12. A b -closed soft subspace (Y, τ_Y, E) of a soft b -normal space (X, τ, E) is soft b -normal.

Proof. Let $(G_1, E), (G_2, E)$ be b -closed soft sets in Y such that $(G_1, E) \cap (G_2, E) = \tilde{\phi}$. Then $(G_1, E) = (Y, E) \cap (F_1, E)$ and $(G_2, E) = (Y, E) \cap (F_2, E)$ for some b -closed soft sets $(F_1, E), (F_2, E)$ in X from Theorem 2.2. Since Y is a b -closed soft subset of X . Then $(G_1, E), (G_2, E)$ are b -closed soft sets in X such that $(G_1, E) \cap (G_2, E) = \tilde{\phi}$. Hence, by soft b -normality there exist b -open soft sets (H_1, E) and (H_2, E) such that $(G_1, E) \subseteq (H_1, E)$, $(G_2, E) \subseteq (H_2, E)$ and $(H_1, E) \cap (H_2, E) = \tilde{\phi}$. Since $(G_1, E), (G_2, E) \subseteq (Y, E)$, then $(G_1, E) \subseteq (Y, E) \cap (H_1, E)$, $(G_2, E) \subseteq (Y, E) \cap (H_2, E)$ and $[(Y, E) \cap (H_1, E)] \cap [(Y, E) \cap (H_2, E)] = \tilde{\phi}$, where $(Y, E) \cap (H_1, E)$ and $(Y, E) \cap (H_2, E)$ are b -open soft sets in Y . Therefore, (Y, τ_Y, E) is a soft b -normal space.

Theorem 3.13. Let (X, τ, E) be a soft topological space. If (X, τ, E) is soft b -normal space and x_E is b -closed soft set in τ for all $x \in X$, then (X, τ, E) is soft $b-T_3$ -space.

Proof. Since x_E is b -closed soft set for all $x \in X$, then (X, τ, E) is soft $b-T_1$ -space from Theorem 3.3. Also (X, τ, E) is soft b -regular space from Theorem 3.7 and Definition 3.5. Hence, (X, τ, E) is soft $b-T_3$ -space.

4 Irresolute b -open soft functions

Theorem 4.1. Let (X, τ_1, A) and (Y, τ_2, B) be soft topological spaces and $f_{pu} : SS(X)_A \rightarrow SS(Y)_B$ be a soft function which is bijective and irresolute b -open soft. If (X, τ_1, A) is soft $b-T_0$ -space, then (Y, τ_2, B) is also a soft $b-T_0$ -space.

Proof. Let $y_1, y_2 \in Y$ such that $y_1 \neq y_2$. Since f_{pu} is surjective, then $\exists x_1, x_2 \in X$ such that $u(x_1) = y_1$, $u(x_2) = y_2$ and $x_1 \neq x_2$. By hypothesis, there exist b -open soft sets (F, A) and (G, A) in X such that either $x_1 \in (F, A)$ and $x_2 \notin (F, A)$, or $x_2 \in (G, A)$ and $x_1 \notin (G, A)$. So, either $x_1 \in F_A(e)$ and $x_2 \notin F_A(e)$ or $x_2 \in G_A(e)$ and $x_1 \notin G_A(e)$

for all $e \in A$. This implies that, either $y_1 = u(x_1) \in u[F_A(e)]$ and $y_2 = u(x_2) \notin u[F_A(e)]$ or $y_2 = u(x_2) \in u[G_A(e)]$ and $y_1 = u(x_1) \notin u[G_A(e)]$ for all $e \in A$. Hence, either $y_1 \in f_{pu}(F,A)$ and $y_2 \notin f_{pu}(F,A)$ or $y_2 \in f_{pu}(G,A)$ and $y_1 \notin f_{pu}(G,A)$. Since f_{pu} is irresolute b-open soft function, then $f_{pu}(F,A), f_{pu}(G,A)$ are b-open soft sets in Y . Hence, (Y, τ_2, B) is also a soft b- T_0 -space.

Theorem 4.2 Let (X, τ_1, A) and (Y, τ_2, B) be soft topological spaces and $f_{pu} : SS(X)_A \rightarrow SS(Y)_B$ be a soft function which is bijective and irresolute b-open soft. If (X, τ_1, A) is soft b- T_1 -space, then (Y, τ_2, B) is also a soft b- T_1 -space.

Proof. It is similar to the proof of Theorem 4.1. **Theorem**

4.3. Let (X, τ_1, A) and (Y, τ_2, B) be soft topological spaces and $f_{pu} : SS(X)_A \rightarrow SS(Y)_B$ be a soft function which is bijective and irresolute b-open soft. If (X, τ_1, A) is soft b- T_2 -space, then (Y, τ_2, B) is also a soft b- T_2 -space.

Proof. $y_1, y_2 \in Y$ such that $y_1 \neq y_2$. Since f_{pu} is surjective, then $\exists x_1, x_2 \in X$ such that $u(x_1) = y_1, u(x_2) = y_2$ and $x_1 \neq x_2$. By hypothesis, there exist b-open soft sets (F,A) and (G,A) in X such that $x_1 \in (F,A), x_2 \in (G,A)$ and $(F,A) \tilde{\cap} (G,A) = \tilde{\phi}_A$. So $x_1 \in F_A(e), x_2 \in G_A(e)$ and $F_A(e) \tilde{\cap} G_A(e) = \phi$ for all $e \in A$. This implies that, $y_1 = u(x_1) \in u[F_A(e)], y_2 = u(x_2) \in u[G_A(e)]$ for all $e \in A$. Hence, $y_1 \in f_{pu}(F,A), y_2 \in f_{pu}(G,A)$ and $f_{pu}(F,A) \tilde{\cap} f_{pu}(G,A) = f_{pu}[(F,A) \tilde{\cap} (G,A)] = f_{pu}[\tilde{\phi}_A] = \tilde{\phi}_B$ from Theorem 2.3. Since f_{pu} is irresolute b-open soft function, then $f_{pu}(F,A), f_{pu}(G,A)$ are b-open soft sets in Y . Thus, (Y, τ_2, B) is also a soft b- T_2 -space.

Theorem 4.4. Let (X, τ_1, A) and (Y, τ_2, B) be soft topological spaces and $f_{pu} : SS(X)_A \rightarrow SS(Y)_B$ be a soft function which is bijective, b-irresolute soft and irresolute b-open soft. If (X, τ_1, A) is soft b-regular space, then (Y, τ_2, B) is also a soft b-regular space.

Proof. Let (G,B) be a b-closed soft set in Y and $y \in Y$ such that $y \notin (G,B)$. Since f_{pu} is surjective and b-irresolute soft, then $\exists x \in X$ such that $u(x) = y$ and $f_{pu}^{-1}(G,B)$ is b-closed soft set in X such that $x \notin f_{pu}^{-1}(G,B)$. By hypothesis, there exist b-open soft sets (F,A) and (H,A) in X such that $x \in (F,A), f_{pu}^{-1}(G,B) \tilde{\subseteq} (H,A)$ and $(F,A) \tilde{\cap} (H,A) = \tilde{\phi}_A$. It follows that, $x \in F_A(e)$ for all $e \in A$ and $(G,B) = f_{pu}[f_{pu}^{-1}(G,B)] \tilde{\subseteq} f_{pu}(H,A)$ from Theorem 2.3. So, $y = u(x) \in u[F_A(e)]$ for all $e \in A$ and $(G,B) \tilde{\subseteq} f_{pu}(H,A)$. Hence, $y \in f_{pu}(F,A)$ and $(G,B) \tilde{\subseteq} f_{pu}(H,A)$ and $f_{pu}(F,A) \tilde{\cap} f_{pu}(H,A) = f_{pu}[(F,A) \tilde{\cap} (H,A)] = f_{pu}[\tilde{\phi}_A] = \tilde{\phi}_B$ from Theorem 2.3. Since f_{pu} is irresolute b-open soft function. Then, $f_{pu}(F,A), f_{pu}(H,A)$ are b-open soft sets in Y . Thus, (Y, τ_2, B) is also a soft b-regular space.

Theorem 4.5. Let (X, τ_1, A) and (Y, τ_2, B) be soft topological spaces and $f_{pu} : SS(X)_A \rightarrow SS(Y)_B$ be a soft function which is bijective, b-irresolute soft and irresolute b-open soft. If (X, τ_1, A) is soft b- T_3 -space, then (Y, τ_2, B) is also a soft b- T_3 -space.

Proof. Since (X, τ_1, A) is soft b- T_3 -space, then (X, τ_1, A) is soft b-regular T_1 -space. It follows that, (Y, τ_2, B) is also a soft b- T_1 -space from Theorem 4.2 and soft b-regular space from Theorem 4.4. Hence, (Y, τ_2, B) is also a soft b- T_3 -space.

Theorem 4.6. Let (X, τ_1, A) and (Y, τ_2, B) be soft topological spaces and $f_{pu} : SS(X)_A \rightarrow SS(Y)_B$ be a soft function which is bijective, b-irresolute soft and irresolute b-open soft. If (X, τ_1, A) is soft b-normal space, then (Y, τ_2, B) is also a soft b-normal space.

Proof. Let $(F,B), (G,B)$ be b-closed soft sets in Y such that $(F,B) \tilde{\cap} (G,B) = \tilde{\phi}_B$. Since f_{pu} is b-irresolute soft, then $f_{pu}^{-1}(F,B)$ and $f_{pu}^{-1}(G,B)$ are b-closed soft set in X such that $f_{pu}^{-1}(F,B) \tilde{\cap} f_{pu}^{-1}(G,B) = f_{pu}^{-1}[(F,B) \tilde{\cap} (G,B)] = f_{pu}^{-1}[\tilde{\phi}_B] = \tilde{\phi}_A$ from Theorem 2.3. By hypothesis, there exist irresolute b-open soft sets (K,A) and (H,A) in X such that $f_{pu}^{-1}(F,B) \tilde{\subseteq} (K,A), f_{pu}^{-1}(G,B) \tilde{\subseteq} (H,A)$ and $(F,A) \tilde{\cap} (H,A) = \tilde{\phi}_A$. It follows that, $(F,B) = f_{pu}[f_{pu}^{-1}(F,B)] \tilde{\subseteq} f_{pu}(K,A)$ and $(G,B) = f_{pu}[f_{pu}^{-1}(G,B)] \tilde{\subseteq} f_{pu}(H,A)$ from Theorem 2.3 and $f_{pu}(K,A) \tilde{\cap} f_{pu}(H,A) = f_{pu}[(K,A) \tilde{\cap} (H,A)] = f_{pu}[\tilde{\phi}_A] = \tilde{\phi}_B$ from Theorem 2.3. Since f_{pu} is irresolute b-open soft function. Then $f_{pu}(K,A), f_{pu}(H,A)$ are b-open soft sets in Y . Thus, (Y, τ_2, B) is also a soft b-normal space.

Corollary 4.1. Let (X, τ_1, A) and (Y, τ_2, B) be soft topological spaces and $f_{pu} : SS(X)_A \rightarrow SS(Y)_B$ be a soft function which is bijective, b-irresolute soft and irresolute b-open soft. If (X, τ_1, A) is soft b- T_4 -space, then (Y, τ_2, B) is also a soft b- T_4 -space.

Proof. It is obvious from Theorem 4.2 and Theorem 4.6.

5 Conclusion

Recently, many scientists have studied the soft set theory, which is initiated by Molodtsov [33] and easily applied to many problems having uncertainties from social life. In the present work, we introduce the notion of soft b-separation axioms. In particular we study the properties of the soft b-regular spaces and soft b-normal spaces. We show that, if x_E is b-closed soft set for all $x \in X$ in a soft topological space (X, τ, E) , then (X, τ, E) is soft b- T_1 -space. Also, we show that if a soft topological space (X, τ, E) is soft b- T_3 -space, then $\forall x \in X, x_E$ is b-closed soft set. Also, we show that the property of being b- T_i -spaces ($i = 1, 2$) is soft topological property under a bijection and irresolute b-open soft mapping. Further, the properties of being soft b-regular and soft b-normal are soft topological properties under a bijection, b-irresolute soft and irresolute b-open soft functions. Finally, we show that the property of being b- T_i -spaces ($i = 1, 2, 3, 4$) is a hereditary property. We hope that, the results in this paper will help researcher enhance and promote the further study on soft topology to carry out a general framework for their applications in practical life.

The authors are grateful to the anonymous referee for a careful checking of the details and for helpful comments that improved this paper.

References

- [1] A. M. Abd El-latif and Serkan Karatas, Supra b-open soft sets and supra b-soft continuity on soft topological spaces, *Journal of Mathematics and Computer Applications Research*, 5 (2015) 1-18.
- [2] B. Ahmad and A. Kharal, Mappings on fuzzy soft classes, *Adv. Fuzzy Syst.* 2009, Art. ID 407890, 6 pp.
- [3] B. Ahmad and A. Kharal, Mappings on soft classes, *New Math. Nat. Comput.*, 7 (3) (2011) 471-481.
- [4] B. Ahmad and A. Kharal, On fuzzy soft sets, *Adv. Fuzzy Syst.* 2009, Art. ID 586507, 6 pp.
- [5] H. Aktas and N. Cagman, Soft sets and soft groups, *Information Sciences*, 1 (77)(2007) 2726-2735.
- [6] M. I. Ali, F. Feng, X. Liu, W. K. Min and M. Shabir, On some new operations in soft set theory, *Computers and Mathematics with Applications*, 57 (2009) 1547-1553.
- [7] S. Atmaca and I.Zorlutuna, On fuzzy soft topological spaces, *Annals of Fuzzy Mathematics and Informatics*, 5 (2013) 377-386.
- [8] Bakir Tanay and M. Burel Kandemir, Topological structure of fuzzy soft sets, *Computer and mathematics with applications*, (61)(2011) 2952-2957.
- [9] N. Cagman, F. Citak and S. Enginoglu, Fuzzy parameterized fuzzy soft set theory and its applications, *Turkish Journal of Fuzzy Systems*, 1(1)(2010) 21-35.
- [10] N. Cagman and S. Enginoglu, Soft set theory and uni-Fint decision making, *European Journal of Operational Research*, 207 (2010) 848-855.
- [11] C. L. Change, Fuzzy topological spaces, *J. Math. Anal. Appl.*, 24 (1968), 182-190.
- [12] S. A. El-Sheikh and A. M. Abd El-latif, Characterization of b-open soft sets in soft topological spaces, *Journal of New Theory*, 2 (2015) 8-18.
- [13] S. A. El-Sheikh and A. M. Abd El-latif, Decompositions of some types of supra soft sets and soft continuity, *International Journal of Mathematics Trends and Technology*, 9 (1) (2014) 37-56.
- [14] S. Hussain and B. Ahmad, Some properties of soft topological spaces, *Comput. Math. Appl.*, 62 (2011) 4058-4067.
- [15] Jianyu Xiao, Minming Tong, Qi Fan and Su Xiao, Generalization of Belief and Plausibility Functions to Fuzzy Sets, *Applied Mathematics Information Sciences*, 6 (2012) 697-703.
- [16] A.Kandil, O. A. E. Tantawy, S. A. El-Sheikh and A. M. Abd El-latif, Fuzzy soft semi connected properties in fuzzy soft topological spaces, *Math. Sci. Lett.*, 4 (2015) 171-179.
- [17] A.Kandil, O. A. E. Tantawy, S. A. El-Sheikh and A. M. Abd El-latif, γ -operation and decompositions of some forms of soft continuity in soft topological spaces, *Ann. Fuzzy Math. Inform.*, 7 (2014) 181-196.
- [18] A.Kandil, O. A. E. Tantawy, S. A. El-Sheikh and A. M. Abd El-latif, γ -operation and decompositions of some forms of soft continuity of soft topological spaces via soft ideal, *Ann. Fuzzy Math. Inform.*, 9 (3) (2015) 385-402.
- [19] A.Kandil, O. A. E. Tantawy, S. A. El-Sheikh and A. M. Abd El-latif, Soft connectedness via soft ideals, *Journal of New Results in Science*, 4 (2014) 90-108.
- [20] A.Kandil, O. A. E. Tantawy, S. A. El-Sheikh and A. M. Abd El-latif, Soft ideal theory, Soft local function and generated soft topological spaces, *Appl. Math. Inf. Sci.*, 8 (4) (2014) 1595-1603.
- [21] A.Kandil, O. A. E. Tantawy, S. A. El-Sheikh and A. M. Abd El-latif, Soft regularity and normality based on semi open soft sets and soft ideals, *Appl. Math. Inf. Sci. Lett.*, 3 (2015) 47-55.
- [22] A.Kandil, O. A. E. Tantawy, S. A. El-Sheikh and A. M. Abd El-latif, Soft semi compactness via soft ideals, *Appl. Math. Inf. Sci.*, 8 (5) (2014) 2297-2306.
- [23] A.Kandil, O. A. E. Tantawy, S. A. El-Sheikh and A. M. Abd El-latif, Soft semi (quasi) Hausdorff spaces via soft ideals, *South Asian J. Math.*, 4 (6) (2014) 265-284.
- [24] A.Kandil, O. A. E. Tantawy, S. A. El-Sheikh and A. M. Abd El-latif, Soft semi separation axioms and irresolute soft functions, *Ann. Fuzzy Math. Inform.*, 8 (2) (2014) 305-318.
- [25] A.Kandil, O. A. E. Tantawy, S. A. El-Sheikh and A. M. Abd El-latif, Some fuzzy soft topological properties based on fuzzy semi open soft sets, *South Asian J. Math.*, 4 (4) (2014) 154-169.
- [26] A.Kandil, O. A. E. Tantawy, S. A. El-Sheikh and A. M. Abd El-latif, Supra generalized closed soft sets with respect to an soft ideal in supra soft topological spaces, *Appl. Math. Inf. Sci.*, 8 (4) (2014) 1731-1740.
- [27] D. V. Kovkov, V. M. Kolbanov and D. A. Molodtsov, Soft sets theory-based optimization, *Journal of Computer and Systems Sciences Finternational* 46 (6) (2007) 872-880.
- [28] J. Mahanta and P.K. Das, Results on fuzzy soft topological spaces, *arXiv:1203.0634v1*, 2012.
- [29] P. K. Maji, R. Biswas and A. R. Roy, Fuzzy soft sets, *Journal of Fuzzy Mathematics*, 9 (3) (2001) 589-602.
- [30] P. K. Maji, R. Biswas and A. R. Roy, intuitionistic fuzzy soft sets, *Journal of Fuzzy Mathematics*, 9 (3) (2001) 677-691.
- [31] P. K. Maji, R. Biswas and A. R. Roy, Soft set theory, *Computers and Mathematics with Applications*, 45 (2003) 555-562.
- [32] P. Majumdar and S. K. Samanta, Generalised fuzzy soft sets, *Computers and Mathematics with Applications*, 59 (2010) 1425-1432.
- [33] D. A. Molodtsov, Soft set theory-firs results, *Computers and Mathematics with Applications*, 37 (1999) 19-31.
- [34] D.Molodtsov, V. Y. Leonov and D. V. Kovkov, Soft sets technique and its application, *Nechetkie Sistemy i Myagkie Vychisleniya*, 1 (1) (2006) 8-39.
- [35] A. Mukherjee and S. B. Chakraborty, On Fintuitionistic fuzzy soft relations, *Bulletin of Kerala Mathematics Association*, 5 (1) (2008) 35-42.
- [36] B. Pazar Varol and H. Aygun, Fuzzy soft topology, *Hacettepe Journal of Mathematics and Statistics*, 41 (3) (2012) 407-419.
- [37] D. Pei and D. Miao, From soft sets to information systems, in: X. Hu, Q. Liu, A. Skowron, T. Y. Lin, R. R. Yager, B. Zhang (Eds.), *Proceedings of Granular Computing*, in: IEEE, vol.2, 2005, pp. 617-621.
- [38] S. Roy and T. K. Samanta, A note on fuzzy soft topological spaces, *Annals of Fuzzy Mathematics and informatics*, 3 (2) (2012) 305-311.

[39] S. Roy and T. K. Samanta, An introduction to open and closed sets on fuzzy topological spaces, *Annals of Fuzzy Mathematics and Informatics*, 6 (2) (2012) 425-431.

[40] M. Shabir and M. Naz, On soft topological spaces, *Comput. Math. Appl.*, 61 (2011) 1786-1799.

[41] B. Tanay and M. B. Kandemir, Topological structure of fuzzy soft sets, *Computer and Math. with appl.*, 61 (2011) 412-418.

[42] Weijian Rong, The countabilities of soft topological spaces, *International Journal of Computational and Mathematical Sciences*, 6 (2012) 159-162.

[43] Won Keun Min, A note on soft topological spaces, *Computers and Mathematics with Applications*, 62 (2011) 3524-3528.

[44] Z. Xiao, L. Chen, B. Zhong and S. Ye, Recognition for soft information based on the theory of soft sets, in: J. Chen (Ed.), *Proceedings of ICSSM-05*, vol. 2, IEEE, 2005, pp. 1104-1106.

[45] Yong Chan Kim and Jung Mi Ko, Fuzzy G-closure operators, *commun Korean Math. Soc.*, 18 (2)(2008) 325-340.

[46] L. A. Zadeh, Fuzzy sets, *Information and Control*, 8 (1965) 338-353.

[47] Y. Zou and Z. Xiao, Data analysis approaches of soft sets under incomplete information, *Knowledge-Based Systems*, 21 (2008) 941-945.

[48] I. Zorlutuna, M. Akdag, W.K. Min and S. Atmaca, Remarks on soft topological spaces, *Annals of Fuzzy Mathematics and Informatics*, 3 (2012) 171-185.



Rodyna Hosny is a Lecturer of pure mathematics of Department of Mathematics at University of Zagazig. she received the PhD degree in Pure Mathematics. Her research interests are in the areas of pure mathematics such as General topology, Soft topology, Rough sets, and Fuzzy topology. She is referee of several international journals in the frame of pure mathematics. She has published research articles in various international journals.



Alaa Mohamed Abd El-Latif received the PhD degree in pure Mathematics (Topology) Ain Shams University, Faculty of Education, Mathematic Department, Cairo, Egypt. His primary research areas are General Topology, Fuzzy Topology, Set theory, Soft set theory and Soft topology. He

is referee of several international journals in the pure mathematics. Dr. Alaa has published many papers in refereed journals.



Sobhy Ahmed Aly El-Sheikh is a Professor of pure Mathematics, Ain Shams University ,Faculty of Education, Mathematic Department, Cairo, Egypt. He born in 1955. He received the Ph.D. degree in Topology from the University of Zagazig. His primary

research areas are General Topology, Fuzzy Topology, double sets and theory of sets. Dr. Sobhy has published over 15 papers in *Fuzzy set and system Journal (FSS)*, *Information science Journal (INFS)*, *Journal of fuzzy Mathematics* and *Egyptian Journal of Mathematical Society*. He was the Supervisor of many PHD and MSC Thesis.