

2021

Risk Factors Of Preterm Birth of Neonates Attended Al- Mukalla Maternity and Childhood Hospital ,Yemen

Hanan Saeed Bin Gouth

College of Medicine and Health Sciences, Hadhramout University

Saleh Awad bahwel

College of Science, Hadhramout University

Mazin Ahmed Jawass

College of Science, Hadhramout University

Ali Salim Bahartha

College of Medicine & Health sciences; Hadhramout University

Fauzia Faraj Bamatraf

College of Medicine & Health sciences, Hadhramout University

Follow this and additional works at: https://digitalcommons.aaru.edu.jo/huj_nas



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Bin Gouth, Hanan Saeed; bahwel, Saleh Awad; Jawass, Mazin Ahmed; Bahartha, Ali Salim; and Bamatraf, Fauzia Faraj (2021) "Risk Factors Of Preterm Birth of Neonates Attended Al- Mukalla Maternity and Childhood Hospital ,Yemen," *Hadhramout University Journal of Natural & Applied Sciences*: Vol. 14 : Iss. 1 , Article 1.

Available at: https://digitalcommons.aaru.edu.jo/huj_nas/vol14/iss1/1

This Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in Hadhramout University Journal of Natural & Applied Sciences by an authorized editor. The journal is hosted on [Digital Commons](#), an Elsevier platform. For more information, please contact rakan@aar.edu.jo, marah@aar.edu.jo, u.murad@aar.edu.jo.

Risk Factors Of Preterm Birth of Neonates Attended Al- Mukalla Maternity and Childhood Hospital ,Yemen

Hanan Saeed Bin Gouth* Saleh Awad bahwel* Mazin Ahmed Jawass*
Ali Salim Bahartha* Fauzia Faraj Bamatraf **

Abstract

The preterm birth continues to be the leading cause of prenatal morbidity and mortality. Neonates born preterm are known to have a certain added risk of death, disease, and disability. The aim of the present study is to identify risk factors associated with preterm birth of neonates in Al-Mukalla Maternity and Childhood Hospital (MCH). Retrospective case-control study at a ratio of 1:1 was conducted in the neonatal unit of Pediatric Ward from October 2012 to October 2013, cases and controls data were collected from medical records. A total of 104 cases and 104 controls were included in the study. The results showed several risk factors significantly associated with preterm birth of neonates which are: bad obstetric history (BOH), with p value 0.014, present maternal diseases including hypertension, pre or/and - eclampsia, urinary tract infection and genital infection (p value= 0.003, 0.002, 0.045, 0.002 respectively) as well as presence of twins and antipartum hemorrhage (p value= 0.000, 0.028 respectively). We concluded that the most common risk factors of preterm birth of neonate were BOH, Maternal diseases in current pregnancy as well as presence of twins and antipartum hemorrhage. It is necessary to improve prenatal care for pregnant women which may decrease the potential of preterm birth of neonates.

Key words: Preterm neonate, Risk factors, Al-Mukalla, Yemen.

Introduction:

Preterm birth (premature birth) is defined by the World Health Organization (WHO) as birth of an infant prior to 37 weeks' (259 days') completed gestation [23]. It continues to be the leading cause of perinatal morbidity and mortality [9]. Infants born preterm are known to have a certain added risk of death, disease, disability, as well as longer-term motor, cognitive, visual, hearing, behavioral, social-emotional, health, and growth problems compared with normal-term infants [17,30].

Based on global reports, 60%–80% of neonatal mortalities (not accompanied by congenital abnormalities) occur in premature infants, resulting in asphyxia in the first week and septicemia in the fourth week [6]. Previous studies have estimated that 11.1% of all live births are preterm worldwide, ranging from about 5% in several European countries to 18% in some African countries [3].

The underlying causes of preterm delivery are multiple and poorly understood. It may include individual-level behavioral and psychosocial factors, neighborhood characteristics, environmental exposures, medical conditions, infertility treatments, biological factors, and genetics [2], many of these factors occurs in combination.

Studies have revealed that mothers' education, age over 36, history of having a premature infant, multiparity, mother hypertension, infant diabetes, oligohydramnios polyhydramnios, placenta previa, anatomic abnormality of uterus, history of organic disorder (cardiac, renal, thyroid), and blood group type A have significant correlation with recurrence of premature labor [7,21].

In other studies, the most prevalent risk factors leading to premature birth consist of insufficient pregnancy care (52%), mothers' age being under 20 (34.7%), third-trimester hemorrhage (23.4%), and eclampsia and preeclampsia (13.1%) [11,27], and the most prevalent complications of preterm labor reported in infants include septicemia (66.7%), hyperbilirubinemia (58.8%), asphyxia (26.8%), and complications regarding the hyaline membrane (23.3%) [11,27]. The aim of the present study is to identify the risk factors associated with preterm birth of neonates (in Al-Mukalla Maternity and Childhood Hospital (MCH)).

Method and Materials:

This study was designed as retrospective case-control study, at a ratio of 1:1. It was conducted in neonatal unit of the pediatric ward in Al-Mukalla Maternity and Childhood Hospital (MCH) in Al-Mukalla city the capital of Hadhramout Governorate, Yemen. This Hospital is a tertiary care teaching hospital responsible for the care of all pediatric and mother patients coming from three Governorates; Hadhramout, Shabowa and Al-Mahra.

*Pediatric department, College of Medicine & Health sciences; Hadhramout University (HUCOM).

** Community Medicine department; College of Medicine & Health sciences; Hadhramout University (HUCOM). Received on 3/4/2017 and Accepted for Publication on 17/5/2017

The data of cases and controls were collected through medical records during one year period from October 2012 to October 2013. Cases were defined as preterm live neonates (29–<37 weeks). Controls were defined as a full-term live neonates (≥ 37 weeks). In general, pre-term cases were diagnosed in advance by the resident specialist in the hospital. The study was included 104 preterm live neonates who are registered during the study period (one year) and equal numbers of full term live neonates as control, registered during the same period.

The risk factors studied in this study were: Sex of baby, bad obstetric history (BOH) of mother including: {previous abortion, preterm labor,} parity (number of pregnancy primary or multi), maternal systemic diseases (Maternal diseases in current pregnancy) including {hypertension, preeclampsia-eclampsia, diabetes mellitus (DM), heart, renal diseases, anemia, urinary tract infection, and genital infection} and localized comorbidities including { history of twins, anti-partum hemorrhage, premature rupture of membrane (PROM), uterine anomalies, cervical incompetence,

polyhydramnios and oligohydramnios.} The study was approved by the Ethics Committee of the hospital of child and mother in Al-Mukalla city.

Statistical Analysis

The collected data were entered into a computer and analyzed using SPSS version 20.0. Frequencies and percentage were calculated and presented in tables and graphs using Excel and Word programs. Chi square test was used to determine the association between preterm birth of neonate and potential predictor factors and statistical significance was achieved when at $P < 0.05$.

Results:

A total of 104 cases and same number for controls were included in the study. The male: female ratio was 1.1:1 with mild male predominance, without any significant association ($p =$ value 0.890). The study demonstrated that there were statistically significant association between the case preterm birth of neonate and the bad obstetric history of mother (BOH), with p value = 0.014 (Table 1)

Table 1: Association between preterm birth of neonates and Sex, BOH and Parity factors

Factors		Case N=104 F (%)		Control N=104 F (%)		P value
Sex	Male	55	52.9	54	51.9	0.890
	Female	49	47.1	50	48.1	
BOH	Yes	23	22.1	10	9.6	0.014
	No	81	77.9	94	90.4	
Parity	Yes	31	29.8	44	42.3	0.060
	No	73	70.2	60	57.7	

The association between preterm birth of neonates and present maternal systemic diseases factors are shown in Table No 2. The study showed that there were statistically significant association between the case preterm birth of neonate and the present maternal systemic diseases including hypertension $p =$ value 0.003, preeclampsia-eclampsia p value = 0

.002, urinary tract infection $p =$ value 0.045, genital infection p value = 0.002, twins p value = 0.000 and antipartum hemorrhage with p value = 0.028. While other factors (parity, D.M, heart and renal diseases, anemia, PROM, uterine anomalies, cervical incompetence, and oligohydramnios) were not significant as factor for preterm birth of neonate.

Table 2: Association between preterm birth of neonates and present maternal systemic diseases factors

Factors		Cases N=104 F (%)		Controls N=104 F (%)		P value
Preeclampsia-eclampsia	Yes	26	25	9	8.7	0.002
	No	78	75	95	91.3	
Hypertension	Yes	11	10.6	1	0.9	0.003
	No	93	89.4	103	99.1	
D.M	Yes	11	10.6	5	4.8	0.118
	No	93	89.4	99	95.2	
Heart disease	Yes	1	0.9	2	1.9	0.561
	No	103	99.1	102	98.1	
Renal disease	Yes	0.0	0.0	3	2.9	0.081
	No	104	100	101	97.1	
Anemia	Yes	7	6.7	2	1.9	0.088
	No	97	93.3	102	98.1	
UTI	Yes	29	27.9	17	16.3	0.045
	No	75	72.1	87	83.7	
Genital infection	Yes	26	25	9	8.7	0.002
	No	78	75	95	91.3	

As shown in Table 3 there are significant association between preterm birth of neonate and the history of Twins and anti-partum

hemorrhage(p value = 0.000 and 0.028 respectively).

Table 3: Association between preterm birth of neonates and localized obstetric factors

Factors		Cases N=104 F (%)		Controls N=104 F (%)		P value
Twins	Yes	23	22.1	3	2.9	0.000
	No	81	77.9	101	97.1	
PROM	Yes	28	26.9	25	24.0	0.633
	No	76	73.1	79	76.0	

Anti-partum hemorrhage	Yes	20	19.2	9	8.7	0.028
	No	84	80.8	95	91.3	
Uterine anomalies	Yes	3	0.9	0	0.0	0.081
	No	101	99.1	104	100.0	
Cervical incompetence	Yes	2	1.9	1	0.9	0.561
	No	102	98.1	103	99.1	
Polyhydramnios	Yes	8	7.7	2	1.9	0.052
	No	96	92.3	102	98.1	
Oligohydramnios.	Yes	6	5.8	6	5.8	1.000
	No	98	94.2	98	94.2	

Discussion:

Preterm birth is a prevalent obstetric complication associated with significant neonatal mortality and morbidity worldwide. Addressing the burden of preterm birth in developing countries is of public health importance due to its high (9 to 16%) prevalence, though the exact etiopathogenesis of preterm birth is still unclear. In developing nations, prediction and/or diagnosis of this multifactorial process is made mainly based on the evidence reported in the western literature on the risk factors and probable pathological mechanisms [5,10,32]. Our study was designed as a matched case-control study, aimed to determine the risk factors of preterm birth of neonate in Al-Mukalla MCH.

The present study found that bad obstetric history (previous abortion, or preterm labor) was significantly associated with preterm birth of a neonate, and this corresponding with many studies [9,32,24, 29]. No significant association was observed in this study between preterm birth of neonate and parity. Some cross-sectional analyses have reported no effect of parity on the occurrence of preterm birth of a neonate [24, 4]. While others showed an association with high parity [15].

Systemic and localized maternal comorbidities factors were studied in the present study and it was found that there were significant associations of some of them. Maternal eclampsia and pregnancy hypertension is a status that is present in 5%–7% of all deliveries, and is

correlated with main fetal disease and premature labor [13]. This study showed significant association between preterm birth of neonate and pre or/ and eclampsia and hypertension which consistent with many previous studies [9,31,4,15,20,1,22].

Maternal Diabetes mellitus was found without any significant association with preterm birth of neonate and this finding corresponding with some studies [4,24] while many studies observed significant association [31,32,25,26].

Cardiovascular and renal disease factors did not show any significant association with preterm birth of neonates in the current study, as opposed to other studies and this in contrary to other studies [15,25] this may be due to low numbers of cases and control with these maternal disease. Although the number of cases with maternal anemia was higher than that in controls, there was not significant association found with preterm birth of neonates in contrary to other studies (1, 19). Maternal Urinary tract infection was found to be a significant risk factor for preterm birth of neonates in this study, and this findings were similar with findings of other studies [4, 15, 24].

Ascending genital tract infections have previously been implicated as cause of preterm birth of neonates [24, 4, 8], our study observed corresponding finding.

Premature rupture of membrane (PROM) was also studied in this study and it was found that no significant association with preterm birth of

neonates, and this consistent with some studies [24] while other study observed significant association [31].

The presence of multiple pregnancy (twins) was observed in our study as highly significant factor associated with preterm birth of neonates and this was corresponding to other studies [4, 24, 25]. Antenatal hemorrhage as (placenta Previa or abruption placenta) was found to have significant association with preterm birth of neonates and this consistent with many studies [15, 19, 18, 20].

Uterine anomalies and cervical incompetence were considered as factors associated with of preterm birth of neonate in different studies [31, 8, 12, 22], but our study not showed significant association, this may be due to low

numbers of mothers in the study complaining of them.

The present study revealed that there was no significant association between polyhydramnios and preterm birth of neonate which was corresponding to some other studies [7, 9, 14], as well as no significant association between oligohydramnios and preterm birth of neonate which was contrary to other studies [19, 16, 28].

Conclusion and recommendations: in conclusion the most common risk factors of preterm birth of neonate were BOH, Maternal diseases in current pregnancy as well as twins and antipartum hemorrhage. It is necessary to improve prenatal care for pregnant women which may decrease the potential for preterm birth of neonate.

References:

- 1- Ananth CV, Vintzileos AM. (2008) Medically indicated preterm birth: recognizing the importance of the problem. *Clin Perinatol.* ;**35**:53–67. doi: 10.1016/j.clp.2007.11.001.
- 2- Behrman RE, Butler AS, eds. (2007). Preterm birth: causes, consequences, and prevention. Washington, D.C.:The National Academies Press. 5 p.
- 3- Blencowe H, Cousens S, Oestergaard M, et al. (2012) National, regional, and worldwide estimates of preterm birth rates in the year 2010 with time trends since 1990 for selected countries: a systematic analysis and implications. *Lancet*;379(9832):2162–2172.
- 4- Chaitanya Tellapragada, Vandana Kalwaje Eshwara, Parvati Bhat, Shashidhar Acharya, Asha Kamath, Shashikala Bhat, Chythra Rao, Sathisha Nayak, Chiranjay Mukhopadhyay.(2016).Risk Factors for Preterm Birth and Low Birth Weight Among Pregnant Indian Women: A Hospital-based Prospective Study. *J Prev Med Public Health* ;49:165-175.
- 5- Chan RL. (2014).Biochemical markers of spontaneous preterm birth in asymptomatic women. *Biomed Res Int*;2014:164081.
- 6- Choi YY, Song ES, Kim YH, Song TB.(2011). Analysis of high-risk infant births and their mortality: ten years' data from Chonnam National University Hospital Medical School, Gwangju, Korea. *Chonnam Med J* ;47(1):31–38.
- 7- Covarrubias LO, Aguirre GE, Chapuz JR, May AI, Velázquez JD, Eguiluz ME.(2008) Maternal factors associated to prematurity. *Ginecol Obstet Mex.* ;76(9):526–536.
- 8- Giraldo PC, Araújo ED, Junior JE, do Amaral RL, Passos MR, Gonçalves AK.(2012)The prevalence of urogenital infections in pregnant women experiencing preterm and full-term labor. *Infect Dis Obstet Gynecol*:878241.
- 9- Goldenberg RL, Culhane JF, Dams J, Romero R. (2008) Epidemiology and causes of preterm birth. *Lancet* 371: 75–84.
- 10- Goldenberg RL, Goepfert AR, Ramsey PS.(2005).(Biochemical markers for the prediction of preterm birth. *Am J Obstet Gynecol*;192(5 Suppl):S36-S46.
- 11- Gupta N.(2011). Perinatal mortality: an analysis of causes and strategies. *Indian Med Assoc.* ;109(4):245–246. 255–257.
- 12- Krymko H, Bashiri A, Smolin A, Sheiner E, Bar-David J, Shoham-Vardi I, Vardi H, Mazor M.(2004): Risk factors for recurrent preterm delivery. *Eur J Obstet Gynecol Reprod Biol*, 113(2):160-163.
- 13- Lazdam M, De la Horra A, Pitcher A, et al.(2010).Elevated blood pressure in offspring born premature to hypertensive pregnancy: is endothelial dysfunction the underlying vascular mechanism? *Hypertension.* 56(1):159–165.
- 14- Lo CC, Hsu JJ, Hsieh CC, Hsieh TT, Hung TH.(2007). Risk factors for spontaneous preterm delivery before 34 weeks of gestation among Taiwanese women. *Taiwan J Obstet Gynecol.* ;46:389–394.
- 15- Maria do Carmo Leal, Ana Paula Esteves-Pereira, Marcos Nakamura-Pereira, Jacqueline Alves Torres, Mariza Theme-Filha, Rosa Maria Soares Madeira Domingues, Marcos Augusto Bastos Dias, Maria Elizabeth Moreira and Silvana Granado Gama (2016). Prevalence and risk factors related to preterm birth in Brazil . *Reproductive Health*, 13(Suppl 3):127.
- 16- McElrath TF, Hecht JL, Markenson G, Harper M, Delpapa E, Allred EN, , Leviton A.(2008). Pregnancy disorders that lead to delivery before the 28th week of gestation: an epidemiologic approach to classification. *Am J Epidemiol.* ;168:980–989.
- 17- Moster D, Lie RT, Markestad T. (2008). Long-term medical and social consequences of preterm birth. *N Engl J Med* 359: 262–73 .
- 18- Moutquin JM.(2003). Classification and heterogeneity of preterm birth. *BJOG.* ;110(Suppl 20):30–33.
- 19- Nathalie Auger, Thi Uyen Nhi Le Alison L Park, and Zhong-Cheng .(2011).Association between maternal comorbidity and preterm birth by severity and clinical subtype: retrospective cohort study *BMC Pregnancy Childbirth*; 11: 67.
- 20- Ofori BD, Le Tiec M, Berard A.(2008). Risk factors associated with preterm birth according to gestational age at birth. *Pharmacoepidemiol Drug Saf*;17: pp556–564. .
- 21- O'Shea TM, Allred EN, Dammann O, et al. (2009).The ELGAN study of the brain and related disorders in extremely low gestational age newborns. *Early Hum Dev.* ;85(11):719–725. .
- 22- Peng XS (2002) A case-control study on risk factors of premature labor. *Chin J Dis Control Prev* 6: 348 (in Chinese).
- 23- Rafael TJ, Hoffman MK, Leiby BE, Berghella V.(2012) Gestational age of previous twin preterm birth as a predictor for subsequent singleton preterm birth. *Am J Obstet Gynecol.* ;206(2):156.e1–156.e6.
- 24- Samim A Al-Dabbagh I and Wafa Y Al-Tae. (2006).Risk factors for pre-term birth in Iraq: a case-control study *BMC Pregnancy and Childbirth*, 6:13
- 25- Seyed Hesamedin Nabavizadeh, Mohammad Malekzadeh, Ali Mousavizadeh Hamid Reza Ghaffarian Shirazi, Parvin Ghaffari, Nooshin Karshenas, Tahmin Malekzadeh, and Mohammad Zoladl. Retr.(2012). Retrospective study of factors related to preterm labor in Yasuj, Iran *Int J Gen Med.*; 5: 1013–101.
- 26- Shingairai AF, Siobán DH, Godfrey BW.(2004). Risk factors for prematurity at Harare Maternity Hospital, Zimbabwe. *Int J Epidemiol.* 33(6):1194–1201.
- 27- Shrestha S, Dangol SS, Shrestha M, Shrestha

- RP.(2010). Outcome of preterm babies and associated risk factors in a hospital. J Nepal Med Assoc. 50(180):286–290.
- 28- Silva AM, de Almeida MF, Matsuo T, Soares DA.(2009)..Risk factors for pre-term birth in Londrina, Paraná State, Brazil. Cad Saude Publica.25(10):2125-38.
- 29- Subapriya K, Rao AA .(2009). Association of risk factors with spontaneous preterm birth after 28 weeks and neonatal outcome--a case-control study in a tertiary care referral centreJ Indian Med Assoc. Mar;107(3):186.
- 30- Yao RY, Zhuang Y, Li HY, Yuan CJ, Hu CL, et al. (2007) Intelligence development of preterm infants in adolescence. Chin J Sch Health 28: 440–1 (in Chinese).
- 31- Zhang XR, Zeng CM, Liu J. Zhongguo Dang Dai Er Ke Za Zhi.(2011). Risk factors for preterm birth and complications in 287 late preterm infants]. Mar;13(3):177-80.
- 32- Zhang YP, Liu XH, Gao SH, Wang JM, Gu YS, Zhang JY, et al.(2012). Risk factors for preterm birth in five Maternal and Child Health hospitals in Beijing. PLoS One;7(12):e52780.

عوامل الخطر للولادة المبكرة لحديثي الولادة (الخدج) في مستشفى المكلا للأمومة والطفولة- اليمن

حنان سعيد بن بريك
علي سالم باحارثه
صالح عوض بحول
مازن أحمد جواس
فوزية فرج بامطرف

الملخص

ما زالت الولادة المبكرة هي السبب الرئيسي للأمراض والوفيات خلال الفترة الأولى للولادة. ومن المعروف أن حديثي الولادة المولودين قبل الأوان (الخدج) لديهم خطر إضافي معين من الموت، والمرض، والإعاقة. والهدف من هذه الدراسة هو التعرف على عوامل الخطر المرتبطة بالولادة المبكرة لحديثي الولادة (الخدج) في مستشفى المكلا للأمومة والطفولة -اليمن. أجريت دراسة استرجاعية لحالات مع ضوابط بنسبة 1:1 في وحدة حديثي الولادة لجناح الأطفال من أكتوبر 2012 إلى أكتوبر 2013، وجمعت بيانات الحالات والضوابط من السجلات الطبية. وشملت الدراسة 104 حالات و 104 ضوابط. أظهرت النتائج العديد من عوامل الخطورة ذات دلالة إحصائية ارتبطت بالولادة المبكرة لحديثي الولادة (الخدج) وهي: تاريخ الولادة السيئ ($p \text{ value} = 0014$)، أمراض الأم الحالية بما في ذلك ارتفاع ضغط الدم، تسمم الحمل، عدوى المسالك البولية والعدوى التناسلية (0.002 , 0.045 , 0.002 and $p \text{ value} = 0.003$ على التوالي) وكذلك التوائم والنزيف في أثناء الحمل ($p < 0.000$ and $p < 0.028$ على التوالي). خلصت الدراسة إلى أن عوامل الخطورة الأكثر شيوعاً للولادة قبل الأوان للخدج هي التاريخ المرضي السيء في أثناء الحمل السابق، أمراض الأم في الحمل الحالي وكذلك التوائم والنزيف في أثناء الحمل. من الضروري تحسين الرعاية الصحية للنساء الحوامل مما قد يقلل من احتمال الولادات المبكرة.

الكلمات المفتاحية: الخدج حديثي الولادة ، عوامل الخطورة ، المكلا، اليمن.