

Computing The Moments Of Order Statistics From Nonidentically Distributed Marshall-Olkin Extended Burr XII Random Variables

M. Gharib^{1,*} and B. I. Mohammed²

¹ Department of Mathematics, Faculty of Science, Ain Shams University, Cairo, Egypt

² Department of Mathematics, Faculty of Science, Al-Azhar University, Nasr City 11884, Cairo, Egypt

Received: 28 Oct. 2016, Revised: 2 Feb. 2017, Accepted: 12 Feb. 2017

Published online: 1 Jul. 2017

Abstract: Order statistics (os) for independent non-identically distributed (inid) random variables (rvs) is widely discussed in the literature, see, for example, Balakrishnan [3], Balakrishnan and Subramanian [4], Barakat and Abdelkader [8] and Jamjoom and Al-Saiary [14]. In this paper a recurrence relation is established for computing all single moments of all os arising from inid Marshall-Olkin extended Burr XII (MOEB XII) rvs. Another proof for the independent identical distributed (iid) rvs case is also presented and numerical examples are given.

Keywords: Order statistics; Moments; Marshall-Olkin extended Burr XII distribution.

1 Introduction

Order statistics is an important branch of statistics which deals with theory and applications of ordered rvs and functions involving them. The subject of os from inid rvs is discussed widely in the literature see for example David [10], Bapat and Beg [5] and David and Nagaraja [11]. Barakat [6] found the limit behavior of bivariate os from inid rvs. Gungor et al. [17] expressed the multivariate os by marginal ordering of inid rvs under discontinuous distribution functions. The moments of order statistics of inid rvs have been treated using three different approaches. The first approach is used when there exists a basic relation between the probability density function (pdf) and the cumulative distribution function (cdf) see Balakrishnan [3]. Applications of this approach are found in the literature for several continuous distributions see Jamjoom and Al-Saiary [15] and the references therein. In particular, Balakrishnan [3] applied this approach to derive recurrence relations for single and product moments of os from inid rvs for the exponential and right truncated exponential distributions. Childs and Balakrishnan [9] found the moments of os from inid rvs for the logistic distribution.

The second approach was introduced by Barakat and Abdelkader [8]. Although this approach is an easier manner to evaluate the moments of os of inid rvs but its application is restricted to distributions having cdfs $F(x)$ that can be written as $F(x) = 1 - \lambda(x)$. Of course this approach can also be applied if the survival function of the considered distribution has an explicit form. The first application of this second approach was by Barakat and Abdelkader [7] to Weibull distribution and then a generalized version of the approach was given by Barakat and Abdelkader [8] where they applied it to Erlang, positive exponential, pareto and laplace distributions. Later this approach is applied by Abdelkader [1,2] to compute the moments of os using the survival function of inid rvs having, respectively, Gamma and Beta distributions. Further, Jamjoom [12], Jamjoom and Al-Saiary [13] have applied this technique to compute the moments of os of inid Burr(XII) distribution as well as Beta three-parameter type I distribution.

The third approach, which referred to as the moment generating function technique, is established by Jamjoom and Al-Saiary [14] and depends mainly on the second approach. The moments of inid os for Burr type II, exponential and Erlang truncated exponential distributions, are computed using this third approach by Jamjoom and Al-Saiary [14].

* Corresponding author e-mail: dr_bahady@yahoo.com

A rv X is said to be has a MOEB(XII) distribution if its cdf is given by

$$F(x) = 1 - \frac{\alpha}{(1+x^c)^{m-\bar{\alpha}}}, \quad x > 0, \alpha, c, m > 0, \bar{\alpha} = 1 - \alpha, \tag{1}$$

In fact the MOEB(XII) distribution is an extended class that includes some distributions as special cases Burr(XII) ($\alpha = 1$), Lomax ($\alpha = 1, c = 1$) and log-logistic or weibull exponential distribution ($\alpha = 1, m = 1$). For the details of the mathematical statistical properties and application fields of the MOEB(XII) distribution see Gharib et al. [16].

In the present paper the problem of computing the moments of os from inid rvs having MOEB(XII) distribution is discussed using the second approach.

Let X_1, X_2, \dots, X_n be independent rvs and let $X_{1:n} \leq X_{2:n} \leq \dots \leq X_{n:n}$ denote the corresponding os. Bapat and Beg [5] have shown that the cdf of the r^{th} os $X_{r:n}$ ($1 \leq r \leq n$) can be expressed in terms of permanents, that is

$$F_{r:n}(x) = \sum_{i=r}^n \frac{1}{i! (n-i)!} per \begin{bmatrix} F(x) & \bar{F}(x) \\ \vdots & \vdots \\ \vdots & \vdots \end{bmatrix}, \quad -\infty < x < \infty, \tag{2}$$

where $F(x)$ and $\bar{F}(x) = 1 - F(x)$ denote the column vectors $(F_1(x), F_2(x), \dots, F_n(x))$ and $(\bar{F}_1(x), \bar{F}_2(x), \dots, \bar{F}_n(x))$ respectively. Moreover if a_1, a_2, \dots are column vectors then

$$\begin{bmatrix} a_1, a_2, \dots \\ i_1 \ i_2 \ \dots \end{bmatrix},$$

will denote the matrix obtained by taking i_1 copies of a_1 , i_2 copies of a_2 and so on. Also, in (2) $per(A)$ denotes the permanent of a square matrix A which is defined similar to the determinants except that all terms in the expansion have a positive sign, see Mine [18].

Assume that the rvs $X_i, 1, 2, \dots, n$ are inid having MOEB(XII) distribution with cdf (1).

In the next section, we derive the k^{th} moments $\mu_{n:n}^{(k)}$ and $\mu_{1:n}^{(k)}$ of the maximum and minimum of a random sample of size n from MOEB(XII) distribution.

2 Main result

Relation (3). For $n = 1, 2, \dots$ and $k = 1, 2, \dots$,

$$\mu_{n:n}^{(k)} = \frac{k}{c} \sum_{j=1}^n (-1)^{j+1} I_j, \tag{3}$$

where

$$I_j = \alpha^j \sum_{1 < i_1 < i_2 < \dots < i_n \leq n} \left(\sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} \dots \sum_{u_n=0}^{\infty} \bar{\alpha}^{\sum_{j=1}^n u_j} \right) B \left(\sum_{j=1}^n m_{i_j} (1 + u_j) - \frac{k}{c}, \frac{k}{c} \right), \tag{4}$$

and

$$\mu_{1:n}^{(k)} = \frac{k}{c} I_n, \tag{5}$$

where I_n is defined in (4) when $j = n$.

Proof. By definition

$$\mu_{n:n}^{(k)} = k \int_0^{\infty} x^{k-1} (1 - F_{n:n}(x)) dx,$$

where $F_{n:n}(x)$ is the cdf of the maximum os from inid rvs $X_i, i = 1, 2, \dots, n$ defined by

$$F_{n:n}(x) = \prod_{i=1}^n F_i(x),$$

and for MOEB(XII) distribution we have

$$F_{n:n}(x) = \prod_{i=1}^n \left(1 - \frac{\alpha}{(1+x^c)^{m_i - \bar{\alpha}}} \right),$$

then,

$$\begin{aligned} \mu_{n:n}^{(k)} &= k \int_0^\infty x^{k-1} \left\{ 1 - \prod_{i=1}^n \left(1 - \frac{\alpha}{(1+x^c)^{m_i - \bar{\alpha}}} \right) \right\} dx, \\ &= k \int_0^\infty x^{k-1} \left\{ \sum_{i=1}^n \frac{\alpha}{((1+x^c)^{m_i} - \bar{\alpha})} - \sum_{1 \leq i_1 < i_2 \leq n} \left[\frac{\alpha^2}{((1+x^c)^{m_{i_1}} - \bar{\alpha})((1+x^c)^{m_{i_2}} - \bar{\alpha})} \right] \right. \\ &\quad + \sum_{1 \leq i_1 < i_2 < i_3 \leq n} \left[\frac{\alpha^3}{((1+x^c)^{m_{i_1}} - \bar{\alpha})((1+x^c)^{m_{i_2}} - \bar{\alpha})((1+x^c)^{m_{i_3}} - \bar{\alpha})} \right] + \dots \\ &\quad \left. + (-1)^{n+1} \left[\frac{\alpha^n}{((1+x^c)^{m_{i_1}} - \bar{\alpha})((1+x^c)^{m_{i_2}} - \bar{\alpha}) \dots ((1+x^c)^{m_{i_n}} - \bar{\alpha})} \right] \right\} dx, \end{aligned}$$

putting $(1+x^c) = y^{-1}$, we get,

$$\begin{aligned} \mu_{n:n}^{(k)} &= \frac{k}{c} \left\{ \alpha \sum_{i=1}^n \left(\sum_{u=0}^\infty (\bar{\alpha})^u \right) B \left(m_i (1+u) - \frac{k}{c}, \frac{k}{c} \right) - \right. \\ &\quad - \alpha^2 \sum_{1 \leq i_1 < i_2 \leq n} \left(\sum_{u_1=0}^\infty \sum_{u_2=0}^\infty (\bar{\alpha})^{u_1+u_2} \right) B \left(m_{i_1}(1+u_1) + m_{i_2}(1+u_2) - \frac{k}{c}, \frac{k}{c} \right) \\ &\quad + \dots + (-1)^{n+1} \alpha^n \sum_{1 \leq i_1 < i_2 < \dots < i_n \leq n} \left(\sum_{u_1=0}^\infty \sum_{u_2=0}^\infty \dots \sum_{u_n=0}^\infty (\bar{\alpha})^{\sum_{i=1}^n u_i} \right) \\ &\quad \left. B \left(\sum_{j=1}^n m_{i_j}(1+u_j) - \frac{k}{c}, \frac{k}{c} \right) \right\}. \end{aligned}$$

This can be written as

$$\mu_{n:n}^{(k)} = \frac{k}{c} \sum_{j=1}^n (-1)^{j+1} I_j,$$

where

$$I_j = \alpha^j \sum_{1 < i_1 < i_2 < \dots < i_n < n} \left(\sum_{u_1=0}^\infty \sum_{u_2=0}^\infty \dots \sum_{u_n=0}^\infty (\bar{\alpha})^{\sum_{i=1}^n u_i} \right) B \left(\sum_{j=1}^n m_{i_j}(1+u_j) - \frac{k}{c}, \frac{k}{c} \right)$$

The proof of (5) follows by using the relation

$$\mu_{1:n}^{(k)} = k \int_0^\infty x^{k-1} (1 - F_{1:n}(x)) dx,$$

where

$$F_{1:n}(x) = 1 - \prod_{i=1}^n (1 - F_i(x)),$$

is the cdf of the smallest os from inid rvs.

Thus for MOEB(XII) distribution we have

$$\begin{aligned} \mu_{1:n}^{(k)} &= k \int_0^\infty x^{k-1} \prod_{i=1}^n \left(\frac{\alpha}{(1+x^c)^{m_i - \bar{\alpha}}} \right) dx \\ &= k \int_0^\infty x^{k-1} \left(\frac{\alpha}{((1+x^c)^{m_1} - \bar{\alpha})} \frac{\alpha}{((1+x^c)^{m_2} - \bar{\alpha})} \dots \frac{\alpha}{((1+x^c)^{m_n} - \bar{\alpha})} \right) dx \end{aligned}$$

putting $(1 + x^c) = y^{-1}$, then,

$$\begin{aligned} \mu_{1:n}^{(k)} &= \frac{k}{c} \alpha^n \int_0^1 y^{\sum_{i=1}^n m_i - \frac{k}{c} - 1} (1 - y)^{\frac{k}{c} - 1} \\ &\quad \left(\sum_{u_1=0}^{\infty} (\bar{\alpha} y^{m_1})^{u_1} \sum_{u_2=0}^{\infty} (\bar{\alpha} y^{m_2})^{u_2} \dots \sum_{u_n=0}^{\infty} (\bar{\alpha} y^{m_n})^{u_n} \right) dy \\ &= \frac{k}{c} \alpha^n \left(\sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} \dots \sum_{u_n=0}^{\infty} (\bar{\alpha})^{\sum_{i=1}^n u_i} \right) B \left(\sum_{i=1}^n m_i (1 + u_i) - \frac{k}{c}, \frac{k}{c} \right) = \frac{k}{c} I_n, \end{aligned}$$

where

$$I_n = \alpha^n \left(\sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} \dots \sum_{u_n=0}^{\infty} (\bar{\alpha})^{\sum_{i=1}^n u_i} \right) B \left(\sum_{i=1}^n m_i (1 + u_i) - \frac{k}{c}, \frac{k}{c} \right)$$

which can also be written as

$$I_n = \sum_{1 \leq i_1 < i_2 < \dots < i_n \leq n} \sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} \dots \sum_{u_n=0}^{\infty} \alpha^n \left(\bar{\alpha} \right)^{\sum_{i=1}^n u_i} B \left(\sum_{i=1}^n m_i (1 + u_i) - \frac{k}{c}, \frac{k}{c} \right)$$

which completes the proof.

Theorem 2.1. For $r = 1, 2, \dots, n$ and $k = 1, 2, \dots$

$$\mu_{r:n}^{(k)} = \mu_{r-1:n}^{(k)} + \sum_{j=1}^r (-1)^{j-1} \binom{n-r+j}{j-1} I_{n-r+j},$$

where $I_j, j = 1, 2, \dots, r$ is given by (4) and with the convention that $\mu_{0:n}^{(k)} = 0$.

Proof. Equation (2) can be rewritten as

$$F_{r-1:n}(x) = F_{r:n}(x) + \frac{1}{(r-1)! (n-r+1)!} \text{per} \begin{bmatrix} F(x) & \bar{F}(x) \\ r-1 & n-r+1 \end{bmatrix},$$

which is equivalent to

$$F_{r-1:n}(x) = F_{r:n}(x) + \sum_p \prod_{j=1}^n F_{i_j}(x) \prod_{j=r}^n F_{i_{n-j+1}}(x),$$

where the summation p extends over all permutations (i_1, i_2, \dots, i_n) of $(1, 2, \dots, n)$ for which

$1 \leq i_1 < i_2 < \dots < i_{r-1} \leq n$ and $1 \leq i_r < i_{r+1} < \dots < i_{n-1} \leq n$. Now let

$$x_{i_0} = \inf\{x : F_i(x) > 0\} \geq 0, \text{ for all } i.$$

Then

$$\begin{aligned} \mu_{r:n}^{(k)} &= E(X_{r:n}^k) = k \int_0^{\infty} x^{k-1} \bar{F}_{r:n}(x) dx \\ &= \mu_{r-1:n}^{(k)} + Q_{r:n}^{(k)}, \end{aligned}$$

where

$$\begin{aligned}
 Q_{r:n}^{(k)} &= k \int_0^\infty x^{k-1} \sum_p \prod_{j=1}^{r-1} (1 - \bar{F}_{i_j}(x)) \prod_{j=r}^n \bar{F}_{i_j}(x) dx \\
 &= k \int_0^\infty x^{k-1} \sum_p \prod_{j=1}^{r-1} \left(1 - \frac{\alpha}{(1+x^c)^{m_{i_j} - \bar{\alpha}}} \right) \prod_{j=r}^n \frac{\alpha}{(1+x^c)^{m_{i_j} - \bar{\alpha}}} dx \\
 &= k \int_0^\infty x^{k-1} \sum_p \left[1 - \sum_{j_1=1}^{r-1} \frac{\alpha}{(1+x^c)^{m_{i_{j_1}} - \bar{\alpha}}} + \frac{\alpha^2}{((1+x^c)^{m_{i_{j_1}} - \bar{\alpha}})((1+x^c)^{m_{i_{j_2}} - \bar{\alpha}})} \right. \\
 &\quad \left. + \dots + (-1)^{r-1} \frac{\alpha^{r-1}}{((1+x^c)^{m_{i_{j_1}} - \bar{\alpha}})((1+x^c)^{m_{i_{j_2}} - \bar{\alpha}}) \dots ((1+x^c)^{m_{i_{j_{r-1}} - \bar{\alpha}})}} \right] \\
 &\quad \prod_{j=r}^n \frac{\alpha}{(1+x^c)^{m_{i_j} - \bar{\alpha}}} dx.
 \end{aligned}$$

Putting $(1+x^c) = y^{-1}$, we get

$$\prod_{j=r}^n \frac{\alpha}{(1+x^c)^{m_{i_j} - \bar{\alpha}}} = \alpha^{n-(r-1)} \left(\sum_{u_r=0}^\infty \sum_{u_{r+1}=0}^\infty \dots \sum_{u_n=0}^\infty (\bar{\alpha})^{\sum_{i=r}^n u_j} \right) y^{\sum_{j=r}^n m_{i_j}(1+u_j)}.$$

Therefore,

$$\begin{aligned}
 Q_{r:n}^{(k)} &= \frac{k}{c} \sum_p \int_0^1 \alpha^{n-(r-1)} \left(\sum_{u_r=0}^\infty \sum_{u_{r+1}=0}^\infty \dots \sum_{u_n=0}^\infty (\bar{\alpha})^{\sum_{i=r}^n u_j} \right) \left[y^{\sum_{j=r}^n m_{i_j}(1+u_j) - \frac{k}{c} - 1} \right. \\
 &\quad - \alpha \sum_{j_1=1}^{r-1} \left(\sum_{u_1=0}^\infty \bar{\alpha}^{u_1} \right) y^{m_{i_{j_1}}(1+u_1) + \sum_{j=r}^n m_{i_j}(1+u_j) - \frac{k}{c} - 1} + \\
 &\quad + \alpha^2 \sum_{1 \leq i_1 < i_2 \leq r-1} \sum_{u_1=0}^\infty \sum_{u_2=0}^\infty (\bar{\alpha})^{u_1+u_2} y^{m_{i_{j_1}}(1+u_1) + m_{i_{j_2}}(1+u_2) + \sum_{j=r}^n m_{i_j}(1+u_j) - \frac{k}{c} - 1} + \\
 &\quad + \dots + (-1)^{r-1} \alpha^{r-1} \left(\sum_{u_r=0}^\infty \sum_{u_{r+1}=0}^\infty \dots \sum_{u_{r-1}=0}^\infty (\bar{\alpha})^{\sum_{i=1}^{r-1} u_j} \right) \\
 &\quad \left. y^{\sum_{j=1}^{r-1} m_{i_j}(1+u_j) + \sum_{j=r}^n m_{i_j}(1+u_j) - \frac{k}{c} - 1} (1-y)^{\frac{k}{c} - 1} \right] dy
 \end{aligned}$$

hence, applying integration we get

$$\begin{aligned}
 Q_{r:n}^{(k)} &= \frac{k}{c} \sum_p \left[\alpha^{n-(r-1)} \left(\sum_{u_r=0}^{\infty} \sum_{u_{r+1}=0}^{\infty} \dots \sum_{u_n=0}^{\infty} (\bar{\alpha})^{\sum_{i=r}^n u_j} \right) B \left(\sum_{i=r}^n m_{i_j} (1+u_j) - \frac{k}{c}, \frac{k}{c} \right) - \right. \\
 &\quad \left. - \alpha^{n-r+2} \sum_{j_1}^{r-1} \left(\sum_{u_1=0}^{\infty} \bar{\alpha}^{u_1} \right) \left(\sum_{u_r=0}^{\infty} \sum_{u_{r+1}=0}^{\infty} \dots \sum_{u_n=0}^{\infty} (\bar{\alpha})^{\sum_{i=r}^n u_j} \right) \right. \\
 &\quad \left. B \left(\sum_{i=r}^n m_{i_j} (1+u_j) + m_{i_{j_1}} (1+u_1) - \frac{k}{c}, \frac{k}{c} \right) \right. \\
 &\quad \left. + \alpha^{n-r+3} \sum_{1 \leq i_1 < i_2 \leq r-1} \sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} (\bar{\alpha})^{u_1+u_2} \right) \left(\sum_{u_r=0}^{\infty} \sum_{u_{r+1}=0}^{\infty} \dots \sum_{u_n=0}^{\infty} (\bar{\alpha})^{\sum_{i=r}^n u_j} \right) \right. \\
 &\quad \left. B \left(\sum_{i=r}^n m_{i_j} (1+u_j) + m_{i_{j_1}} (1+u_1) + m_{i_{j_2}} (1+u_2) - \frac{k}{c}, \frac{k}{c} \right) + \dots + \right. \\
 &\quad \left. + (-1)^{r-1} \alpha^n \left(\sum_{u_r=0}^{\infty} \sum_{u_{r+1}=0}^{\infty} \dots \sum_{u_n=0}^{\infty} (\bar{\alpha})^{\sum_{i=1}^n u_j} \right) B \left(\sum_{i=1}^n m_{i_j} (1+u_j) - \frac{k}{c}, \frac{k}{c} \right) \right].
 \end{aligned}$$

Now using the facts that $\sum_p (1) = \binom{n}{r-1}$ and that $\sum_{1 \leq i_1 < i_2 < \dots < i_m \leq n} (1) = \binom{n}{m}$, for all $n \geq m$, the above relation reduces to:

$$Q_{r:n}^{(k)} = \frac{k}{c} \alpha^n \sum_{j=1}^n (-1)^{j-1} a_j I_{n-r-j}$$

where $I_j, j = 1, 2, \dots, r$ is given by (4) and $a_j = \frac{(n-r+j)!}{(n-r+1)! (j-1)!}$ since,

$$\binom{n}{r-1} \binom{r-1}{j-1} = a_j \binom{n}{r-1}.$$

This completes the proof of the theorem.

To sum up the computations for obtaining the kth moments of all os, one needs to compute the sequence $\{I_j\}_{j=1}^{j=n}$ which is given by (4). Then recursively applying theorem 2.1, starting with the maximum $\mu_{n:n}^{(k)}$ in (3) one can obtain all moments of all os $\mu_{r:n}^{(k)}, r \leq n$ from MOEB(XII) distribution. For example if $n = 3$, we get

$$\mu_{3:3}^{(k)} = \frac{k}{c} (I_1 - I_2 + I_3),$$

where

$$\begin{aligned}
 I_1 &= \alpha^3 \left(\sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} \sum_{u_3=0}^{\infty} (\bar{\alpha})^{\sum_{i=1}^3 u_i} \right) \left[B \left(m_1(1+u_1) - \frac{k}{c}, \frac{k}{c} \right) + B \left(m_2(1+u_2) - \frac{k}{c}, \frac{k}{c} \right) \right. \\
 &\quad \left. + B \left(m_3(1+u_3) - \frac{k}{c}, \frac{k}{c} \right) \right] \\
 I_2 &= \alpha^2 \left(\sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} \sum_{u_3=0}^{\infty} (\bar{\alpha})^{\sum_{i=1}^3 u_i} \right) \left[B \left(m_1(1+u_1) + m_2(1+u_2) - \frac{k}{c}, \frac{k}{c} \right) \right. \\
 &\quad \left. + B \left(m_1(1+u_1) + m_3(1+u_3) - \frac{k}{c}, \frac{k}{c} \right) + B \left(m_2(1+u_2) + m_3(1+u_3) - \frac{k}{c}, \frac{k}{c} \right) \right] \\
 I_3 &= \alpha^3 \left(\sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} \sum_{u_3=0}^{\infty} (\bar{\alpha})^{\sum_{i=1}^3 u_i} \right) B \left(m_1(1+u_1) + m_2(1+u_2) + m_3(1+u_3) - \frac{k}{c}, \frac{k}{c} \right) \tag{6}
 \end{aligned}$$

$$\mu_{1:3}^{(k)} = \frac{k}{c} I_3$$

$$\mu_{2:3}^{(k)} = \frac{k}{c} (I_2 - 2I_3).$$

These results can be put in the following table.

$\mu_{3:3}^{*(k)}$	I_1	$-I_2$	$+I_3$
$\mu_{2:3}^{*(k)}$	I_2	$-2I_3$	
$\mu_{1:3}^{*(k)}$			I_3

The moments $\mu_{r:n}^{(k)}$, $r \leq n$ of order statistics arising from non-identically MOEB(XII) random variables with $n = 3$.

Where $\mu_{r:n}^{*(k)} = \frac{c}{k} \mu_{r:n}^{(k)}$.

For a general form of this table see Barakat and Abdelkader (2000).

3 Independent identically distributed case

In this section, the moments of the os arising from iid MOEB(XII) rvs are derived in the following theorem.

Theorem 3.1. For the case of a sample of n iid arising from MOEB(XII) distributin the kth moment ($k = 1, 2, \dots$) of the rth ($1 \leq r \leq n$) os is given by

$$\mu_{r:n}^{(k)} = k \sum_{j=1}^r (-1)^{j-(n-r+1)} \binom{j-1}{n-r} I_j,$$

where

$$I_j = \alpha^j \binom{n}{j} \left(\sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} \dots \sum_{u_n=0}^{\infty} (\bar{\alpha})^{\sum_{i=1}^j u_i} \right) B \left(m \left(\sum_{i=1}^j u_i + 1 \right) - \frac{k}{c}, \frac{k}{c} \right).$$

Proof.

$$\begin{aligned} I_j &= \binom{n}{j} \int_0^{\infty} x^{k-1} [\bar{F}(x)]^j dx \\ &= \binom{n}{j} \int_0^{\infty} x^{k-1} \left[\frac{\alpha}{(1+x^c)^m - \bar{\alpha}} \right]^j dx \\ &= \alpha^j \binom{n}{j} \left(\sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} \dots \sum_{u_n=0}^{\infty} (\bar{\alpha})^{\sum_{i=1}^j u_i} \right) B \left(m \left(\sum_{i=1}^j u_i + 1 \right) - \frac{k}{c}, \frac{k}{c} \right). \end{aligned}$$

Corollary 3.1. For iid MOEB(XII) rvs, $\mu_{1:n}^{(k)}$ becomes

$$\begin{aligned} \mu_{1:n}^{(k)} &= k \int_0^{\infty} x^{k-1} \left(\frac{\alpha}{(1+x^c)^m - \bar{\alpha}} \right)^n dx \\ &= k \alpha^n \left(\sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} \dots \sum_{u_n=0}^{\infty} (\bar{\alpha})^{\sum_{i=1}^n u_i} \right) B \left(m \left(\sum_{i=1}^n u_i + 1 \right) - \frac{k}{c}, \frac{k}{c} \right). \end{aligned}$$

4 Numerical application

The following examples are computed when $k=1$.

Case 1: independent identically distributed

Example 4.1. Let $n=3$ and $m=2, 3, 4,$ and 5 table 2 shows the results:

m	2	3	4	5
$\mu_{3:3}$	4.88419	1.65829	0.926742	0.62209

For example when $m=3,$

$$\mu_{3:3} = \frac{1}{c} (I_1 - I_2 + I_3),$$

$$I_1 = \alpha^n \left(\sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} \sum_{u_3=0}^{\infty} (\bar{\alpha})^{\sum_{i=1}^3 u_i} \right) B \left(m \left(\sum_{i=1}^3 u_i + 1 \right) - \frac{1}{c}, \frac{1}{c} \right) = 0.994973$$

$$I_2 = \alpha^2 \left(\sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} \sum_{u_3=0}^{\infty} (\bar{\alpha})^{\sum_{i=1}^3 u_i} \right) B \left(m \left(\sum_{i=1}^3 u_i + 1 \right) - \frac{1}{c}, \frac{1}{c} \right) = 0.663315$$

$$I_3 = \alpha^3 \left(\sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} \sum_{u_3=0}^{\infty} (\bar{\alpha})^{\sum_{i=1}^3 u_i} \right) B \left(m \left(\sum_{i=1}^3 u_i + 1 \right) - \frac{1}{c}, \frac{1}{c} \right) = 0.994973$$

Therefore, $\mu_{3:3} = 1.65829.$

Case 2: independent nonidentically distributed

Example 4.2.

(a) Setting $n=2, \alpha=1.5, c=0.8$ and $m_1=2, m_2=3,$ in (3), (4) we get

$$\mu_{2:2} = \frac{1}{c} (I_1 - I_2),$$

where

$$I_1 = \alpha^2 \left(\sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} (\bar{\alpha})^{\sum_{i=1}^2 u_i} \right) \left[B \left(m_1(1 + u_1) - \frac{1}{c}, \frac{1}{c} \right) + B \left(m_2(1 + u_2) - \frac{1}{c}, \frac{1}{c} \right) \right] = 2.07087,$$

$$I_2 = \alpha^2 \left(\sum_{u_1=0}^{\infty} \sum_{u_2=0}^{\infty} (\bar{\alpha})^{\sum_{i=1}^2 u_i} \right) \left[B \left(m_1(1 + u_1) + m_2(1 + u_2) - \frac{1}{c}, \frac{1}{c} \right) \right] = 0.240458.$$

Therefore $\mu_{2:2} = 2.28802.$

(b) Let $n=3, \alpha=1.5, c=0.8$ and $m_1=2, m_2=3, m_3=4$ in (3), (4) we get

$$\mu_{3:3} = \frac{1}{c} (I_1 - I_2 + I_3) = 0.62209.$$

Where I_1, I_2 and I_3 are given by (6).

Acknowledgement

The authors would like to thank the Editor-in-Chief, the Associate Editor, and the referee for their careful reading and comments which greatly improved the paper.

References

[1] Y. Abdelkader, Computing the moments of order statistics from nonidentically distributed Gamma variables with applications. Int. J. Math. Game Theo. Algebra., vol. 14, no. 1, pp. 1-8. (2004).

[2] Y. Abdelkader, Computing the moments of order statistics from independent nonidentically distributed Beta random variables. Stat. Pap., vol. 49, pp. 136-149.(2008).

[3] N. Balakrishnan, Order statistics from non-identical exponential random variables and some applications. Comput. Stat. & Data Anal., vol. 18, no. 2, pp. 203-253.(1994).

- [4] N. Balakrishnan and K. Balasubramanian, Order statistics from non-identically power function random variables. *Commun. Statistic. Theory Meth.* vol. 24, no. 6, pp. 1443-1454. (1995).
- [5] R. B. Bapat and M. I. Beg, Order statistics for nonidentically distributed variables and permanents. *Sankhyā Ser. A*, vol 51, no. 1, pp. 79-93.(1989).
- [6] H. M. Barakat, The limit behavior of bivariate order statistics from Non-identical distributed random variables. *J. Appl. Math. Polish Academy Sci.*, vol. 29, pp. 371-386 (2002).
- [7] H. M. Barakat and Y. H. Abdelkader, Computing the moments of order statistics from nonidentically distributed Weibull variables. *J. Comp. Applied Math.*,vol. 117, no. 1, pp. 85-90 (2000).
- [8] H. M. Barakat and Y. H. Abdelkader, Computing the moments of order statistics from nonidentical random variables. *Stat. Meth. Appl.*,vol. 13, no. 1, pp. 15-26 (2004).
- [9] A. Childs and N. Balakrishnan, Relations for order statistics from non-identical logistic random variables and assessment of the effect of multiple outliers on the bias of linear estimators. *J. of Stat. Plan. and Infer.*, vol. 136, no.7, pp. 2227-2253 (2006).
- [10] H. A. David, Order statistics. 2nd ed., John Wiley and Sons Inc., New York (1981).
- [11] H. A. David and H. N. Nagaraja, Order statistics. 3rd ed., Wiley Interscience. Hoboken, NJ (2003).
- [12] A. A. Jamjoom, Computing the moments of order statistics from independent non-identically distributed Burr type XII random variables. *J. of Math. and Stat.*, vol. 2, no. 3, pp. 432-438. (2006).
- [13] A. A. Jamjoom and Z. A. Al-Saiary, Computing the moments of order statistics from independent nonidentically distributed Beta type I and Erlang truncated exponential variables. *J. of Math. and Stat.*, vol. 6, no. 4, pp. 442-448 (2010).
- [14] A. A. Jamjoom and Z. A. Al-Saiary, Moment generating function technique for moments of order statistics from nonidentically distributed random variables. *Int. J. of Stat. and Sys.*, vol. 6, no. 2, pp. 77-188 (2011).
- [15] A. A. Jamjoom and Z. A. Al-Saiary, Computing the moments of order statistics from independent nonidentically distributed exponentiated Fréchet variables. *J. of Prob. and Stat.*, Volume 2012, Article ID 248750, 14 pages (2012).
- [16] M. Gharib, M. M. Mohie El - Din and B. I. Mohammed, An extended Burr (XII) distribution and its application to censored data, *Journal of Advanced Research in Statistics and Probability*, Vol. 2, Issue 1, 37 - 50 (2010).
- [17] M. Güngör, A. Gokhan and Y. Bulut, Multivariate order statistics by marginal ordering of (INID) random vectors under discontinuous df's. *J. Math. Stat.*, vol. 1, pp. 49-50. ISSN: 1549-3644 (2005).
- [18] H. Minc, Theory of permanents. *Linear and multilinear Algebra*, vol. 12, pp. 227-263 (1987).



Mohamed Gharib is Professor of Mathematical Statistics at Ain Shams University (Egypt). He received the PhD degree in Mathematical Statistics at Tashkent University (Uzbekistan.). Head of Mathematics Department at the Faculty of Science (Ain Shams University 2008 -2011). Member (founder) of the Egyptian Mathematical Society since 1992. He is a referee of several mathematical journals. His main research interests are Limit theorems for Markov chains, Distribution Theory, Characterization Theory.



Bahady Ibrahim is Associate Professor of Mathematical Statistical at Al-azhar University. He received the PhD degree in Mathematical Statistical at Al-Azhar University (Egypt). He is referee of several international journals in the frame of Mathematical Statistical and probability theory. His main research interests are: Stochastic Processes, Theory of Distributions including (Univariate and Multivariate), Estimation Theory, Reliability Theory and Order Statistics.