

2011

Design , Synthesis and Biological Evaluation Of Sulfonic Acid Esters / Benzene sulfonamides As Potential CETP Inhibitors

Reema Abu Khalaf

Al-Zaytoonah University, ReemaAbuKhalaf@hotmail.com

Ghassan Abu Sheikha

Al-Zaytoonah University, GhassanAbuSheikha@yahoo.com

Mutasem Taha

University of Jordan, MutasemTaha@yahoo.com

Follow this and additional works at: <https://digitalcommons.aaru.edu.jo/albalqa>

Recommended Citation

Khalaf, Reema Abu; Sheikha, Ghassan Abu; and Taha, Mutasem (2011) "Design , Synthesis and Biological Evaluation Of Sulfonic Acid Esters / Benzene sulfonamides As Potential CETP Inhibitors," *Al-Balqa Journal for Research and Studies* **البلقاء للبحوث والدراسات**: Vol. 14 : Iss. 1 , Article 11.

Available at: <https://digitalcommons.aaru.edu.jo/albalqa/vol14/iss1/11>

This Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in Al-Balqa Journal for Research and Studies **البلقاء للبحوث والدراسات** by an authorized editor. The journal is hosted on [Digital Commons](#), an Elsevier platform. For more information, please contact rakan@aarj.edu.jo, marah@aarj.edu.jo, u.murad@aarj.edu.jo.

DESIGN, SYNTHESIS AND BIOLOGICAL EVALUATION OF SULFONIC ACID ESTERS/ BENZENESULFONAMIDES AS POTENTIAL CETP INHIBITORS

Reema Abu Khalaf¹, Ghassan Abu Sheikha¹ and Mutasem Taha²

¹ Department of Pharmacy, Faculty of Pharmacy, Al-Zaytoonah University of Jordan, Amman, Jordan

² Department of Pharmaceutical Sciences, Faculty of Pharmacy, University of Jordan, Amman, Jordan

Cholesteryl ester transfer protein (CETP) is a glycoprotein involved in transporting lipoprotein particles and neutral lipids between high-density lipoprotein (HDL) and low density lipoproteins (LDL) and therefore it's a proper target for treating dyslipidemia and related disorders. Guided by our previously-reported pharmacophore and QSAR models for CETP inhibition, we synthesized and bioassayed a series of toluene-4-sulfonic acid 4-benzylamino-phenyl ester (8a-8m)/ N-(4-benzyl-phenyl)-4-methyl-benzenesulfonamide (6a-6l) derivatives. The proposed structures for compounds 6a-l and 8a-m were confirmed via elemental analyses, IR spectroscopy, mass spectroscopy, ¹H- and ¹³C-NMR spectra. The most potent synthesized compound 6j illustrated anti-CETP IC₅₀ of 3.4 μM.

