

Helicobacter pylori and Intestinal Parasites Co-infection: Estimation of Risk Factors among Dyspeptic Patients in Mukalla city, Hadhramout, Yemen

Eidha Ali Bin-Hameed Bin-Hameed
Hadhramout University, eidha6@gmail.com

Huda Mohammed Barajash Barajash
Hadhramout University, sun2080moon@gmail.com

Follow this and additional works at: https://digitalcommons.aaru.edu.jo/huj_nas



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Bin-Hameed, Eidha Ali Bin-Hameed and Barajash, Huda Mohammed Barajash () "Helicobacter pylori and Intestinal Parasites Co-infection: Estimation of Risk Factors among Dyspeptic Patients in Mukalla city, Hadhramout, Yemen," *Hadhramout University Journal of Natural & Applied Sciences*: Vol. 20: Iss. 1, Article 1.

Available at: https://digitalcommons.aaru.edu.jo/huj_nas/vol20/iss1/1

This Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in Hadhramout University Journal of Natural & Applied Sciences by an authorized editor. The journal is hosted on [Digital Commons](#), an Elsevier platform. For more information, please contact rakan@aar.edu.jo, marah@aar.edu.jo, u.murad@aar.edu.jo.

Helicobacter pylori and Intestinal Parasites Co-infection: Estimation of Risk Factors among Dyspeptic Patients in Mukalla city, Hadhramout, Yemen

Cover Page Footnote

Dear the Editor-in-Chief, Hadhramout University Journal of Natural & Applied Sciences Subject: SUBMISSION OF NEW MANUSCRIPT FOR PUBLICATION I am enclosing herewith a manuscript entitled “[Helicobacter pylori and Intestinal Parasites Co-infection: Estimation of Risk Factors among Dyspeptic Patients in Mukalla city, Hadhramout, Yemen]” for possible evaluation and publication. With the submission of this manuscript, I would like to undertake that the research article is an original and has not been submitted or accepted for publication elsewhere, there is no conflict of interest, and the manuscript has been read and approved by all the authors.

Article

 Digital Object Identifier:
 Received 20 October 2022,
 Accepted 12 Augustus 2023,
 Available online 13 September 2023

Helicobacter pylori and Intestinal Parasites Co-infection: Estimation of Risk Factors among Dyspeptic Patients in Mukalla city, Hadhramout, Yemen

 Eidha Ali Bin-Hameed^{1,2*} and Huda Mohammed Barajash¹
¹Biology Department, Faculty of Science, Hadhramout University, Yemen

²Health Sciences Department, Faculty of Medicine and Health Sciences, University of Science and Technology, Aden, Yemen

 *Corresponding author: eidha6@gmail.com

 This is an open-access article under production of [Hadhramout University Journal of Natural & Applied Science](#) with eISSN 2790-7201

Abstract : Patients with dyspepsia often have *Helicobacter pylori* (*H. pylori*) infections, which are more common in the countries of the third world. *H. pylori* has ability to colonizing the gastric mucosa, and the infections often lead to development of various gastrointestinal complications. In addition, *H. pylori* infection and intestinal parasite colonization in humans is a regular occurrence. The purpose of the current research was to determine the prevalence of intestinal parasites and *H. pylori* co-infection, as well as related risk factors, in dyspeptic patients who were sent to hospitals in Mukalla City, Hadhramout governorate, Yemen. This research used a cross-sectional design to collect and analyze 100 stool samples. The purpose was to detect *H. pylori* infection using a quick immunochromatographic stool-based antigen test, and also to identify any presence of parasite infection. To obtain data, we used a structured questionnaire. The data were examined using SPSS version 24, and logistic regression analysis was utilized to identify the independent predictive risk variables, and P-value considered significant at 0.05. The total prevalence of *H. pylori* infection among dyspeptic patients was 30%. Males had a higher prevalence infection of *H. pylori* (COR=0.386, 95%CI=0.160-0.934, P=0.035). Furthermore, positive *H. pylori* results were significantly linked with unfiltered drinking water (COR=3.67; 95%CI=1.436-9.363; P=0.007), regurgitation and heartburn symptoms (COR=0.865, 95%CI=0.034-0.536, P=0.004), and antibiotic use (COR=0.312, 95%CI=0.125-0.780, P=0.013). This study indicates that intestinal protozoa *Entamoeba histolytica* and *Giardia lamblia* were the commonest co-infected parasites with *H. pylori* in dyspeptic patients with no significant association. Finally, gender, unfiltered water source, regurgitation and heartburn symptoms, and antibiotics used were risk variables associated with co-infection with the intestinal parasites *E. histolytica* and *G. lamblia* and *H. pylori*.

Keywords: Co-infection, Dyspeptic patients, *Entamoeba histolytica*, *Giardia lamblia*, *Helicobacter pylori*, Hadhramout, Prevalence, Yemen

1. Introduction:

H. pylori colonization of the human gastrointestinal mucosa leads to the development of gastric cancer, chronic gastritis, peptic ulcers, gastric mucosa associated lymphoid tissue (MALT) lymphoma, and peptic ulcers [1]. Some developing countries have 90%, while developed countries have values ranging from 25 to 40% prevalence of *H. pylori* infection [2]. Dyspepsia is a prevalent health issue that is seen globally [3], and is the most common symptoms of the upper gastrointestinal [4]. The incidence rate of dyspepsia associated with *H. pylori* infection is 25% [3,5]. Uncertain factors, including viral and non-infectious agents, may contribute to dyspepsia [6]. The major causes include gastroduodenitis, peptic ulcer, malignancies, esophagitis, parasitic infection and functional dyspepsia [7].

The poly-microbial that causes the gastrointestinal disturbances have clinical significance [8]. Co-infection of *H. pylori* and intestinal parasites are commonly causing of

gastrointestinal symptoms and disturbances [9]. Intestinal parasites have a global distribution that affects millions of people worldwide [10]. Intestinal parasites and co-infection with *H. pylori* are both prevalent causes of gastrointestinal illnesses and important infectious pathogens for global public health. Intestinal parasite infection is common in Uganda [11], Pakistan [12], Ethiopia [13], Sudan [14], Nigeria [15], and Yemen [16], and some studies have indicated a high frequency of *H. pylori* infection among these individuals. In Yemen, some studies reported the prevalent infection of *H. pylori* in dyspepsia, gastritis, duodenal and gastric ulcers patients in Mukalla city, Hadhramout [17], Sana'a city [18], Taiz city [19], and Dhamar [20-21]. However, there are little data available on the prevalence of co-infection with intestinal parasites and *H. pylori*. In addition to *H. pylori* infection, dyspeptic patient's clinical outcomes must be evaluated for intestinal parasite infection. The aim of this research is to look at intestinal parasites and *H. pylori* co-

infections, along with the risk factors associated with these co-infections, in patients in Mukalla City, Hadhramout Governorate, Yemen who have dyspeptic symptoms.

2. Materials and methods:

2.1 Sample size, population, and study design:

This research was planned as a cross-sectional study and done in Mukalla city, which is located in Hadhramout Governorate, Yemen. One hundred stool samples were taken from patients who were referred to several major hospitals in Mukalla city, Hadhramout with probable dyspepsia.

2.2 Inclusion and exclusions criteria:

In this study, patients who showed symptoms of dyspepsia (heartburn and acid regurgitation) were included. Patients when they were without symptoms of dyspepsia were excluded.

2.3 Collection of data:

For the purpose of gathering information about the risk factors of the infection of H. pylori, a standard structured questionnaire was utilized. It was composed of an interview with the participant and a set of systematic questions about the study's variables.

2.4 Stool antigen H. pylori test:

The InTec Products, INC, USA offers the H. pylori one-step antigen test, a quick immunochromatographic assay for the identification of H. pylori antigen in samples of human feces. Based on the manufacturer's criteria, this test was utilized to generate qualitative and visual results with a high degree of specificity (98.6%) and sensitivity (98.4%).. Feces sample passes across the conjugate pad when it is introduced to the sample pad, mobilizing the gold anti-H. pylori conjugate that is placed on the conjugation pad. The combination undergoes a reaction with the anti-H. pylori that is present on the test section of the membrane as it moves downwards due to capillary action [17].

2.5 Detection technique for intestinal parasites:

Direct saline wet-mounting preparation was used. A little amount of stool sample was located on a glass slide microscope. Using a light microscope at Olympus, Philippines, a drop of Lugol iodine was applied, covered with

a glass coverslip, and the specimen was examined at 40× and 100× magnification to look for parasite cysts and eggs [22].

2.6 Participant consent and ethical research permission:

Faculty of Science at Hadhramout University granted this project ethical permission. Before beginning the study, written consent was obtained. The hospital's management provided letters of authorization. The participants' information was collected after they verbally consented to it in accordance with the informed consent with confidentially the findings of each participant.

2.7 Analysis of Data:

Version 24 of SPSS was utilized for the analysis of the data, a statistical software package for social sciences. The characteristics of the participants were described using descriptive statistics of frequencies and percentages. Logistic regression analysis was used so that a determination could be made about the nature of the connection between the dependent and independent variables. The 95% confidence intervals (CI) and crude odds ratio/adjusted odds ratio (OR) were used to show the relationship between the variables. At 0.05, the P-value has been described as significant.

3. Results

3.1 .Risk factors and H. pylori prevalence in dyspeptic patients.

In this examination, it was discovered that H. pylori infection had a prevalence of 30%. A higher prevalence of H. pylori infection was seen among men (63% vs. 37%; COR=0.386, 95%CI=0.160-0.934, P=0.035), although this figure shows a statistically significant difference between males and females. Age groups 19 to 32 years show the highest rate 40.0% of H. pylori infections, followed by age group 33 to 46 years that show 33.0% of H. pylori infections without significant related factors. Unfiltered water sources increased the incidence of H. pylori infection in positive patients by 3.667 times (95% CI: 1.436–9.363, P=0.007). Food habits and H. pylori infection were not substantially correlated (P > 0.05). Clinically, there is a strong correlation between heartburn and regurgitation (COR=0.865, 95%CI=0.034-0.536, P=0.004). Additionally, as indicated in Table 1, there is a significant association between using antibiotics and H. pylori infection (COR=0.312, 95%CI=0.125-0.780, P=0.013).

Table 1. H. pylori infection prevalence and risk factors in individuals with dyspepsia.

	Variable	No. of cases	Positive fecal Ag(%)	COR	CI(95%)	P-value
Gender	Male	47	19(63.0)	0.386	0.160-0.934	0.035*
	Female	53	11(37.0)	1		
Age group (years)	5 – 18	3	1(3.0)	0.067	0.072-12.105	0.958
	19 – 32	38	12(40.0)	1.011	0.327-3.124	0.985
	33 – 46	37	10(33.0)	1.260	0.397-3.995	0.695
	47 – 60	22	7(24.0)	1		
Food patterns	Fatty, citrus and spicy	93	27(90.0)	1.833	0.384-.8.746	0.447
	Nothing	7	3(10.0)	1		
Water sources	Unfiltered	48	8(27.0)	3.667	1.436-9.363	0.007*
	Filtered	52	22(73.0)	1		
Clinical symptoms	Regurgitation	4	0(0.0)	269245810.7	0.000	0.999
	Heartburn	37	6(20.0)	0.139	0.258-5.219	0.845
	Heartburn and regurgitation	38	21(70)	0.865	0.034-0.536	0.004*
	No symptoms	21	3(10.0)	1		
Antibiotics used	Used	29	14(47.0)	0.312	0.125-0.780	0.013*
	Non-used	71	16(53.0)	1		

*Statistically significant P-value <0.05, Abbreviations: CI, confidence interval, Crude Odds Ratio; COR

3.2 Intestinal protozoa co-infection with H. pylori:

The intestinal protozoa *Entamoeba histolytica* (*E. histolytica*) cyst and the trophozoite and *Giardia lamblia* (*G. lamblia*)

trophozoite were detected 6(20%) in dyspeptic patients infected with H. pylori without statistically significant association (P = 0.890) as presented in Table 2.

Table 2. Co-infection of H. pylori and intestinal protozoa among dyspeptic patients

Type of intestinal protozoa infection	No. of cases	Positive fecal Ag(%)	COR	CI(95%)	P-value
<i>E. histolytica</i> cyst	9	2(7.0)	3.500	0.284-43.161	0.328
<i>E. histolytica</i> trophozoite	5	2(7.0)	1.500	0.106-21.312	0.765
<i>G. lamblia</i> trophozoite	2	1(3.0)	1		
<i>E. histolytica</i> cyst and trophozoite	1	0(0.0)	1615474864	0.000	1.000
<i>E. histolytica</i> cyst and <i>G. lamblia</i> trophozoite	2	1(3.0)	1		
No parasitic infection	81	24(80.0)	2.375	0.316-17.853	0.401
Total	100	30(100.0)			

For those related risk factors that demonstrated significance in the estimation of the crude odds ratio, the logistic regression of multivariate analysis was carried out. Table 3 demonstrates

that there is still a significant association between the incidence of H. pylori infection, sex, unfiltered water, regurgitation and heartburn symptoms, and antibiotic usage.

Table 3. Analysis of multivariate logistic regression for significant associated risk factors with H. pylori infection among dyspeptic patients

Variable		Fecal antigen test		
		AOR	CI(95%)	P-value
Gender	Male	2.591	1.071-6.267	0.035*
	Female	1		
Water source	Non filtered	0.727	0.107-0.696	0.007*
	Filtered	1		
Clinical symptoms	Regurgitation	-	-	-
	Heartburn	0.161	0.258-5.219	0.845
	Heartburn and regurgitation	7.412	1.866-29.444	0.004*
	No symptoms	1		
Antibiotics used	Used	3.208	1.283-8.024	0.013*
	Non-used	1		

*Statistically significant P-value <0.05

Abbreviations: CI, confidence interval; AOR Adjusted odds ratio

4. Discussion:

The prevalence of H. pylori infection among patients with dyspepsia was estimated to be 30% in the present research. Other investigations exhibited similar findings regarding the frequency of H. pylori infection within varying ranges: 23.5% [23], 24.3% [24], 29.6% [25], 37.8% [26], and 80.3% [27]. The variance in the prevalence of H. pylori infections may be attributed to lifestyle factors or different levels of exposure to risk factors.

Our results revealed a high prevalent infection of H. pylori observed among males than among females with significant association (P=0.032). Similar findings revealed that infection of H. pylori was more prevalent among men [28], other more prevalent H. pylori infection showed in males 78% [29]. On the other hand, 35.7% of females found to be positive for H.

pylori infection [23]. Another study showed that in general 62% of women were infected with H. pylori [30], while a study showed 63.6% infected females with H. pylori [31]. According to a study, women are more likely than men to have H. pylori infection [25].

The present investigation revealed that the largest prevalence of H. pylori infections was seen among individuals aged 19–32 and 33–46 years. Furthermore, there was no significant association between infection frequency and age among those aged 47 to 60 years (P = 0.969). Similar research revealed that 61.6% of H. pylori-positive individuals were in the 31–60 age range [29]. Infection of H. pylori starts early in childhood and adolescence and peaks in adults between the ages of 35 and 44 [30]. A peak prevalence was seen, according to another study in the group of people aged 40 to 49 years [32].

The prevalent infection of *H. pylori* was substantially greater in patients aged 5-50 years [33]. In the age range of 46 to 55 years, the highest prevalence of *H. pylori* was noted [34]. Another study found that the age range of 18 to 30 years had a higher frequency of *H. pylori* infection [28], while a third study showed that the age range of 30-39 years had a 90.8% prevalence of *H. pylori* infection [35].

Our research revealed a lack of substantial link between the consumption of food meals and the occurrence of *H. pylori* infection ($P=0.349$). However, we did observe a statistically significant link between the source of water and the frequency of *H. pylori* infection ($P=0.012$). Some studies showed *H. pylori* infection prevalence was associated with some foods [35-36]. Another study carried out in Yemen revealed a statistically significant correlation between *H. pylori* positivity and fat rich meals and Qat chewing [18]. Other study results showed prevalence of *H. pylori* infection were high significantly among individuals with consumption of unboiling and unclean water sources [24,33,37-38].

Participants who reported regurgitation and heartburn showed substantial clinical evidence of *H. pylori* infection ($P = 0.001$). Additionally, the prevalence of *H. pylori* infection was substantially correlated with the usage of antibiotics ($P=0.011$). Similar findings demonstrated that *H. pylori* infection affected every patient with upper stomach pain and frequent burping [30]. Other investigations revealed that the most typical symptoms of *H. pylori* infection were dyspepsia, heartburn, and epigastric discomfort [18,28,37].

According to a different study, symptomatic participants frequently experienced heartburn, followed by nausea, vomiting, abdominal pain, and black, unpleasant-smelling stools [23]. According to a study, esophagitis, peptic ulcers, and gastritis are all substantially correlated with *H. pylori* infection [38]. Consistent with previous research, the antibiotics applied were shown to be a significant contributing factor to the probability of *H. pylori* infection [18,39].

The most prevalent parasites found to be co-infected with *H. pylori* in dyspeptic patients in the current investigation were intestinal protozoa, with no evidence of a significant correlation, which is consistent with the findings of prior studies [40]. Intestinal protozoa and *H. pylori* infection may be related, according to several research [6,41]. In Yemen, patients claiming to have gastric giardiasis had *H. pylori* infection rate of 75% [16].

Another study has shown a high *H. pylori* infection prevalence associated with intestinal parasitic infections; protozoal, helminthic and both [42], while other study showed the patients with *H. pylori* infection have *G. lamblia* and *E. histolytica*/dispar [43]. Another study found that *E. histolytica* was the most common parasite co-infected with *H. pylori* infection [28]. In addition to the infection of *H. pylori*, *G. lamblia* was reported among patients with dyspepsia [6,44]. There is a high association between concurrent intestinal parasites *G. lamblia* and *H. pylori* infections in rates of 38.3–70.2% [13].

Furthermore, intestinal parasites *E. histolytica*, *Entamoeba coli*, and *G. lamblia* were more common among patients infected with *H. pylori* [14]. Other studies showed involvement of *H. pylori* infection and intestinal parasites *E. histolytica*, *G. lamblia*, and other parasites with significant

differences as probable causes of duodenal, gastric, and peptic ulcer [15]. Other study concluded a high of *H. pylori* prevalence and *G. lamblia* [45]. The prevalence of *G. lamblia* was significantly high among patients infected with *H. pylori* [46]. The prevalence of *H. pylori* infection was associated risk factor for *G. intestinalis* infection [47]. A study showed 69.4% *H. pylori*, 51.4% co-infected with *G. lamblia* or *E. histolytica* [48]. *H. pylori* co-infected with *E. histolytica*/*E. dispar* [39]. Therefore, *H. pylori* infection may provide appropriate conditions for infection of giardiasis [49].

In our study, the rate of co-infections of *H. pylori* and *G. lamblia* or *E. histolytica* could be due to the role of *H. pylori*-produce urease enzyme that converts the stomach wall urea to ammonia resulting in increased in pH of the stomach [50]. It is found a significant increased urease enzyme activity among co-infected Giardiasis and *H. pylori* compared to single *G. lamblia* infected individuals [51]. Additionally, possible transmission routes similar to the fecal–oral route could explain the observed the incidence of intestinal parasites among infected participants with *H. pylori* in this study.

This research has numerous limitations: (1) the cross-sectional study design limits the ability to reveal a true association between dyspepsia and *H. pylori* infection, (2) *H. pylori* testing based on stool antigen may exaggerate the prevalence of *H. pylori* infection, (3) The approach of direct wet mount preparation may underestimate the prevalence of intestinal parasites, (4) the prevalence of co-infection *H. pylori* with intestinal parasites could not be representative the general population since only dyspeptic symptomatic individuals were included in the study.

5. Conclusion

The present study showed 30% of dyspepsia and *H. pylori* infection. Sex, unfiltered drinking water, heartburn and regurgitation symptoms and antibiotics drug used were significantly associated with intestinal parasites and *H. pylori* accompaniment co-infection. The reported findings need more research to study the observed correlation in greater depth, as well as reasons for this association. Screening for *E. histolytica* and *G. lamblia* infection in the intestine is advised for effective care of symptomatic dyspeptic patients.

Acknowledgements

We appreciate all the participants and laboratory technicians who contributed to this study. Special thanks to Hospitals administrations for its unreserved supporting during collection of data. Much gratitude is extended to the Biology Department of the Faculty of Science at Hadhramout University for their contributions to the advancement of scientific research.

References:

- [1] A. Sethi, M. Chaudhuri, L. Kelly, W. Hopman, "Prevalence of *Helicobacter pylori* in a first nations population in Northwestern Ontario," *Canadian Family Physician*, vol. 59, no. 4, p. e182–e187, 2013.
- [2] F. Francesco, A. Tortora, R.T. Di, G. D'Angelo, I. Gianluca, "Role of *Helicobacter pylori* infection on nutrition and metabolism," *World Journal Gastroenterology*, vol. 20, p. 12809-12817, 2014, DOI: 10.3748/wjg.v20.i36.12809.

- [3] F. Naz, S. Malik, S. Afzal, S.A. Anwar, "Frequency of Seropositivity of *Helicobacter pylori* in Patients Presenting with Dyspepsia," *Journal of Ayub Medical College Abbottabad*, vol. 25, no. 3-4, p. 50-54, 2013.
- [4] B. Tepeš, "Subgroups of dyspepsia, In: Duvnjak M, editor. *Dyspepsia in clinical practice*," Berlin: Springer, p. 9–18.
- [5] K. Yazdanpanah, N. Moghimi, V. Yousefinejad, E. Ghaderi, A. Azizi, S.F. Nazem, "Dyspepsia prevalence in general population aged over 20 in the west part of Iran," *Journal of the Pakistan Medical Association*, vol. 62, no. 7, p. 672–676, 2012.
- [6] S.A. Fouad, S. Esmat, M.M.A. Basyoni, F.M. Salah, M.H. Kobaisi, "Molecular identification of *Giardia intestinalis* in patients with Dyspepsia," *Digestion*, vol. 90, p. 63–71, 2014, doi: 10.1159/000362644.
- [7] S.M. Ayana, B. Swai, Maro V.P. G.S. Kibiki, "Upper gastrointestinal endoscopic findings and prevalence of *Helicobacter pylori* infection among adult patients with dyspepsia in northern Tanzania," *Tanzania Journal of Health Research*, vol. 16, no.1, p. 1-9, 2014, doi: 10.4314/thrb.v16i1.3.
- [8] B.M. Peters, M.A. Jabra-Rizk, G.A. O'May, C.J. William, M.E. Shirliff, "Polymicrobial interactions: impact on pathogenesis and human disease," *Clinical Microbiology Review*, vol. 25, no. 1, p. 193–213, 2012, doi: 10.1128/CMR.00013-11.
- [9] A.G. Gravina, R.M. Zagari, C. De Musis, L. Romano, C. Loguercio, M. Romano, "*Helicobacter pylori* and extragastric diseases," *World Journal Gastroenterology*, vol. 24, no. 29, p. 3204–3221, 2018, <https://doi.org/10.3389/fmicb.2022.972777>.
- [10] C.W. Liao, C.J. Fu, C.Y. Kao, Y.L. Lee, P.C. Chen, T.W. Chuang, T. Naito, C-M. Chou, Y-C Huang, I. Bonfim, C-K. Fan, "Prevalence of intestinal parasitic infections among school children in capital areas of the Democratic Republic of São Tomé and Príncipe, West Africa," *African Health Sciences*, vol. 16, no. 3, p. 690–697, 2016, doi: 10.4314/ahs.v16i3.8.
- [11] J. Ankarklev, E. Hestvik, M. Lebbad, J. Lindh, D.H. Kaddu-Mulindwa, J.O. Andersson, T. Tylleskär, J.K. Tumwine, S. G Svärd, "Common Coinfections of *Giardia intestinalis* and *Helicobacter pylori* in non-symptomatic Ugandan children," *PLoS Neglected Tropical Diseases*, vol. 6, 8, p. e1780, 2012, doi: 10.1371/journal.pntd.0001780.
- [12] J. Yakoob, Z. Abbas, R. Khan, K. Tariq, S. Awan, M.A. Beg, "Association of *Helicobacter pylori* and protozoal parasites in patients with chronic diarrhoea," *British Journal of Biomed Sciences*, vol. 75, n. 3, p. 105–109, 2018, <https://doi.org/10.1080/09674845.2017.1420129>.
- [13] A. Seid, Z. Tamir, B. Kasanew, M. Senbetay, "Coinfection of intestinal parasites and *Helicobacter pylori* among upper gastrointestinal symptomatic adult patients attending Mekanesalem Hospital, northeast Ethiopia," *BMC Research Notes*, vol. 11, no. 144, 2018, <https://doi.org/10.1186/s13104-018-3246-4>.
- [14] Y. Abd Elbagi, A.B. Abd Alla, M.B.E. Saad, "The relationship between *Helicobacter pylori* infection and intestinal parasites in individuals from Khartoum state, Sudan: a case-control study," *F1000Research*, vol. 8, 2019, <https://doi.org/10.12688/f1000research.21397.1>.
- [15] O.C. Ifeanyi, O. Prosper, O.U. Adogu, "Prevalence of *Helicobacter pylori* and other intestinal parasites amongst duodenal and gastric ulcer patients at Imo state University Teaching Hospital, Orlu, south eastern Nigeria," *Journal of Medical Sciences*, vol. 4, 9, p. 362-369, 2013, DOI: 10.14303/jmms.2013.120.
- [16] M.A. Bin Mohanna, L.M. Al-Zubairi, A.K. Sallam, "Prevalence of *Helicobacter pylori* and parasites in symptomatic children examined for *Helicobacter pylori* antibodies, antigens, and parasites in Yemen" *Saudi Medical Journal*, vol. 35, no. 11, p. 1408–1411, 2014.
- [17] E.A. Bin-Hameed, H.M. Barajash, "Screening for the prevalence of *Helicobacter pylori* infection among dyspeptic patients using simple fecal antigen and serum antibody diagnostic methods at Mukalla city Hospitals, Hadhramout, Yemen," *African Journal of Microbiology Research*, vol. 15, no. 6, p. 325-333, 2021, DOI: 10.5897/AJMR2020.9413.
- [18] A.S.S. Moharram, A. Alqadym, A.A. Alhetary, "Prevalence of *Helicobacter pylori* among Gastritis Patients in Sana'a, Yemen," *International Journal of Current Microbiology and Applied Sciences*, vol. 4, no. 3, p. 769-778, 2015.
- [19] G.A. A-Ameri, M.N. Alkadasi, "The prevalence of *Helicobacter pylori* and risk factors infection associated in Taiz city, Yemen," *International Journal of Current Microbiology and Applied Sciences*, vol. 2, no. 8, p. 226-23, 2013.
- [20] D.A. Almashhadany, S.M. Mayas, H.I. Mohammed, A.A. Hassan, I.U.H. Khan, "Population and Gender-Based Investigation for Prevalence of *Helicobacter pylori* in Dhamar, Yemen," *Canadian Journal of Gastroenterology and Hepatology*, 2023, 3800810. doi: 10.1155/2023/3800810.
- [21] D.A. Almashhadany, S.M. Mayass, "Prevalence of *Helicobacter pylori* in Human in Dhamar Governorate / Yemen," *Journal of Medical and Pharmaceutical Sciences*, vol. 2, no. 1, p. 1-10, 2018.
- [22] B. Chakarova, "Comparative evaluation of the diagnostic methods for detection of *Giardia intestinalis* in human fecal samples, *Trakia Journal of Sciences*, vol. 8, p. 174-179, 2010, DOI: 10.13140/RG.2.1.4433.2966.
- [23] E.S. Samson, O.J. Okeleke, A.Y. Richard, F.T. Gideon, A.L. Olutoyosi, O. Damilola, "Screening for *Helicobacter pylori* Infection among Undergraduate Students of a Tertiary Institution using serum Antibody and Stool Antigen Detection Methods," *ScienTechn Research*, vol. 3, no. 2, p. 10 pages, 2018, DOI: 10.26717/BJSTR.2018.03.000883.
- [24] P. Aitila, M. Mutyaba, S. Okeny, M.N. Kasule, R. Kasule, F. Ssedyabane, B. Okongo, "Prevalence and Risk Factors of *Helicobacter pylori* Infection among Children Aged 1 to 15 Years at Holy Innocents Children's Hospital, Mbarara, South Western Uganda," *Journal of Tropical Medicine*, ID 9303072, 2019, <https://doi.org/10.1155/2019/9303072>.
- [25] A. Alim, M. Ataş, T. Güneş, S. Özkan, N. Dündar, "Comparison of antigen and antibody detection tests used for diagnosing the *Helicobacter pylori* infection in symptomatic patients," *Basic Clinical Sciences*, vol. 4, p. 61-70, 2010.
- [26] A. Iranikhah, M-R. Ghadir, S. Sarkeshikian, H. Saneian, A. Heiari, M. Mahvari, "Stool Antigen Tests for the Detection of *Helicobacter pylori* in Children," *Iranian Journal of Pediatric*, vol. 23, no. 2, p. 138–142, 2013.

- [27] M. Calik, O. Karamese, S.A. Acar, Y. Karamese, F. Dicle, S. Albayrak, B. Can, A. Guvendi, M. Tuegut, C.H. Yazgi, "Investigation of *Helicobacter pylori* antigen in stool samples of patients with upper gastrointestinal complaints," *Brazilian Journal of Microbiology*, vol. 47, no. 1, 167-171, 2016.
- [28] M.A. Abo-Shadi, T.A. El-Shazly, M.S. Al-Johani, "Clinical, Endoscopic, Pathological and Serological Findings of *Helicobacter pylori* Infection in Saudi Patients with Upper Gastrointestinal Diseases," *British Journal of Medical Research*, vol. 3, no. 4, p. 1109-1124, 2013, DOI: 10.9734/BJMMR/2013/2650.
- [29] D. Chandrababu, H.P. Nandeesh, D. Suvarna, H.V. Aradya, T.R. Vijaykumar, I. Suresh, S.R. Kothe, "Association of *H. pylori* Infection with Gastrointestinal Disease: A cross Sectional Study from Mysuru, India," *National Journal of Medical Research*, vol. 6, no. 3, p. 292-295, 2016.
- [30] L.B.K. Mabeku, M.L.N. Nganga, H. Leundji, "Potential risk factors and prevalence of *Helicobacter pylori* infection among adult patients with dyspepsia symptoms in Cameroon," *BMC Infectious Diseases*, vol. 18, p. 278, 2018, doi: 10.1186/s12879-018-3146-1.
- [31] J.A. Aminde, G.A. Dedino, C.A. Ngwasiri, K.S. Ombaku, M.C.A. Mahop, L.N. Aminde, "Helicobacter pylori infection among patients presenting with dyspepsia at a primary care setting in Cameroon: seroprevalence, five-year trend and predictors," *BMC Infectious Diseases*, vol. 19, no. 1, p. 30, 2019, doi: 10.1186/s12879-019-3677-0.
- [32] C.G. Simón, F.C.A. Andrés, C.G. Tomás, A.C.L. Luis, I.G.G. Héctor, E.M. Santiago, "Prevalence of *Helicobacter pylori* and Histopathological Features in Gastric Biopsies from Patients with Dyspeptic Symptoms at a Referral Center in Medellín," *Revista Colombiana Gastroenterología*, vol. 31, no. 1, 9-14, 2016.
- [33] A.F. Syam, M. Miftahussurur, D. Makmun, I.A. Nusi, L.H. Zain, F. ZulkhairiAkil, W.B. Uswan, D. Simanjuntak, T. Uchida, P. Adi, A.P. Utari, Y.A.A. Rezkiha, P. Subsomwong, S.R. Nasronudin, Y. Yamaoka, "Risk Factors and Prevalence of *Helicobacter pylori* in Five Largest Islands of Indonesia: A Preliminary Study," *PLOS ONE*, vol. 23, no. 2, 14 pages, 2015, doi: 10.1371/journal.pone.0140186.
- [34] M. Javed, K. Amin, D. Muhammad, A. Husain, N. Mahmood, "Prevalence of *H. pylori*," *Professional Medical Journal*, vol. 17, no. 3, 431-439, 2010.
- [35] Y. Zhu, X. Zhou, J. Wu, J. Su, G. Zhang, "Risk Factors and Prevalence of *Helicobacter pylori* Infection in Persistent High Incidence Area of Gastric Carcinoma in Yangzhong City," *Gastroenterology Research Practice*, vol. ID 481365, 10 pages, 2014, doi: 10.1155/2014/481365.
- [36] S. Gul, L. Jawed, S. Tariq, S. Aziz, "Helicobacter pylori Association with Acid Peptic Disease: It's Incidence in Population Having Increase Junk Food Intake," *Translational Biomedicine*, vol. 7, no. 4, p. 90, 2016, doi: 10.21767/2172-0479.100090.
- [37] A.S. Yadav, U.B. Kulkarni, B.C. Kumar, U.V. Takalkar, "Prevalence of *Helicobacter pylori* infection among dyspepsia patients with mucosal lesion in tertiary care hospital," *Inter Sur Journal*, vol. 5, p. 2264-2267, 2018, DOI: <https://doi.org/10.18203/2349-2902.isj20182234>.
- [38] A.M. Al-Makdad, M.H. Al-Dholae, A.A.K. Thabet, M.A. Al-Haimi, O.S. Balfaqih, A.M. Al-Haddad, "Prevalence of *Helicobacter pylori* Infection in Yemeni Patients," *Yemeni Journal for Medical Sciences*, vol. 7, p. 33-38, 2013, DOI: 10.20428/YJMS.7.1.5.
- [39] T.H. Nguyen, T.B. Phan, V.B. Nguyen, T.V.H. Nguyen, T.T.B. Phan, V.B. Nguyen, T.T.H. Hoang, T.L.A. Le, T.T.M. Nguyen, S.N. Vu, "Prevalence and Risk Factors of *Helicobacter pylori* Infection in Muong Children in Vietnam," *Annals and Clinical Laboratory Research*, vol. 5, no. 1, 2017.
- [40] A.S. Rahman, S.A. Sarker, T. Ahmed, R. Islam, M.A. Wahed, D.A. Sack, "Relationship of Intestinal Parasites, *H. pylori* Infection with Anemia or Iron Status Among School Age Children in Rural Bangladesh," *Journal of Gastroenterology and Hepatology Research*, vol. 2, no. 9, p. 769-773, 2013, DOI: 10.21767/2386-5180.1000159.
- [41] H.A. Shavalipour, R. Mohebi, S. Ghafurian, S. Aslani, A. Maleki, J. Kardan, H. Hiedari, N. Sadeghifard, "Estimation of the parasitic infection prevalence in children with *Helicobacter pylori* infection in Ilam City (2012–2013)," *Archives of Pediatric Infectious Diseases*, 2014, DOI: <https://doi.org/10.5812/pedinfect.15294>.
- [42] F.M. Shaldoum, "Anemia Related to Infection with *Helicobacter pylori* and Intestinal Parasites in Egypt," *Middle East of Journal Applied Sciences*, vol. 05, no. 04, p. 1287-1296, 2015, DOI: 10.21608/idj.2016.146765.
- [43] H. Kazemian, H.H. Kardan, J. Yamchi, A. Shavalipour, S. Ghafourian, R. Mohebi, H.R. Hourri, N. Sadeghifard, "Relationship between *Helicobacter pylori* Infection and Parasitic Infection in Patients in Ilam," *Infection Epidemiology and Medicine*, vol. 2, no. 2, p.15-17, 2016, DOI: 10.18869/modares.iem.2.2.15.
- [44] F.I. Hassanein, A.I. Shehata, R. Abdul-Ghani, "Giardia lamblia and *Helicobacter pylori* infections among mentally challenged individuals in rehabilitation centers in Alexandria, Egypt" *Journal of Infectious Devel Country*, vol. 11, no. 7, p. 577-582, 2017, doi: 10.3855/jidc.8783.
- [45] S.M. Hasan, A. Shabbir, A. Shaikh, Z. Irshad, T.N. Khan, "Endoscopic Biopsy Proven Co-infection of *Helicobacter pylori* and *Giardia lamblia* in Adult Population of Karachi City," *Pakistan Journal of Medicine and Dentistry*, vol. 8, no. 02, p. 45-48, 2019.
- [46] A. Seid, Z. Tamir, B. Kasanew, M. Senbetay, "Co-infection of intestinal parasites and *Helicobacter pylori* among upper gastrointestinal symptomatic adult patients attending Mekanesealem Hospital, northeast Ethiopia," *BMC Research Notes*, vol. 11, p. 144, 2018, <https://doi.org/10.1186/s13104-018-3246-4>.
- [47] J. Ankarklev, E. Hestvik, M. Lebbad, J. Lindh, D.H. Kaddu-Mulindwa, J.O. Andersson, T. Tylleskär, "Common Coinfections of *Giardia intestinalis* and *Helicobacter pylori* in Non-Symptomatic Ugandan Children," *PLoS Neglected Tropical Diseases*, vol. 6, no. 8, p. e1780, 2012, doi:10.1371/journal.pntd.0001780.
- [48] A.A. Sabah, M.R. Gneidy, N.M.K. Saleh, "Prevalence of *Helicobacter pylori* infection among adult patients with different gastrointestinal parasites in Tanta City district," *Journal of the Egyptian Society of Parasitology*, vol. 45, p. 101-106, 2015, doi:10.12816/0010855.

- [49] N.F.M. Ahmed, T.E.M. Elfaki, M. Elsayid, "Prevalence rate of *Giardia lamblia*/*Helicobacter pylori* co-infections in Khartoum State, Sudan, International Journal of Scientific Technology Research, vol. 5, no. 3, p. 181–90, 2016.
- [50] J.G. Kuster, A.H. Vanviret, E.J. Kulpers, "Pathogenesis of *H. pylori* infections," Clinical Microbiology Review, vol. 19, no. 3, p. 449–90, 2006, doi: 10.1128/CMR.00054-05.

- [51] M.M. Abou El-Hoda, H.M. Osman, M.M. Rasha, N.L. Doudar, A.Y. Enany, "Impact of *Helicobacter pylori* infection on the activities of urease and lipase enzymes in patients with giardiasis," Journal of the Egyptian Public Health Association, vol. 82, no. 3–4, p. 273–82, 2007.

العدوى المختلطة لبكتيريا الحلزونية البوابية والطفيليات المعوية: تحديد عوامل الخطورة لدى مرضى عسر الهضم في مدينة المكلا، حضرموت، اليمن

عيزة علي بن حميد^{1,2} ، هدى محمد بارجاش¹

الملخص: غالباً ما يصاب المرضى الذين يعانون من عسر الهضم بعدوى بكتيريا الحلزونية البوابية، وهي أكثر شيوعاً في دول العالم الثالث. تمتلك بكتيريا الحلزونية البوابية القدرة على استعمار الغشاء المخاطي في المعدة، وغالباً ما تؤدي العدوى إلى تطور مضاعفات الجهاز الهضمي المختلفة. بالإضافة إلى ذلك، تُعد عدوى بكتيريا الحلزونية البوابية والطفيليات المعوية لدى البشر أمراً شائعاً. هدفت هذه الدراسة إلى تحديد مدى انتشار الطفيليات المعوية والعدوى المصاحبة للبكتيريا الحلزونية البوابية، بالإضافة إلى دراسة عوامل الخطورة المتعلقة بالعدوى لدى مرضى عسر الهضم في المستشفيات الرئيسية في مدينة المكلا، محافظة حضرموت، اليمن. في هذه الدراسة المقطعية تم جمع وتحليل 100 عينة من البراز. وكان الغرض هو الكشف عن عدوى بكتيريا الحلزونية البوابية باستخدام اختبار مستضد البراز المناعي السريع، وكذلك تحديد وجود عدوى طفيلية مترافقة مع عدوى بكتيريا الحلزونية البوابية. تم الحصول على بيانات المستهدفين باستخدام أداة الاستبيان، وتم فحص البيانات باستخدام نظام التحليل الإحصائي SPSS الإصدار 24، وتم استخدام تحليل الانحدار اللوجستي لتحديد متغيرات الخطورة التنبؤية المستقلة، واعتمدت القيمة المعنوية عند 0.05. كان إجمالي معدل انتشار عدوى بكتيريا الحلزونية البوابية بين مرضى عسر الهضم 30%. كان لدى الذكور معدل انتشار أعلى للعدوى الحلزونية (COR = 0.386، 95% CI = 0.160-0.934، P = 0.035). علاوة على ذلك، ارتبطت النتائج الإيجابية لعدوى بكتيريا الحلزونية البوابية بشكل كبير بمياه الشرب غير المفلترة (COR = 3.67، 95% CI = 1.436-9.363، P = 0.007)، وأعراض الارتجاع والحرقة (COR = 0.865، 95% CI = 0.034-0.536، P = 0.004)، واستخدام المضادات الحيوية (COR = 0.312، 95% CI = 0.125-0.780، P = 0.013). تشير هذه الدراسة إلى أن الأوليات المعوية المتحولة للحالة للنسيج والجيارديا اللامبليية كانت أكثر الطفيليات المرتبطة بعدوى بكتيريا الحلزونية البوابية شيوعاً في المرضى الذين يعانون من عسر الهضم دون وجود ارتباط معنوي. كانت متغيرات الجنس ومصادر المياه غير المفلترة وأعراض الارتجاع والحرقة واستخدام المضادات الحيوية هي أبرز عوامل الخطورة المرتبطة بعدوى بكتيريا الحلزونية البوابية المصاحبة لعدوى الطفيليات المعوية المتحولة للحالة للنسيج والجيارديا اللامبليية.

الكلمات المفتاحية: العدوى المشتركة، مرضى عسر الهضم، المتحولة للحالة للنسيج، الجيارديا اللامبليية، الحلزونية البوابية، حضرموت، الانتشار، اليمن