

Stability and boundedness for a kind of non-autonomous differential equations with constant delay

Cemil Tunç

Department of Mathematics, Faculty of Sciences, Yüzüncü Yıl University, 65080, Van, Turkey

Received: 2 Jan. 2012; Revised 8 May 2012; Accepted 17 Jul. 2012

Published online: 1 Jan. 2013

Abstract: We establish some new sufficient conditions to the uniform asymptotically stability and boundedness of the solutions for a kind of non-autonomous differential equations of third order with constant delay. By defining an appropriate Lyapunov functional, we prove two new theorems on the subject. Our results improve and form a complement to some known recent results in the literature.

Keywords: Uniform asymptotically stability; boundedness; nonlinear differential equation; third order; delay.

1. Introduction

Differential equations of higher order have proved to be valuable tools in the modeling of many phenomena in various fields of science and engineering. Indeed, we can find some applications such as nonlinear oscillations (Afuwape et al. [7], Andres [9] and Fridedrichs [22]), electronic theory Rauch [31], biological model and other models Cronin-Scanlon [12] and Gopalsamy [23], and etc.. Just as above, in the past few decades, there has been much attention paid to the discussion of the qualitative behaviors of the solutions of various nonlinear differential equations of third order without and with delay. For a comprehensive treatment of the subject, we refer the readers to the book of Reissig et al. [32] as a survey and the papers of Ademola and Arawomo ([1], [2]), Ademola et al. [3], Afuwape [5], Afuwape and Castellanos [6], Antiova [10], Chukwu [11], Ezeilo ([13]-[19]), Ezeilo and Tejumola ([20], [21]), Hara [24], Mehri and Shadman [25], Ogundare [26], Ogundare and Okecha [27], Omeike ([28]-[30]), Rauch [31], Swick ([33]-[35]), Tejumola ([36], [37]), Tunç ([38]-[55]), Tunç and Ateş ([56], [57]), Tunç and Ergören [58], Tunç and Tunç [59], Yao and Meng [60], Wu and Shi [62] and the references cited in these sources.

On the other hand, by a recent paper published in 2010, Ademola et al. [4] proved two new results on the boundedness and the uniform ultimate boundedness of the solutions of a nonlinear third order differential equation with

out delay ,

$$\ddot{x}(t) + f(\ddot{x}(t)) + g(\dot{x}(t)) + h(x(t)) = p(t, x(t), \dot{x}(t), \ddot{x}(t)). \tag{1}$$

In this paper, instead of Eq. (1), we consider the non-autonomous and nonlinear third order delay differential equations of the form

$$\begin{aligned} \ddot{x}(t) + f(t, x(t), x(t-r), \dot{x}(t), \dot{x}(t-r), \ddot{x}(t), \ddot{x}(t-r)) \\ + g(\dot{x}(t-r)) + h(x(t-r)) \\ = p(t, x(t), x(t-r), \dot{x}(t), \dot{x}(t-r), \ddot{x}(t), \ddot{x}(t-r)). \end{aligned} \tag{2}$$

We write Eq. (2) in system form as follows

$$\begin{aligned} \dot{x} &= y, \\ \dot{y} &= z, \\ \dot{z} &= -f(t, x, x(t-r), y, y(t-r), z, z(t-r)) - g(y) \\ &\quad - h(x) + \int_{t-r}^t g'(y(s))z(s)ds + \int_{t-r}^t h'(x(s))y(s)ds \\ &\quad + p(t, x, x(t-r), y, y(t-r), z, z(t-r)), \end{aligned} \tag{3}$$

in which the dots denote differentiation with respect to t , $t \in \mathbb{R}_+$, $\mathbb{R}_+ = [0, \infty)$; the functions f, g, h and p are

* Corresponding author: e-mail: cemtunc@yahoo.com

continuous in their respective arguments on $\mathbb{R}_+ \times \mathbb{R}^6$, \mathbb{R} , \mathbb{R} and $\mathbb{R}_+ \times \mathbb{R}^6$, respectively, and $\mathbb{R} = (-\infty, \infty)$. It is also assumed that the functions g and h are differentiable.

This paper is motivated by the papers of Ademola et al. [4] and that mentioned above. We define a new Lyapunov functional for the results to be established here. Then, using that Lyapunov functional, we discuss the uniform asymptotically stability and boundedness of the solutions of Eq. (2) for the cases, $p(\cdot) \equiv 0$ and $p(\cdot) \neq 0$, respectively. Obviously, the equation discussed in [4], Eq. (1), is a particular case of our equation, Eq. (2). Here, by this work, we improve the boundedness result obtained in [4] obtained for ordinary nonlinear differential Eq. (1) without delay to the nonlinear differential Eq. (2) with delay. It should be noted that Ademola et al. [4] discussed the boundedness of the solutions of Eq. (1). In addition to the boundedness of the solutions, we also discuss the uniform asymptotically stability of the solutions. Our results will be also different from that mentioned above. It should be noted that the basic reason to investigate these topics here is that functional differential equations play a key role in applied sciences. However, we only study the theoretical aspects of the topics here.

2. Preliminaries

Consider the functional differential equation

$$\dot{x}(t) = F(t, x_t), x_t = x(t + \theta), -r \leq \theta \leq 0, t \geq 0, \quad (4)$$

with $F : \mathbb{R}_+ \times C_H \rightarrow \mathbb{R}^n$ being continuous, $F(t, 0) = 0$, and we suppose that F takes closed bounded sets into bounded sets of \mathbb{R}^n . Here $(C, \|\cdot\|)$ is the Banach space of continuous function $\varphi : [-r, 0] \rightarrow \mathbb{R}^n$ with supremum norm, $r > 0$; C_H is the open H -ball in C ; $C_H := \{\varphi \in C([-r, 0], \mathbb{R}^n) : \|\varphi\| < H\}$.

Theorem 1. (Yoshizawa [61]). Suppose that there exists a continuous Lyapunov functional $V(t, \varphi)$ defined on $t \in \mathbb{R}_+$, $\|\varphi\| < H_1$, $0 < H_1 < H$, which satisfies the following conditions;

(i) $a(\|\varphi\|) \leq V(t, \varphi) \leq b(\|\varphi\|)$, where $a(r) \in CIP$ and $b(r) \in CIP$ (CIP denotes the families of continuous increasing, positive definite functions),

(ii) $\dot{V}(t, \varphi) \leq -c(\|\varphi\|)$, where $c(r)$ is continuous and positive for $r > 0$.

Then, the zero solution of (4) is uniform-asymptotically stable.

3. Main results

Let $p(\cdot) = 0$.

Our first result is given by the following theorem.

Theorem 2. Suppose that there exist positive constants a, b, b_1, c, δ_0 and L such that the following conditions hold:

(i)

$$h(0) = 0, \delta_0 \leq \frac{h(x)}{x}, (x \neq 0), h'(x) \leq c,$$

$$g(0) = 0, b \leq \frac{g(y)}{y} \leq b_1, (y \neq 0), |g'(y)| \leq L,$$

(ii)

$$a \leq \frac{f(t, x, x(t-r), y, y(t-r), z, z(t-r))}{z}, (z \neq 0).$$

Then, the zero solution of Eq. (2) is uniform-asymptotically stable provided that

$$r < \min \left\{ \frac{a\delta_0}{\alpha(c+L)}, \frac{7(\alpha a + ab - c)}{4(c\alpha + 2ac + c + aL)}, \frac{\alpha}{L(\alpha + a + 2) + c} \right\}.$$

The proof of Theorem 2 and that of the subsequent result depend on some certain fundamental properties of a continuously differentiable Lyapunov functional $V = V(x_t, y_t, z_t)$ defined by

$$\begin{aligned} 2V &= 2a \int_0^x h(\xi) d\xi + 2 \int_0^y g(\tau) d\tau + 2yh(x) + \alpha bx^2 \\ &+ (\alpha + a^2)y^2 + z^2 + 2\alpha axy + 2\alpha xz \\ &+ 2ayz + 2\lambda_1 \int_{-r}^0 \int_{t+s}^t y^2(\theta) d\theta ds + 2\lambda_2 \int_{-r}^0 \int_{t+s}^t z^2(\theta) d\theta ds, \end{aligned}$$

where α a is positive fixed constant satisfying

$$0 < \alpha < b - ca^{-1}, \quad (5)$$

and λ_1 and λ_2 are some positive constants which will be determined later in the proof.

Proof. We observe that the above Lyapunov functional can be rewritten as follows

$$2V = V_1 + V_2 + 2\lambda_1 \int_{-r}^0 \int_{t+s}^t y^2(\theta) d\theta ds + 2\lambda_2 \int_{-r}^0 \int_{t+s}^t z^2(\theta) d\theta ds, \quad (6)$$

where

$$V_1 = 2a \int_0^x h(\xi) d\xi + 2 \int_0^y g(\tau) d\tau + 2yh(x)$$

and

$$V_2 = \alpha bx^2 + (\alpha + a^2)y^2 + z^2 + 2\alpha axy + 2\alpha xz + 2ayz.$$

In view of assumption (i) of Theorem 2, we have $g(y) \geq by$ for all $y \neq 0$. Hence

$$\begin{aligned} 2 \int_0^y g(\tau) d\tau + 2yh(x) &\geq 2 \int_0^y b\tau d\tau + 2yh(x) \\ &= (by + h(x))^2 b^{-1} - b^{-1}h^2(x) \\ &\geq -b^{-1}h^2(x). \end{aligned}$$

Moreover, assumption (i) of Theorem 2 implies

$$\begin{aligned} 2a \int_0^x h(\xi) d\xi &= 2b^{-1} \int_0^x (ab - h'(\xi))h(\xi) d\xi + b^{-1}h^2(x) \\ &\geq (ab - c)b^{-1}\delta_0 x^2 + b^{-1}h^2(x). \end{aligned}$$

Combining the above estimates into V_1 , we obtain

$$V_1 \geq (ab - c)b^{-1}\delta_0 x^2 = k_1 x^2, \quad (k_1 = (ab - c)b^{-1}\delta_0 > 0).$$

We can also rearrange V_2 as follows

$$V_2 = XQ_0X^T,$$

where $X = (x, y, z)$, $Q_0 = \begin{pmatrix} \alpha b & \alpha a & \alpha \\ \alpha a & \alpha + a^2 & a \\ \alpha & a & 1 \end{pmatrix}$ and $\det Q_0 = \alpha^2(b - \alpha) > 0$ since $b - \alpha > 0$. Hence, we get

$$V_2 \geq \alpha^2(x^2 + y^2 + z^2).$$

Gathering the estimates for V_1 and V_2 into (6), it follows that

$$\begin{aligned} 2V &\geq (\alpha^2 + k_1)x^2 + \alpha^2(y^2 + z^2) \\ &+ 2\lambda_1 \int_{-r}^0 \int_{t+s}^t y^2(\theta) d\theta ds + 2\lambda_2 \int_{-r}^0 \int_{t+s}^t z^2(\theta) d\theta ds \\ &\geq k_2(x^2 + y^2 + z^2) \\ &+ 2\lambda_1 \int_{-r}^0 \int_{t+s}^t y^2(\theta) d\theta ds + 2\lambda_2 \int_{-r}^0 \int_{t+s}^t z^2(\theta) d\theta ds, \end{aligned}$$

where $k_2 = \alpha^2$. On the other hand, by the assumptions of Theorem 2 and the estimate $2|m| |n| \leq m^2 + n^2$, it can be easily obtained for a positive constant k_3 that

$$\begin{aligned} 2V &\leq k_3(x^2 + y^2 + z^2) \\ &+ 2\lambda_1 \int_{-r}^0 \int_{t+s}^t y^2(\theta) d\theta ds + 2\lambda_2 \int_{-r}^0 \int_{t+s}^t z^2(\theta) d\theta ds, \end{aligned}$$

so that

$$\begin{aligned} &\bar{k}_2(x^2 + y^2 + z^2) + \lambda_1 \int_{-r}^0 \int_{t+s}^t y^2(\theta) d\theta ds \\ &+ \lambda_2 \int_{-r}^0 \int_{t+s}^t z^2(\theta) d\theta ds \leq V \\ &\leq \bar{k}_3(x^2 + y^2 + z^2) + \lambda_1 \int_{-r}^0 \int_{t+s}^t y^2(\theta) d\theta ds \\ &+ \lambda_2 \int_{-r}^0 \int_{t+s}^t z^2(\theta) d\theta ds, \end{aligned}$$

where $\bar{k}_2 = 2^{-1}k_2$ and $\bar{k}_3 = 2^{-1}k_3$.

Thus, subject to the above discussion, it can be shown that condition (i) of Theorem 1 holds.

Let $(x, y, z) = (x(t), y(t), z(t))$ be a solution of (3). Along this solution, it follows from (6) and (3) that

$$\begin{aligned} \frac{dV}{dt} &= -\alpha x h(x) - \{ayg(y) - y^2 h'(x)\} - \alpha \{g(y) - by\}x \\ &- (\alpha x + ay + z)\{f(\cdot) - az\} \\ &+ (\alpha x + ay + z) \int_{t-r}^t h'(x(s))y(s) ds \\ &+ (\alpha x + ay + z) \int_{t-r}^t g'(y(s))z(s) ds + \lambda_1 y^2 r + \lambda_2 z^2 r \\ &- \lambda_1 \int_{t-r}^t y^2(s) ds - \lambda_2 \int_{t-r}^t z^2(s) ds + \alpha Y Q_1 Y^T, \end{aligned}$$

where $Y = (y, z)$, $Q_1 = \begin{pmatrix} a & 1 \\ 1 & 0 \end{pmatrix}$ and $\det Q_1 = -1$.

Making use of the assumptions of Theorem 2, we get

$$\begin{aligned} \frac{dV}{dt} &\leq -\frac{1}{2}\alpha\delta_0 x^2 - \frac{7}{8}(\alpha a + ab - c)y^2 - \frac{1}{2}\alpha z^2 \\ &+ (\alpha x + ay + z) \int_{t-r}^t h'(x(s))y(s) ds \\ &+ (\alpha x + ay + z) \int_{t-r}^t g'(y(s))z(s) ds + \lambda_1 y^2 r \\ &+ \lambda_2 z^2 r - \lambda_1 \int_{t-r}^t y^2(s) ds - \lambda_2 \int_{t-r}^t z^2(s) ds - W_i, \end{aligned}$$

($i = 1, 2, 3$), where

$$W_1 = \alpha \left\{ \frac{1}{4}\delta_0 x^2 + (g(y) - by)x + \frac{1}{16\alpha}(\alpha a + ab - c)y^2 \right\},$$

$$W_2 = \alpha \left\{ \frac{1}{4}\delta_0 x^2 + (f(\cdot) - az)x + \frac{1}{4}z^2 \right\},$$

$$W_3 = a \left\{ \frac{1}{16a} (\alpha a + ab - c) y^2 + (f(\cdot) - az) y + \frac{\alpha}{4a} z^2 \right\}.$$

Using the estimates

$$\{g(y) - by\}^2 < \frac{\delta_0(\alpha a + ab - c)}{16\alpha} y^2,$$

$$\{f(\cdot) - az\}^2 < \frac{\delta_0}{4\alpha^2} z^2,$$

$$\{f(\cdot) - az\}^2 < \frac{\alpha(\alpha + ab - c)}{16a^2} z^2,$$

respectively, we conclude

$$W_1 \geq \frac{\alpha}{16} \left(2\sqrt{\delta_0} |x| - \sqrt{\frac{\alpha a + ab - c}{\alpha}} |y| \right)^2 \geq 0,$$

$$W_2 \geq \frac{\alpha}{4} (\sqrt{\delta_0} |x| - |z|)^2 \geq 0,$$

$$W_3 \geq \frac{a}{16} \left(\sqrt{\frac{\alpha a + ab - c}{a}} |y| - 2\sqrt{\frac{\alpha}{a}} |z| \right)^2 \geq 0.$$

By noting the assumptions $0 < h'(x) \leq c$, $|g'(y)| \leq L$ and the estimate $2|m| |n| \leq m^2 + n^2$, it is followed that

$$(\alpha x + ay + z) \int_{t-r}^t h'(x(s)) y(s) ds \leq$$

$$\frac{r}{2} (\alpha x^2 + acy^2 + cz^2) + \frac{c}{2} (\alpha + a + 1) \int_{t-r}^t y^2(s) ds,$$

$$(\alpha x + ay + z) \int_{t-r}^t g'(y(s)) z(s) ds \leq$$

$$\frac{rL}{2} (\alpha x^2 + ay^2 + z^2) + \frac{L}{2} (\alpha + a + 1) \int_{t-r}^t z^2(s) ds.$$

In view of the above estimates, we get

$$\frac{dV}{dt} \leq -2^{-1} \{ \alpha \delta_0 - (\alpha c + \alpha L) r \} x^2$$

$$- \left\{ \frac{7}{8} (\alpha a + ab - c) - \frac{1}{2} (2\lambda_1 + ac + aL) r \right\} y^2$$

$$- 2^{-1} \{ \alpha - (2\lambda_2 + c + L) r \} z^2$$

$$- \{ \lambda_1 - 2^{-1} c (\alpha + a + 1) \} \int_{t-r}^t y^2(s) ds$$

$$- \{ \lambda_2 - 2^{-1} L (\alpha + a + 1) \} \int_{t-r}^t z^2(s) ds.$$

Let

$$\lambda_1 = \frac{1}{2} c (\alpha + a + 1)$$

and

$$\lambda_2 = \frac{1}{2} L (\alpha + a + 1).$$

Then, we get

$$\frac{dV}{dt} \leq -\frac{1}{2} \{ \alpha \delta_0 - (\alpha c + \alpha L) r \} x^2$$

$$- \left\{ \frac{7}{8} (\alpha a + ab - c) - \frac{1}{2} (c\alpha + 2ac + c + aL) r \right\} y^2$$

$$- \frac{1}{2} \{ \alpha - [L(\alpha + a + 2) + c] r \} z^2.$$

Hence, we conclude

$$\frac{d}{dt} V(x_t, y_t, z_t) \leq -k_4 x^2 - k_5 y^2 - k_6 z^2 \leq 0$$

for some positive constants k_4, k_5 and k_6 provided that

$$r < \min \left\{ \frac{\alpha \delta_0}{\alpha(c+L)}, \frac{7(\alpha a + ab - c)}{4(c\alpha + 2ac + c + aL)}, \frac{\alpha}{L(\alpha + a + 2) + c} \right\}.$$

This completes the proof of Theorem 2.

Let $p(\cdot) \neq 0$.

Our second result is given by the following theorem.

Theorem 3. We assume that all the assumptions of Theorem 2 and

$$|p(t, x, x(t-r), y, y(t-r), z, z(t-r))| \leq |q(t)|,$$

$$\int_0^t |q(s)| ds \leq P_0 < \infty$$

hold, where P_0 is a positive constant.

Then, there exists a positive constant M such that the solution $x(t)$ of Eq. (2) defined by the initial function

$$x(t) = \phi(t), x'(t) = \phi'(t), x''(t) = \phi''(t)$$

satisfies

$$|x(t)| \leq M, |x'(t)| \leq M, |x''(t)| \leq M$$

for all $t \geq t_0 \geq 0$, where $\phi \in C^2([t_0 - r, t_0], \mathfrak{R})$, provided that

$$r < \min \left\{ \frac{\alpha \delta_0}{\alpha(c+L)}, \frac{7(\alpha a + ab - c)}{4(c\alpha + 2ac + c + aL)}, \frac{\alpha}{L(\alpha + a + 2) + c} \right\}.$$

Proof. Let $(x(t), y(t), z(t))$ be any solution of (3). Then, by an easy calculation, it is obtained that

$$\begin{aligned} \frac{dV}{dt} &\leq -k_4x^2 - k_5y^2 - k_6z^2 + (\alpha x + ay + z) \\ &\quad \times p(t, x, x(t-r), y, y(t-r), z, z(t-r)) \\ &\leq \max(\alpha, a, 1)(|x| + |y| + |z|) |q(t)| \\ &\leq \delta_1(|x| + |y| + |z|) |q(t)| \\ &\leq \delta_1(3 + x^2 + y^2 + z^2) |q(t)|, \end{aligned}$$

where $\delta_1 \equiv \max(\alpha, a, 1)$. Making use of the estimate $x^2 + y^2 + z^2 \leq \bar{k}_2^{-1}V(x_t, y_t, z_t)$, we get

$$\frac{d}{dt}V(x_t, y_t, z_t) \leq \delta_2 |q(t)| + \delta_2 V(x_t, y_t, z_t) |q(t)|,$$

where $\delta_2 = \max(3\delta_1, \delta_1 \bar{k}_2^{-1})$. Integrating the above estimate from 0 to t , it follows that

$$\begin{aligned} V(x_t, y_t, z_t) &\leq V(x_0, y_0, z_0) + \delta_2 \int_0^t |q(s)| ds \\ &\quad + \delta_2 \int_0^t V(x_s, y_s, z_s) |q(s)| ds. \end{aligned}$$

Using the Gronwall-Reid-Bellman inequality, (see Ahmad and Rama Mohana Rao [8]), and the assumption

$$\int_0^t |q(s)| ds \leq P_0 < \infty,$$

we can conclude the result of Theorem 3.

References

- [1] T. A. Ademola; P. O. Arawomo, Stability and uniform ultimate boundedness of solutions of a third-order differential equation. *Int. J. Appl. Math.* 23 (2010), no. 1, 11-22.
- [2] T. A. Ademola; P. O. Arawomo, Stability and ultimate boundedness of solutions to certain third-order differential equations. *Appl. Math. E-Notes* 10 (2010), 61-69.
- [3] T. A. Ademola; R. Kehinde; O. M. Ogunlaran, A boundedness theorem for a certain third order nonlinear differential equation. *J. Math. Stat.* 4 (2008), no. 2, 88-93.
- [4] T. A. Ademola; M. O. Ogundiran; P. O. Arawomo; O. A. Adesina, Boundedness results for a certain third order nonlinear differential equation. *Appl. Math. Comput.* 216 (2010), no.10, 3044-3049.
- [5] A. U. Afuwape, Further ultimate boundedness results for a third-order nonlinear system of differential equations. *Boll. Un. Mat. Ital. C* (6) 4 (1985), no. 1, 347-361.
- [6] A. U. Afuwape; J. E. Castellanos, Asymptotic and exponential stability of certain third-order non-linear delayed differential equations: frequency domain method. *Appl. Math. Comput.* 216 (2010), no. 3, 940-950.
- [7] A. U. Afuwape; P. Omari; F. Zanolin, Nonlinear perturbations of differential operators with nontrivial kernel and applications to third-order periodic boundary value problems. *J. Math. Anal. Appl.* 143 (1989), no. 1, 35-56.
- [8] S. Ahmad; M. Rama Mohana Rao, Theory of ordinary differential equations. With applications in biology and engineering. Affiliated East-West Press Pvt. Ltd., New Delhi, 1999.
- [9] J. Andres, Periodic boundary value problem for certain nonlinear differential equations of the third order. *Math. Slovaca* 35 (1985), no. 3, 305-309.
- [10] E. S. Anitaova, On the boundedness of the solutions of a system of third-order differential equations. (Russian) *Vestnik Leningrad. Univ. Ser. Mat. Meh. Astronom* 19 (1964), no. 2, 149-151.
- [11] E. N. Chukwu, On the boundedness and the existence of a periodic solution of some nonlinear third order delay differential equation. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.* (8) 64 (1978), no. 5, 440-447.
- [12] J. Cronin-Scanlon, Some mathematics of biological oscillations. *SIAM Rev.* 19 (1977), no. 1, 100-138.
- [13] J. O. C. Ezeilo, On the boundedness of solutions of a certain differential equation of the third order. *Proc. London Math. Soc.* (3) 9 (1959) 74-114.
- [14] J. O. C. Ezeilo, A note on a boundedness theorem for some third order differential equations. *J. London Math. Soc.* 36 (1961) 439-444.
- [15] J. O. C. Ezeilo, A boundedness theorem for some non-linear differential equations of the third order. *J. London Math. Soc.* 37 (1962) 469-474.
- [16] J. O. C. Ezeilo, A boundedness theorem for a certain third-order differential equation. *Proc. London Math. Soc.* (3) 13 (1963) 99-124.
- [17] J. O. C. Ezeilo, An elementary proof of a boundedness theorem for a certain third order differential equation. *J. London Math. Soc.* 38 (1963) 11-16.
- [18] J. O. C. Ezeilo, A stability result for a certain third order differential equation. *Ann. Mat. Pura Appl.* (4) 72 (1966) 1-9.
- [19] J. O. C. Ezeilo, A generalization of a boundedness theorem for a certain third-order differential equation. *Proc. Cambridge Philos. Soc.* 63 (1967) 735-742.
- [20] J. O. C. Ezeilo; H. O. Tejumola, Boundedness and periodicity of solutions of a certain system of third-order non-linear differential equations. *Ann. Mat. Pura Appl.* (4) 74 (1966) 283-316.
- [21] J. O. C. Ezeilo; H. O. Tejumola, Boundedness theorems for certain third order differential equations. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.* (8) 55 (1973), 194-201 (1974).
- [22] K. O. Fridedrichs, On nonlinear vibrations of third order. *Studies in Nonlinear Vibration Theory*, pp. 65-103. Institute for Mathematics and Mechanics, New York University, 1946.
- [23] K. Gopalsamy, Stability and oscillations in delay differential equations of population dynamics. *Mathematics and its Applications*, 74. Kluwer Academic Publishers Group, Dordrecht, 1992.
- [24] T. Hara, On the uniform ultimate boundedness of the solutions of certain third order differential equations. *J. Math. Anal. Appl.* 80 (1981), no. 2, 533-544.

- [25] B. Mehri; D. Shadman, Boundedness of solutions of certain third order differential equation. *Math. Inequal. Appl.* 2 (1999), no. 4, 545-549.
- [26] B. S. Ogundare, On the boundedness and the stability results for the solutions of certain third order non-linear differential equations. *Kragujevac J. Math.* 29 (2006), 37-48.
- [27] B. S. Ogundare; G. E. Okecha, On the boundedness and the stability of solution to third order non-linear differential equations. *Ann. Differential Equations* 24 (2008), no. 1, 1-8.
- [28] M. O. Omeike, New result in the ultimate boundedness of solutions of a third-order nonlinear ordinary differential equation. *JIPAM. J. Inequal. Pure Appl. Math.* 9 (2008), no. 1, Article 15, 8 pp.
- [29] M. O. Omeike, Stability and boundedness of solutions of some non-autonomous delay differential equation of the third order. *An. Stiint. Univ. Al. I. Cuza Iași. Mat. (N.S.)* 55 (2009), suppl. 1, 49-58.
- [30] M. Omeike, New results on the asymptotic behavior of a third-order nonlinear differential equation. *Differ. Equ. Appl.* 2 (2010), no. 1, 39-51.
- [31] L. L. Rauch, Oscillation of a third order nonlinear autonomous system. *Contributions to the Theory of Nonlinear Oscillations*, pp. 39-88. *Annals of Mathematics Studies*, no. 20. Princeton University Press, Princeton, N.J., 1950.
- [32] R. Reissig; G. Sansone; R. Conti, *Non-linear Differential Equations of Higher Order*, Translated from the German. Noordhoff International Publishing, Leyden, 1974.
- [33] K. E. Swick, A boundedness result for the solutions of certain third order differential equations. *Ann. Mat. Pura Appl.* (4) 86 (1970) 169-180.
- [34] K. E. Swick, Asymptotic behavior of the solutions of certain third order differential equations. *SIAM J. Appl. Math.* 19 (1970) 96-102.
- [35] K. E. Swick, Boundedness and stability for a nonlinear third order differential equation. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur.* (8) 56 (1974), no. 6, 859-865.
- [36] H. O. Tejumola, On the boundedness and periodicity of solutions of certain third-order non-linear differential equations. *Ann. Mat. Pura Appl.* (4) 83 (1969) 195-212.
- [37] H. O. Tejumola, A note on the boundedness and the stability of solutions of certain third-order differential equations. *Ann. Mat. Pura Appl.* (4) 92 (1972), 65-75.
- [38] C. Tuñç, Uniform ultimate boundedness of the solutions of third-order nonlinear differential equations. *Kuwait J. Sci. Engrg.* 32 (2005), no. 1, 39-48.
- [39] C. Tuñç, Boundedness of solutions of a third-order nonlinear differential equation. *JIPAM. J. Inequal. Pure Appl. Math.* 6 (2005), no. 1, Article 3, 6 pp.
- [40] C. Tuñç, On the asymptotic behavior of solutions of certain third-order nonlinear differential equations. *J. Appl. Math. Stoch. Anal.* 2005, no. 1, 29-35.
- [41] C. Tuñç, On the boundedness of solutions of certain nonlinear vector differential equations of third order. *Bull. Math. Soc. Sci. Math. Roumanie (N. S.)* 49(97) (2006), no. 3, 291-300.
- [42] C. Tuñç, On the stability and boundedness of solutions of nonlinear vector differential equations of third order. *Nonlinear Anal.* 70 (2009), no. 6, 2232-2236.
- [43] Tuñç, C., On the qualitative behaviors of solutions to a kind of nonlinear third order differential equations with a retarded argument. *An. Stiint. Univ. "Ovidius" Constanta Ser. Mat.* 17 (2), (2009), 215-230.
- [44] C. Tuñç, Some new results on the boundedness of solutions of a certain nonlinear differential equation of third order. *Int. J. Nonlinear Sci.* 7 (2009), no. 2, 246-256.
- [45] C. Tuñç, Stability criteria for certain third order nonlinear delay differential equations. *Port. Math.* 66 (2009), no. 1, 71-80.
- [46] C. Tuñç, A new result on the stability of solutions of a nonlinear differential equation of third-order with finite lag. *Southeast Asian Bull. Math.* 33 (2009), no. 5, 947-958.
- [47] C. Tuñç, On the stability and boundedness of solutions to third order nonlinear differential equations with retarded argument. *Nonlinear Dynam.* 57 (2009), no. 1-2, 97-106.
- [48] C. Tuñç, The boundedness of solutions to nonlinear third order differential equations. *Nonlinear Dyn. Syst. Theory* 10 (2010), no. 1, 97-102.
- [49] C. Tuñç, On the stability and boundedness of solutions of nonlinear third order differential equations with delay. *Filomat* 24 (2010), no. 3, 1-10.
- [50] C. Tuñç, Some stability and boundedness conditions for non-autonomous differential equations with deviating arguments. *Electron. J. Qual. Theory Differ. Equ.* 2010, No. 1, 12 pp.
- [51] C. Tuñç, On some qualitative behaviors of solutions to a kind of third order nonlinear delay differential equations. *Electron. J. Qual. Theory Differ. Equ.* 2010, No. 12, 19 pp.
- [52] C. Tuñç, Bound of solutions to third-order nonlinear differential equations with bounded delay. *J. Franklin Inst.* 347 (2010), no. 2, 415-425.
- [53] C. Tuñç, On the existence of periodic solutions to nonlinear third order ordinary differential equations with delay. *J. Comput. Anal. Appl.* 12 (2010), no. 1-B, 191-201.
- [54] C. Tuñç, Asymptotic stable and bounded solutions of a kind of nonlinear differential equations with variable delay. *Funct. Differ. Equ.* 17 (2010), no. 3-4, 345-354.
- [55] C. Tuñç, Stability and bounded of solutions to non-autonomous delay differential equations of third order. *Nonlinear Dynam.* 62 (2010), no.4, 945-953.
- [56] C. Tuñç; M. Ateş, Stability and boundedness results for solutions of certain third order nonlinear vector differential equations. *Nonlinear Dynam.* 45 (2006), no. 3-4, 273-281.
- [57] C. Tuñç, M. Ateş, Boundedness and stability of solutions of a kind of nonlinear third order differential equations. *J. Appl. Func. Anal.* 5, no.3, (2010), 242-250.
- [58] C. Tuñç; H. Ergören, On boundedness of a certain non-linear differential equation of third order. *J. Comput. Anal. Appl.* 12 (2010), no. 3, 687-694.
- [59] C. Tuñç; E. Tuñç, New ultimate boundedness and periodicity results for certain third-order nonlinear vector differential equations. *Math. J. Okayama Univ.* 48 (2006), 159-172.
- [60] H. Yao; W. Meng, On the stability of solutions of certain non-linear third-order delay differential equations. *Int. J. Nonlinear Sci.* 6 (2008), no. 3, 230-237.
- [61] T. Yoshizawa, Stability theory by Liapunov's second method. *Publications of the Mathematical Society of Japan*, No. 9. The Mathematical Society of Japan, Tokyo 1966.
- [62] C. X. Wu; B. Shi, Globally asymptotic stability of a third-order nonlinear delay system. *(Chinese) Math. Pract. Theory* 38 (2008), no. 16, 227-230.

Cemil Tunç was born in Yeşilöz Köyü (Kalbulas), Horasan-Erzurum, Turkey, in 1958. He received the Ph. D. degree in Applied Mathematics from Erciyes University, Kayseri, in 1993. His research interests include qualitative behaviors of solutions of differential equations. At present he is Professor of Mathematics at Yüzüncü Yıl University, Van-Turkey.

