

2022

Relative Booster Ideals of Distributive p-algebras

M. M. Atallah

Department of Mathematics, Lawson State Community College, Birmingham, Alabama, 35221, USA,
eman.rezk@science.tanta.edu.eg

E. G. Rezk

*Department of Mathematics, Faculty of Science, Tanta University, Tanta, Egypt \\ Mathematical Sciences
Department, College of Science, Princess Nourah bint Abdulrahman University, P.O.Box 84428, Riyadh
11671, Saudi Arabia, eman.rezk@science.tanta.edu.eg*

Follow this and additional works at: <https://digitalcommons.aaru.edu.jo/isl>

Recommended Citation

M. Atallah, M. and G. Rezk, E. (2022) "Relative Booster Ideals of Distributive p-algebras," *Information Sciences Letters*: Vol. 11 : Iss. 6 , PP -.

Available at: <https://digitalcommons.aaru.edu.jo/isl/vol11/iss6/7>

This Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in Information Sciences Letters by an authorized editor. The journal is hosted on Digital Commons, an Elsevier platform. For more information, please contact rakan@aarj.edu.jo, marah@aarj.edu.jo, u.murad@aarj.edu.jo.

Relative Booster Ideals of Distributive p -algebras

M. M. Atallah¹ and E. G. Rezk^{2,3,*}

¹Department of Mathematics, Lawson State Community College, Birmingham, Alabama, 35221, USA

²Department of Mathematics, Faculty of Science, Tanta University, Tanta, Egypt

³Mathematical Sciences Department, College of Science, Princess Nourah bint Abdulrahman University, P.O.Box 84428, Riyadh 11671, Saudi Arabia

Received: 12 Mar. 2022, Revised: 22 May 2022, Accepted: 2 Jun 2022

Published online: 1 Nov. 2022

Abstract: In this article, the definition and characterization of relative booster ideals in distributive p -algebras are given. The relationship between disjunctive relative booster ideals and normal relative booster ideals is established in the distributive p -algebras. A lattice congruence relation defined via the relative boosters is given and its quotient lattice structure is obtained.

Keywords: Booster ideals, Congruence Relations, Disjunctive lattices, Distributive p -algebras, Ideals, Normal lattices, Stone algebras.

1 Introduction

O. Frink introduced the notion of pseudo-complementation of meet semi-lattices to generalize the class of Boolean algebras. Many authors introduced this concept of pseudo-complementation in different classes for example, the class of bounded lattices to get the class of p -algebras, consequently distributive p -algebras, Stone algebras, modular p -algebras, quasi-modular p -algebras. Ideals are investigated in the class of pseudo-complemented semi-lattice by T.S. Blyth, refer to [1]

A. Badawy and M. S. Raw [2] introduced the definition of booster ideals in distributive p -algebras in the sense of annihilators with pseudo-complemented elements and discussed some properties. W.H. Cornish studied a normal lattice [3] and announced that disjunctive normal distributive lattices are important in the compactification theory. Y. S. Pawar and S. S. Khopade studied the characterization of disjunctive ideals in 0-distributive lattices, see ([4], Theorem 3.11). Recently, many articles study properties of types of boosters in different classes of algebras, e.g., MS-algebra in [5] and [6], Stone almost distributive lattice in [7], [8] and so on.

In the present article, we generalize the concept of booster which was defined by A.Badawy et al.in [2]. The

relative booster ideals in distributive p -algebras is defined, and its important properties are shown. The behavior of disjunctive relative booster and normal relative booster ideals of distributive p -algebras is discussed, and hence the disjunctive booster and normal booster distributive p -algebras are given. Furthermore, the lattice congruence relations defined via the relative boosters is established.

After preliminaries in section 2, the definition and characterization of relative booster ideals of distributive p -algebra L are introduced. Some related properties are proved and the structure of relative booster ideals of L are given in section 3. In section 4, the relationship between disjunctive relative booster ideals and normal relative booster ideals in distributive p -algebra is established. Consequently, the disjunctive and normal booster distributive p -algebras are given, and every Stone algebra is a normal booster. Concluding by section 5, a relationship between the lattice congruence relation Θ and booster ideals on L is given. Moreover, the structure $(L/\Theta; \wedge, \vee, [0]\Theta, D(L))$ is investigated.

2 Preliminaries

In the present section, we recall some of the basic definitions and theorems that will be needed.

* Corresponding author e-mail: eman.rezk@science.tanta.edu.eg

Definition 1.[1] Let $(L; \wedge, \vee, 0, 1)$ be a bounded lattice. An element $a^\star \in L$ is called a pseudo-complement of a if for all $x \in L$

$$x \wedge a = 0 \text{ iff } x \leq a^\star$$

Therefore, in a bounded lattice the pseudo-complement of a is the largest element $x \in L$ with the property that $a \wedge x = 0$.

The bounded lattice L is said to be pseudo-complemented lattice (a p -algebra, for short) if every element in it have at most one pseudo-complement.

It is clear that $0^\star = 1$ and $1^\star = 0$. A p -algebra is called distributive (modular) if the underlying lattice is distributive (modular).

A distributive p -algebra is called a Stone algebra if it satisfies Stone identity

$$x^\star \vee x^{\star\star} = 1, \text{ for all } x \in L$$

Theorem 1.[1] Let L be a p -algebra. For all $a, b \in L$,

- (1) $a \leq b$ implies $b^\star \leq a^\star$,
- (2) $a \leq a^{\star\star}$,
- (3) $a^\star = a^{\star\star\star}$,
- (4) $(a \vee b)^\star = a^\star \wedge b^\star$,
- (5) $(a \wedge b)^{\star\star} = a^{\star\star} \wedge b^{\star\star}$,
- (6) $(a \wedge b)^\star \geq a^\star \vee b^\star$,
- (7) $(a \vee b)^{\star\star} = (a^{\star\star} \vee b^{\star\star})^{\star\star} \geq a^{\star\star} \wedge b^{\star\star}$.

Moreover, if L be a Stone algebra, then

- (8) $(a \wedge b)^\star = a^\star \vee b^\star$,
- (9) $(a \vee b)^{\star\star} = (a^{\star\star} \vee b^{\star\star})^{\star\star} = a^{\star\star} \wedge b^{\star\star}$.

Let L be a p -algebra, an element $a \in L$ is called a closed element if it satisfies the condition $a^{\star\star} = a$. The set of all closed elements of L denoted by $B(L)$. It is known that, $(B(L); \wedge, \vee, \star, 0, 1)$ forms a Boolean algebra where

$$a \nabla b = (a^\star \wedge b^\star)^\star$$

An element $d \in L$ is said to be a dense element if it satisfies the condition $d^\star = 0$. The set of all dense elements of L denoted by $D(L)$.

Let $I(L)$ be the set of all ideals of L . Then $(I(L); \subseteq)$ forms a complete lattice, and the lattice operations defined on $I(L)$ are given by, for $I, J \in I(L)$,

$$I \wedge J = I \cap J \text{ and } I \vee J = \{x \in L : x \geq a \vee b, \\ \text{for some } a \in I \text{ and } b \in J\}$$

It is known that, $(I(L); \wedge, \vee, (0), L)$ forms a complete lattice. In the special case, $I = \{a\}$, $a \in L$, the ideal $(a) = \{x \in L : x \leq a\}$ is called the principal ideal generated by $a \in L$. One can see that

$$(a) \wedge (b) = (a \wedge b) \text{ and } (a) \vee (b) = (a \vee b),$$

refer to [1].

Definition 2.[2] Let a be an element in a distributive p -algebra L . Then the booster $(a)^\Delta$ of a is the set:

$$(a)^\Delta = \{x \in L : x \wedge a^\star = 0\}.$$

Clearly, $(0)^\Delta = \{0\}$ and $(1)^\Delta = L$. In addition to boosters have the following properties:

Proposition 1.[2] Let L be a distributive p -algebra. Then for any $a, b \in L$:

- (1) $(a)^\Delta$ is an ideal of L containing a ,
- (2) $(a)^\Delta = (a^{\star\star})^\Delta = (a^{\star\star})^\Delta$,
- (3) $(a)^\Delta = \{a\}$ iff $a \in B(L)$,
- (4) $(a)^\Delta = L$ iff $a \in D(L)$,
- (5) If $a \in (b)^\Delta$, then $(a)^\Delta \subseteq (b)^\Delta$,
- (6) If $a \leq b$, then $(a)^\Delta \subseteq (b)^\Delta$,
- (7) $a^\star = b^\star$ iff $(a)^\Delta = (b)^\Delta$,
- (8) $(a)^\Delta = \{0\}$ iff $a = 0$.

Let $Con(L)$ be the set of all congruence relations on L , it is complete distributive lattice under set inclusion. For any $a \in L$, the congruence class of a with respect to the congruence Θ on L is given by $[a]_\Theta = \{x \in L : x \equiv a(\Theta)\}$. The set of all congruence classes is denoted by L/Θ . The operations are defined on L/Θ by $[a]_\Theta \wedge [b]_\Theta = [a \wedge b]_\Theta$ and $[a]_\Theta \vee [b]_\Theta = [a \vee b]_\Theta$ for all $[a]_\Theta, [b]_\Theta \in L/\Theta$. Then the algebraic system $(L/\Theta; \wedge, \vee, [0]_\Theta, [1]_\Theta)$ forms a lattice called a quotient lattice of L modulo Θ , refer to [1].

Throughout the present article, L stands for a distributive p -algebra unless otherwise mentioned

3 Relative Boosters

In the present section, the concept of relative booster and related properties are given.

Definition 3. Let A be a non-empty subset of L and I be a fixed ideal in L . A relative booster of A in an ideal I is the set A^{bo} defined as:

$$A^{bo} = \{x \in I : x \wedge a^\star = 0 \text{ for all } a \in A\}$$

where a^\star represents the pseudo-complemented element of a in L .

In the special case, if $A = \{a\}$, we write $(a)^{bo} = \{x \in I : x \wedge a^\star = 0\}$ is called the relative booster of a in I . If $I = (a)$ is a principal ideal generated by a in L , then $(0)^{bo} = \{0\}$, $(a)^{bo} = (a)$. If $I = L$, then $(a)^{bo}$ coincide with the booster $(a)^\Delta$ as in [2].

Lemma 1. If A is a non-empty subset of L and I is a fixed ideal of L . Then A^{bo} is an ideal of I .

Proof. Suppose that $x, y \in A^{bo}$, then

$$(x \vee y) \wedge a^\star = (x \wedge a^\star) \vee (y \wedge a^\star) = 0, \text{ for all } a \in A.$$

So, $x \vee y \in A^{bo}$. Let $x \in A^{bo}$ and $z \in I, z \leq x$, meeting both sides with a^\star , then

$$z \wedge a^\star \leq x \wedge a^\star \text{ implies that } z \wedge a^\star = 0, \text{ for all } a \in A.$$

Thus $z \in A^{bo}$ and A^{bo} is an ideal of I .

Corollary 1. The relative booster $(a)^{bo}$ of $a \in L$ is an ideal of I containing a .

Proposition 2. For any two elements $a, b \in L$ and a fixed ideal I in L . The following properties are true:

- (1) $(a)^{bo} \subseteq (a^{\star\star})$,
- (2) If $a \leq b$ then $(a)^{bo} \subseteq (b)^{bo}$,
- (3) $(a)^{bo} \cap (b)^{bo} = (a \wedge b)^{bo}$,
- (4) If $b \in (a)^{bo}$ then $(b)^{bo} \subseteq (a)^{bo}$,
- (5) $I = \cup_{a \in I} (a)^{bo}$,
- (6) If $a \in B(L)$, then $(a)^{bo} \subseteq (a)$,
- (7) If $a \in I \cap D(L)$, then $(a)^{bo} = I$,
- (8) $(a)^{bo} = (0)^{bo}$ iff $a = 0$,
- (9) If $a, b \in L$ and $(a)^{bo} = (b)^{bo}$, then :

$$(a \wedge c)^{bo} = (b \wedge c)^{bo} \quad \text{and} \quad (a \vee c)^{bo} = (b \vee c)^{bo} \quad \text{for all } c \in L.$$

Proof.

- (1) we have that, $(a)^{bo} = (a)^\Delta \cap I$. Then, $(a)^{bo} \subseteq (a)^\Delta = (a^{\star\star})$.
- (2) Let $a \leq b$ and $x \in (a)^{bo} \subseteq I$. Then $x \wedge b^{\star\star} \leq x \wedge a^{\star\star} = 0$. Accordingly $x \in (b)^{bo}$ and $(a)^{bo} \subseteq (b)^{bo}$.
- (3) Clearly, we have $(a \wedge b)^{bo} \subseteq (a)^{bo}, (b)^{bo}$. Now, let $x \in (a)^{bo} \cap (b)^{bo} \subseteq (a^{\star\star}) \cap (b^{\star\star})$, i.e., $x \leq a^{\star\star}, x \leq b^{\star\star}$. Then $x \leq a^{\star\star} \wedge b^{\star\star} = (a \wedge b)^{\star\star}$. Hence, $x \wedge (a \wedge b)^{\star\star} = 0$, and then $x \in (a \wedge b)^{bo}$. Therefore, $(a)^{bo} \cap (b)^{bo} = (a \wedge b)^{bo}$.
- (4) If $b \in (a)^{bo}$, then $b \wedge a^{\star\star} = 0$ which implies $a^{\star\star} \leq b^{\star\star}$. Now, let $x \in (b)^{bo}$. Then $x \wedge a^{\star\star} \leq x \wedge b^{\star\star} = 0$. As a result, $x \wedge a^{\star\star} = 0$ and so $x \in (a)^{bo}$.
- (5) Since $a \in (a)^{bo}$, it means that $I = \cup_{a \in I} \{a\} \subseteq \cup_{a \in I} (a)^{bo} \subseteq I$.
- (6) If $a \in B(L)$, then $a = a^{\star\star}$. By using (1), $(a)^{bo} \subseteq (a^{\star\star}) = (a)$.
- (7) If $a \in I \cap D(L)$, then $a^{\star\star} = 0$. Thus, for all $x \in I$ we get $x \wedge a^{\star\star} = 0$. Therefore $I = (a)^{bo}$.
- (8) $a \in (a)^{bo} = (0)^{bo} = \{0\}$ iff $a = 0$.
- (9) Let $(a)^{bo} = (b)^{bo}$, and $x \in (a \wedge c)^{bo}$. Then $(x \wedge a^{\star\star}) \vee (x \wedge c^{\star\star}) = x \wedge (a^{\star\star} \vee c^{\star\star}) \leq x \wedge (a \wedge c)^{\star\star} = 0$. Hence $(x \wedge a^{\star\star}) = (x \wedge c^{\star\star}) = 0$, it implies $x \in (a)^{bo} = (b)^{bo}$ and $x \in (c)^{bo}$. Accordingly $x \in (b)^{bo} \cap (c)^{bo} = (b \wedge c)^{bo}$ (from (3)). As a result, $(a \wedge c)^{bo} \subseteq (b \wedge c)^{bo}$. Similarly, $(b \wedge c)^{bo} \subseteq (a \wedge c)^{bo}$. If $x \in (a \vee c)^{bo}$, then $x \wedge (a \vee c)^{\star\star} = x \wedge a^{\star\star} \wedge c^{\star\star} = 0$. Thus, $x \wedge a^{\star\star} \in (a)^{bo} = (b)^{bo}$ and $x \wedge b^{\star\star} \wedge c^{\star\star} = x \wedge (b \vee c)^{\star\star} = 0$. Therefore, $x \in (b \vee c)^{bo}$ and $(a \vee c)^{bo} \subseteq (b \vee c)^{bo}$. Similarly, $(b \vee c)^{bo} \subseteq (a \vee c)^{bo}$.

Let the set $B^{bo}(I) = \{(a)^{bo} : a \in L\}$ be the set of all relative boosters of a fixed ideal I in L , and the operations on $B^{bo}(I)$ are defined as the following:

$$(a)^{bo} \cap (b)^{bo} = (a \wedge b)^{bo} \quad \text{and} \quad (a)^{bo} \sqcup (b)^{bo} = (a \vee b)^{bo} \quad \text{for all } (a)^{bo}, (b)^{bo} \in B^{bo}(I).$$

Theorem 2. The structure $(B^{bo}(I); \wedge, \sqcup, ^c, \{0\}, I)$ forms a Boolean algebra, where $((a)^{bo})^c = (a^{\star\star})^{bo}$ represents the complemented element of $(a)^{bo}$ in $B^{bo}(I)$.

Proof. Since for any $a, b \in L, (0)^{bo} \subseteq (a)^{bo}, (b)^{bo} \subseteq I$, and by Proposition 2(9), the meet operation in $B^{bo}(I)$ is well defined.

Now, for any $a, b \in L, a, b \leq a \vee b$, by using Proposition 2 (2), $(a)^{bo}, (b)^{bo} \subseteq (a \vee b)^{bo}$. That is $(a \vee b)^{bo}$ is an upper bound of both $(a)^{bo}$ and $(b)^{bo}$. Let $(c)^{bo}$ be another an upper bound of both $(a)^{bo}$ and $(b)^{bo}$. Then $(a)^{bo}, (b)^{bo} \subseteq (c)^{bo}$. Suppose that $x \in (a \vee b)^{bo}$, i.e., $x \wedge (a \vee b)^{\star\star} = x \wedge (a^{\star\star} \vee b^{\star\star}) = 0$. So, $x \wedge a^{\star\star} \in (b)^{bo} \subseteq (c)^{bo}$. It implies that $x \wedge a^{\star\star} \wedge c^{\star\star} = 0$ and $x \wedge c^{\star\star} \in (a)^{bo} \subseteq (c)^{bo}$. Hence $x \in (c)^{bo}$ and $(a \vee b)^{bo} \subseteq (c)^{bo}$. As a result, $(a \vee b)^{bo} = (a)^{bo} \sqcup (b)^{bo}$.

The distributive condition can be proved as the following, let $(a)^{bo}, (b)^{bo}, (c)^{bo} \in B^{bo}(I)$. Then

$$\begin{aligned} (a)^{bo} \cap [(b)^{bo} \sqcup (c)^{bo}] &= (a)^{bo} \cap (b \vee c)^{bo} = (a \wedge (b \vee c))^{bo} \\ &= ((a \wedge b) \vee (a \wedge c))^{bo} \\ &= (a \wedge b)^{bo} \sqcup (a \wedge c)^{bo} \\ &= ((a)^{bo} \cap (b)^{bo}) \sqcup ((a)^{bo} \cap (c)^{bo}). \end{aligned}$$

Hence, $B^{bo}(I)$ is a bounded distributive lattice.

Since for any $(a)^{bo} \in B^{bo}(I)$ there exists $(a^{\star\star})^{bo} \in B^{bo}(I)$ such that $(a)^{bo} \wedge (a^{\star\star})^{bo} = (a \wedge a^{\star\star})^{bo} = (0)^{bo} = \{0\}$, and if $(x)^{bo} \in B^{bo}(I)$ satisfies the property $(x)^{bo} \wedge (a)^{bo} = \{0\}$, we get $(x \wedge a)^{bo} = \{0\}$, and so $x \wedge a = 0$, that means $x \leq a^{\star\star}$, by Proposition 2(2), $(x)^{bo} \subseteq (a^{\star\star})^{bo}$. Hence, $(a^{\star\star})^{bo}$ represents the pseudo-complementation of $(a)^{bo} \in B^{bo}(I)$, and $(a)^{bo} \sqcup (a^{\star\star})^{bo} = (a \vee a^{\star\star})^{bo} = I$. Consequently, $((a)^{bo})^c = (a^{\star\star})^{bo}$ is the complement of $(a)^{bo} \in B^{bo}(I)$. Therefore, $B^{bo}(I)$ is a Boolean algebra.

Example 1. Consider L in Figure 1. The relative booster of the set $A = \{a, c\}$ in the ideal $I = \{\zeta\}$ is $A^{bo} = \{0, a\}$ and its booster is $A^\Delta = \{0, a, c\}$. Figure 2 represent the boolean algebra of the set of all relative doosters in the ideal I .




Figure 1: distributive p-algebra L




Figure 2: The Boolean algebra $(B^{bo}(I); \wedge, \sqcup, ^c, (0], I)$

Lemma 2. Let I be a fixed ideal of L and J be an ideal of $B^{bo}(I)$. Then:

- (1) For any ideal K of L , the set $\{(a)^{bo} : a \in K\}$ is an ideal of $B^{bo}(I)$,
- (2) The set $\{a \in L : (a)^{bo} \in J\}$ is an ideal of L .

Proof. (1) For all $a, b \in K$ we get $a \vee b \in K$. So, if $(a)^{bo}, (b)^{bo} \in \{(a)^{bo} : a \in K\}$, then $(a)^{bo} \sqcup (b)^{bo} = (a \vee b)^{bo} \in \{(a)^{bo} : a \in K\}$. Now let $(a)^{bo} \in \{(a)^{bo} : a \in K\}$ and $(c)^{bo} \in B^{bo}(I)$ such that $(c)^{bo} \subseteq (a)^{bo}$. Then $(c)^{bo} \cap (a)^{bo} = (c)^{bo} = (c \wedge a)^{bo} \in \{(a)^{bo} : a \in K\}$, since $a \wedge c \in K$. Thus $\{(a)^{bo} : a \in K\}$ is an ideal of $B^{bo}(I)$.

(2) If $(a)^{bo}, (b)^{bo} \in J$, implies $(a)^{bo} \sqcup (b)^{bo} = (a \vee b)^{bo} \in J$. Accordingly, if $a, b \in \{a \in L : (a)^{bo} \in J\}$, then $a \vee b \in \{a \in L : (a)^{bo} \in J\}$. Assume $a \in \{a \in L : (a)^{bo} \in J\}$ and $c \in L$ such that $c \leq a$. Hence $(c)^{bo} \cap (a)^{bo} = (c)^{bo} = (c \wedge a)^{bo} \in J$ and $c \in \{a \in L : (a)^{bo} \in J\}$. Therefore $\{a \in L : (a)^{bo} \in J\}$ is an ideal of L .

Definition 4. Let $I(L)$ be a lattice of ideals of L and $I(B^{bo}(I))$ be a lattice of the ideals of $B^{bo}(I)$, for a fixed ideal I of L . Then define the maps:

- $\phi : I(L) \rightarrow I(B^{bo}(I))$ as: $\phi(K) = \{(a)^{bo} : a \in K\}$, for any ideal K of L and,
- $\psi : I(B^{bo}(I)) \rightarrow I(L)$ as: $\psi(J) = \{a \in L : (a)^{bo} \in J\}$, for any ideal J of $B^{bo}(I)$.

Theorem 3. The maps ϕ and ψ are satisfy the following conditions:

- (1) Maps ϕ and ψ are isotones,
- (2) The map $\phi(\psi)$ is an identity map,
- (3) The map $\psi(\phi)$ is a clousure operator,
- (4) $\psi(\phi(K \cap G)) = \psi(\phi(K)) \cap \psi(\phi(G))$, for any two ideals $K, G \in I(L)$.

Proof. (1) If $K, G \in I(L)$ and $K \subseteq G$, then $\phi(K) \subseteq \phi(G)$. On the other side, suppose $J, H \in I(B^{bo}(I))$ such that $J \subseteq H$. Then $\{a \in L : (a)^{bo} \in J\} \subseteq \{a \in L : (a)^{bo} \in H\}$. I.e., $\psi(J) \subseteq \psi(H)$.

(2) Assume $J \in I(B^{bo}(I))$ implies $\psi(J) \in I(L)$. Accordingly, $(a)^{bo} \in J$ iff $a \in \psi(J)$ iff $(a)^{bo} \in \phi(\psi(J))$. Therefore $\phi(\psi(J)) = J$.

(3) Now we show that $\psi(\phi)$ is extensive, isotone and idempotent.

- (i) Let $a \in K \in I(L)$. Then $(a)^{bo} \in \phi(K)$. It means $a \in \psi(\phi(K))$ and $K \subseteq \psi(\phi(K))$. Hence $\psi(\phi)$ is extensive.

(ii) Let $K, G \in I(L)$ such that $K \subseteq G$ and $a \in \psi(\phi(K))$. Then $(a)^{bo} \in \phi(K)$ and there exist $b \in K \subseteq G$ such that $(a)^{bo} = (b)^{bo} \in \phi(G)$. Since $\phi(G)$ is an ideal of $B^{bo}(I)$, hence $a \in \psi(\phi(G))$. As a result, $\psi(\phi(K)) \subseteq \psi(\phi(G))$ and $\psi(\phi)$ is isotone.

(iii) We get $\psi(\phi(K)) \subseteq \psi(\phi(\psi(\phi(K))))$. Since $\phi(\psi(\phi(K))) \in I(B^{bo}(I))$. Conversely, suppose $a \in \psi(\phi(\psi(\phi(K))))$ implies $(a)^{bo} \in \phi(\psi(\phi(K)))$. Let $b \in \psi(\phi(K))$ and $(b)^{bo} = (a)^{bo} \in \phi(K)$. Therefore $a \in \psi(\phi(K))$ and it is idempotent.

(4) Suppose $a \in \psi(\phi(K)) \cap \psi(\phi(G))$. It means $(a)^{bo} \in \phi(K) \cap \phi(G) = \phi(K \cap G)$. So, $a \in \psi(\phi(K \cap G))$ and $\psi(\phi(K \cap G)) \supseteq \psi(\phi(K)) \cap \psi(\phi(G))$. The convers is clear.

4 Disjunctive and Normal Relative Boosters

In the present section, the behavior of a disjunctive relative booster ideal and a normal relative booster ideal in distributive p -algebra are established.

Definition 5. An ideal I of L is a disjunctive relative booster if it satisfies

$$(a)^{bo} = (b)^{bo} \text{ implies that } a = b \text{ for all } a, b \in L.$$

Definition 6. An ideal I of L is a normal relative booster ideal if it satisfies

$$(a)^{bo} \vee (b)^{bo} = (a \vee b)^{bo} \text{ for all } a, b \in L$$

In particular case $I = L$, we can say that L is called disjunctive (or normal) booster.

Example 2. Consider L in Figure (1). The ideal $(\tau]$ is disjunctive and normal relative booster while the ideal $(\eta]$ is normal relative booster but not disjunctive. The ideal $(\xi]$ is neither disjunctive nor normal relative booster in L .

Theorem 4. Let I be a disjunctive relative booster ideal of L . Then I can be embedded into $B^{bo}(I)$.

Proof. Suppose that $\varphi : I \rightarrow B^{bo}(I)$ is defined by: $\varphi(a) = (a)^{bo}$, for all $a \in I$. It is easy to see that φ is well defined. Moreover,

$$\varphi(a \wedge b) = \varphi(a \wedge b)^{bo} = (a)^{bo} \cap (b)^{bo} = \varphi(a) \cap \varphi(b), \text{ and}$$

$$\varphi(a \vee b) = (a \vee b)^{bo} = (a)^{bo} \sqcup (b)^{bo} = \varphi(a) \sqcup \varphi(b).$$

Also, φ is one-to-one follows from the disjunctive of I .

Corollary 2. L is disjunctive booster iff L is a Boolean algebra.

A characterization of a normal relative booster ideal is given in the following results:

Theorem 5. An ideal I of L is normal relative booster iff $B^{bo}(I)$ is a Boolean sub-lattice of $I(L)$.

*Proof.*The necessary condition follows from Theorem 2. Conversely, suppose that $B^+(I)$ is a Boolean sub-lattice of $I(L)$ and for all $(a)^{bo}, (b)^{bo} \in B^{bo}(I)$

$$(a)^{bo} \vee (b)^{bo} = (c)^{bo}, \text{ for some } c \in L.$$

So, $(a)^{bo}, (b)^{bo} \subseteq (c)^{bo}$, and $a, b \leq a \vee b$, by Proposition 2(2), $(a)^{bo}, (b)^{bo} \subseteq (a \vee b)^{bo}$. Thus, $(c)^{bo}$ and $(a \vee b)^{bo}$ are upper bounds of both $(a)^{bo}$ and $(b)^{bo}$. Since L is distributive, then $I(L)$ so, and

$$\begin{aligned} (c)^{bo} \wedge (a \vee b)^{bo} &= (c)^{bo} \wedge [(a)^{bo} \sqcup (b)^{bo}] \\ &= [(c)^{bo} \wedge (a)^{bo}] \sqcup [(c)^{bo} \wedge (b)^{bo}] \\ &= (a)^{bo} \sqcup (b)^{bo} = (a \vee b)^{bo}. \end{aligned}$$

Therefore $(a \vee b)^{bo} \subseteq (c)^{bo}$ and $(a)^{bo} \vee (b)^{bo} = (a \vee b)^{bo}$.

Corollary 3. L is normal booster iff $B^{bo}(I)$ is a Boolean sub-lattice of $I(L)$.

Theorem 6. If L is a Stone algebra, then L is a normal booster.

Proof. Since, $a, b \leq a \vee b$, then by Proposition 2(2), $(a)^\Delta, (b)^\Delta \subseteq (a \vee b)^\Delta$ and so $(a)^\Delta \vee (b)^\Delta \subseteq (a \vee b)^\Delta$. Now, suppose that $x \in (a \vee b)^\Delta$, i.e., $x \wedge (a \vee b)^\star = x \wedge a^\star \wedge b^\star = 0$. It implies that $(x \wedge a^\star \wedge b^\star) = (x \wedge a^\star) \wedge (b^\star) = (0)$. Hence $(x \wedge a^\star) = (x) \wedge (a^\star) \subseteq (b^\star)^\star = (b^{\star\star}) = (b)^\Delta$. But $(x) \wedge (a^{\star\star}) \subseteq (a^{\star\star}) = (a)^\Delta$. Accordingly, $((x) \wedge (a^{\star\star})) \vee ((x) \wedge (a^\star)) = (x) \wedge ((a^\star) \vee (a^{\star\star})) = (x) \wedge (a^\star \vee a^{\star\star}) = (x) \subseteq (a)^\Delta \vee (b)^\Delta$. Therefore, $x \in (a)^\Delta \vee (b)^\Delta$ and $(a)^\Delta \vee (b)^\Delta = (a \vee b)^\Delta$

Corollary 4. A disjunctive booster distributive p -algebra is a normal booster

The convers of the above corollary is not true. For example, the Stone algebra of three elements chain is normal booster but not disjunctive booster.

5 Congruence Relations Via The Relative Boosters

In the present section a lattice congruence defined via the relative boosters is studied and some properties are derived.

Definition 7.[6] A Gilvenko congruence relation Φ on L is defined as

$$a \equiv b(\Phi) \text{ iff } a^{\star\star} = b^{\star\star}.$$

Theorem 7. Let Θ be a binary relation defined on L by the rule, for a fixed element $z \in L$ and fixed ideal I of L ,

$$a \equiv b(\Theta) \text{ iff } (a)^{bo} \cap (z)^{bo} = (b)^{bo} \cap (z)^{bo} \text{ for all } a, b \in L.$$

Then Θ is a lattice congruence on L .

Proof. It is clear that, Θ is an equivalence relation on L . Now, let $a \equiv b(\Theta)$ and $c \equiv d(\Theta)$ for all $a, b, c, d \in L$ by the definition, we get

$$(a)^{bo} \cap (z)^{bo} = (b)^{bo} \cap (z)^{bo} \text{ and } (c)^{bo} \cap (z)^{bo} = (d)^{bo} \cap (z)^{bo},$$

thus

$$\begin{aligned} (a \wedge c)^{bo} \cap (z)^{bo} &= (a)^{bo} \cap (c)^{bo} \cap (z)^{bo} = (b)^{bo} \cap (d)^{bo} \cap (z)^{bo} \\ &= (b \wedge d)^{bo} \cap (z)^{bo}, \text{ and} \\ (a \vee c)^{bo} \cap (z)^{bo} &= ((a \vee c) \wedge z)^{bo} = ((a \wedge z) \vee (c \wedge z))^{bo} \\ &= (a \wedge z)^{bo} \sqcup (c \wedge z)^{bo} = (b \wedge z)^{bo} \sqcup (d \wedge z)^{bo} \\ &= ((b \wedge z) \vee (d \wedge z))^{bo} = ((b \vee d) \wedge z)^{bo} \\ &= (b \vee d)^{bo} \cap (z)^{bo} \end{aligned}$$

Therefore, $a \wedge c \equiv b \wedge d(\Theta)$ and $a \vee c \equiv b \vee d(\Theta)$. That is, Θ is a lattice congruence relation on L .

Example 3. Consider relative boosters of elements of lattice L in ideal $I = \{\zeta\}$. Closed curves in Figure 3 represent the congruence classes of relation Θ which is defined as $a \equiv b(\Theta)$ iff $(a)^{bo} \cap (\alpha)^{bo} = (b)^{bo} \cap (\alpha)^{bo}$ for a fixed element $\alpha \in L$ and for all $a, b \in L$.




Figure 3: Congruence classes of relation Θ

Corollary 5. If the fixed element z in L , $z \in I \cap D(L)$ is dense element, then the congruence relation Θ on L defined as

$$a \equiv b(\Theta) \text{ iff } (a)^{bo} = (b)^{bo} \text{ for all } a, b \in L$$

In particular case, whenever $I = L$, we have

$$a \equiv b(\Theta) \text{ iff } (a)^\Delta = (b)^\Delta \text{ for all } a, b \in L$$

A Gilvinko congruence is an example of this congruence.

Theorem 8. Let Θ be a congruence relation on L defined as

$$a \equiv b(\Theta) \text{ iff } (a)^{bo} = (b)^{bo} \text{ for all } a, b \in L.$$

Then, the structure $(L/\Theta; \wedge, \vee, ^c, (0), D(L))$ forms a Boolean algebra.

Proof. From the definition of the class $[a]_{\Theta} = \{x \in L : (x)^{bo} = (a)^{bo}\}$, we have

$$[0]_{\Theta} = \{x \in L : (x)^{bo} = (0)^{bo}\} = \{0\} = (0).$$

So, $[0]_{\Theta}$ is the smallest congruence class of L . Also, let $d \in D(L)$,

$$[d]_{\Theta} = \{x \in L : (x)^{bo} = (d)^{bo}\} = \{x \in L : (x)^{bo} = (1)\} \\ = \{x \in L : x^{\star} = (0)\} = D(L).$$

$D(L)$ is a class of L/Θ , since the set of dense elements $D(L)$ forms a filter of L , then for any $x \in L$ and $d \in D(L)$, we have $x \vee d \in D(L)$. Thus $[x]_{\Theta} \vee [d]_{\Theta} = [x \vee d]_{\Theta} = D(L)$. Therefore, $D(L)$ is the largest element of L/Θ . Moreover,

$$[a]_{\Theta} \wedge [a^{\star}]_{\Theta} = [a \wedge a^{\star}]_{\Theta} = [0]_{\Theta} \\ \text{and } [a]_{\Theta} \vee [a^{\star}]_{\Theta} = [a \vee a^{\star}]_{\Theta} = D(L).$$

It means that the complement $([a]_{\Theta})^c$ of $[a]_{\Theta}$ is $[a^{\star}]_{\Theta}$. As a result, L/Θ is a Boolean algebra.

Example 4. Consider relative boosters of elements of lattice L in ideal $I = (\zeta)$. Closed curves in Figure 4 represent the congruence classes of relation Θ which is defined as $a \equiv b(\Theta)$ iff $(a)^{bo} \cap (\delta)^{bo} = (b)^{bo} \cap (\delta)^{bo}$ for all $a, b \in L$. It is equivalent to $a \equiv b(\Theta)$ iff $(a)^{bo} = (b)^{bo}$, since $\delta \in D(L)$. Figure 5 shows the Boolean algebra of L/Θ .




Figure 4: Congruence classes of relation Θ




Figure 5: The Boolean algebra $(L/\Theta; \wedge, \vee, ^c, (0), D(L))$

Corollary 6. Let $B^{bo}(I)$ be a Boolean algebra of relative boosters of fixed ideal I of L and Θ be a congruence defined as

$$a \equiv b(\Theta) \text{ iff } (a)^{bo} = (b)^{bo} \text{ for all } a, b \in L.$$

Then, $L/\Theta \cong B^{bo}(I)$.

6 Conclusion

- (1) The structure of relative booster ideals of a distributive p -algebra L forms a Boolean algebra, and hence it is a generalization of the booster ideals of L . Moreover, there is a closure operator generated by $I(B^{bo}(I))$ on the lattice ideal of L .
- (2) A distributive p -algebra L is disjunctive booster iff L is a Boolean algebra, and L is normal booster iff the structure of booster ideals is a sublattice of the structure of all ideals of L .
- (3) If the congruence relation Θ defined via the relative booster ideal of a distributive p -algebra L . Then the structure $(L/\Theta; \wedge, \vee, ^c, (0), D(L))$ forms a Boolean algebra isomorphic to the Boolean algebra of relative boosters.

Conflicts of Interests

The authors declare that they have no conflicts of interests

References

- [1] G.Gratzer, *Lattice Theory: Foundation*, Springer Basel AG,(2011).
- [2] A. Badawy and M.S. Rao, σ -Ideal of Distributive p -algebras, *Chamchuri Journal of Mathematics*,**6**, 17-30(2014).
- [3] W.H.Cornish, Annulets and α -Ideals In A Distributive Lattice, *J. Aust., Math. Soc.*,**15**, 70-77(1973).
- [4] Y.S.Pawar and S.S.Khopade, α -ideals and annihilator ideals in 0-distributive lattices, *Acta Univ. Palacki. Olomuc. Fac. Res. Nat. Mathematica*, **49**(1), 63-74(2010).
- [5] A. Badawy and R. El-Fawal, Closure Filters of Decomposable MS-Algebras, *Southeast Asian Bulletin of Mathematics*, **44**(2), 177-194(2020).
- [6] R. Noorbhasha, B. Ravikumar and K.P. Shum, Closure Filters and Prime Fuzzy Closure Filters of MS-Algebra, *Korean J. Math.*, **28**(3), 509-524(2020).
- [7] N. Rafi, Ravi Kummur Bandaru and G.C. Rao, e -filters in a Stone Almost Lattices, *Chamehuri Journal of Mathematics*,**7**, 16-28(2015).
- [8] N. Rafi and B. Ravikumar, β Prime Spectrum of Stone Almost Distributive Lattices, *Discussiones Mathematicae General Algebra and Applications*, **40**, 311-326(2020).