

New Symmetric Identities Involving q -Zeta Type Functions

Serkan Araci^{1,*}, Armen Bagdasaryan², Cenap Özel³ and H. M. Srivastava⁴

¹ Faculty of Economics, Administrative and Social Sciences, Hasan Kalyoncu University, 27410 Gaziantep, Turkey

² Institute for Control Sciences, Russian Academy of Sciences, 65 Profsoyuznaya Street, 117997 Moscow, Russian Federation

³ Department of Mathematics, Abant İzzet Baysal University, TR-14268 Bolu, Turkey

⁴ Department of Mathematics and Statistics, University of Victoria, Victoria, British Columbia V8W 3R4, Canada

Received: 17 Oct. 2013, Revised: 15 Jan. 2014, Accepted: 16 Jan. 2014

Published online: 1 Nov. 2014

Abstract: The main object of this paper is to obtain several symmetric properties of the q -zeta type functions. As applications of these properties, we give some new interesting identities for the modified q -Genocchi polynomials. Finally, our applications are shown to lead to a number of interesting results which we state in the present paper.

Keywords: Genocchi numbers and polynomials, generating functions, q -Genocchi polynomials, Euler and q -Euler zeta functions, q -zeta type functions.

1 Introduction

Throughout this paper, we use the following standard notations:

$$\mathbb{N} := \{1, 2, 3, \dots\} \quad \text{and} \quad \mathbb{N}_0 := \{0, 1, 2, \dots\} = \mathbb{N} \cup \{0\}.$$

Also, as usual, \mathbb{R} denotes the set of real numbers and \mathbb{C} denotes the set of complex numbers.

The Genocchi polynomials $G_n(x)$ and the Genocchi numbers $G_n := G_n(0)$ are given by the following generating functions:

$$\sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!} = \left(\frac{2t}{e^t + 1} \right) e^{xt} \quad (1)$$

$$\sum_{n=0}^{\infty} G_n \frac{t^n}{n!} = \frac{2t}{e^t + 1}, \quad (|t| < \pi),$$

respectively. In particular, the second generating function in (1) can be restated as follows:

$$e^{Gt+t} + e^{Gt} = 2t$$

by using the *umbral* (symbolic) convention exhibited by $G^n := G_n$. By utilizing the Taylor-Maclaurin expansion,

one finds that

$$(G+1)^n + G_n = \begin{cases} 2 & (n=1) \\ 0 & (\text{otherwise}). \end{cases} \quad (2)$$

It follows from (2) that (see, for details, [29])

$$G_1 = 1, \quad G_2 = -1, \quad G_3 = 0, \quad G_4 = 1, \quad G_5 = 0, \\ G_6 = -3, \quad G_7 = 0, \quad G_8 = 17, \dots$$

and (in general)

$$G_{2n+1} = 0 \quad (n \in \mathbb{N}).$$

The history of the Genocchi polynomials $G_n(x)$ and the Genocchi numbers G_n can be traced back to the Italian mathematician, Angelo Genocchi (1817–1889). From Genocchi to the present time, the Genocchi polynomials and the Genocchi numbers have been extensively studied in many different contexts in such branches of Mathematics as, for instance, Elementary Number Theory, Complex Analytic Number Theory, Homotopy Theory (especially stable Homotopy groups of spheres), Differential Topology (especially differential structures on spheres), Theory of Modular Forms (especially Eisenstein series), p -Adic Analytic Number

* Corresponding author e-mail: mtsrxn@hotmail.com

Theory (especially p -adic L -functions) and Quantum Physics (especially quantum groups). Investigations involving the Genocchi polynomials and their associated combinatorial relations have received considerable attention in recent years (see, for details, [1], [2], [3], [6], [7], [8], [16], [24], [30] and [26]).

Araci *et al.* [6] studied the modified q -Genocchi polynomials which are given by the following generating function:

$$F_q(x, t) = \sum_{n=0}^{\infty} \mathcal{G}_{n,q}(x) \frac{t^n}{n!} = [2]_q t \sum_{m=0}^{\infty} (-q)^m e^{(x+[m]_q)t}, \quad (3)$$

where the q -number $[\lambda]_q$ is given by

$$[\lambda]_q := \frac{1 - q^\lambda}{1 - q} \quad (0 < q < 1; \lambda \in \mathbb{C}), \quad (4)$$

so that, obviously, we have

$$\lim_{q \rightarrow 1^-} \{[\lambda]_q\} = \lambda \quad (\lambda \in \mathbb{C}).$$

In the case when $x = 0$ in (3), it leads to

$$\mathcal{G}_{n,q}(0) := \mathcal{G}_{n,q},$$

that is, to the modified q -Genocchi numbers $\mathcal{G}_{n,q}$. In addition to this, by letting $q \rightarrow 1^-$, $\mathcal{G}_{n,q}$ reduces to the Genocchi numbers G_n :

$$\lim_{q \rightarrow 1^-} \{\mathcal{G}_{n,q}\} = G_n.$$

The Genocchi numbers $G_n(x)$ possess a number of important properties and are well known in Number Theory. In fact, these numbers are related to the values at negative integers of the Euler Zeta function defined by (see [20], [22], [23], [28], [29]; see also [31])

$$\zeta(s, x) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(x+n)^s} = \Phi(-1, s, x) \quad (5)$$

$$(s \in \mathbb{C}; x \in \mathbb{C} \setminus \mathbb{Z}_0^-; \mathbb{Z}_0^- := \{0, -1, -2, \dots\}),$$

where $\Phi(z, s, a)$ denotes the widely- and extensively-studied general Hurwitz-Lerch Zeta function defined by (see, for example, [28, p. 121 *et seq.*] and [29, p. 194 *et seq.*]; see also [27], [31] and [32])

$$\Phi(z, s, a) := \sum_{n=0}^{\infty} \frac{z^n}{(n+a)^s} \quad (6)$$

$$(a \in \mathbb{C} \setminus \mathbb{Z}_0^-; s \in \mathbb{C} \text{ when } |z| < 1; \Re(s) > 1 \text{ when } |z| = 1).$$

Recently, Kim [20] defined the q -Euler Zeta function as follows:

$$\zeta_q(s, x) = [2]_q \sum_{n=0}^{\infty} \frac{(-1)^n q^n}{[x+n]_q^s}, \quad (s \in \mathbb{C}; x \in \mathbb{C} \setminus \mathbb{Z}_0^-). \quad (7)$$

On the other hand, Araci *et al* [6] introduced the q -Zeta type function $\tilde{\zeta}_q(s, x)$ which is slightly different from Kim's q -Zeta function $\zeta_q(s, x)$ defined by (7):

$$\begin{aligned} \tilde{\zeta}_q(s, x) &:= \frac{1}{\Gamma(s)} \int_0^{\infty} t^{s-2} \{-F_q(x, -t)\} dt \\ &= [2]_q \sum_{n=0}^{\infty} \frac{(-1)^n q^n}{(x+[n]_q)^s}, \quad (8) \end{aligned}$$

$$(s \in \mathbb{C}; x \neq -[n]_q \quad (n \in \mathbb{N}_0)),$$

where $F_q(x, -t)$ is given by (3). From (3) and (8), we find that (see [6])

$$\tilde{\zeta}_q(-n, x) = \frac{\mathcal{G}_{n+1,q}(x)}{n+1} \quad (n \in \mathbb{N}_0). \quad (9)$$

Moreover, by using (7) and (8), we have

$$q^{-sx} \tilde{\zeta}_q(s, q^{-1}[x]_{q^{-1}}) = \zeta_q(s, x). \quad (10)$$

The Zeta functions play a crucially important rôle in Analytic Number Theory and have applications in such areas as (for example) physics, probability theory, applied statistics, complex analysis, mathematical physics, p -adic analysis and other related areas. In particular, the Zeta functions occur within the concept of knot theory, quantum field theory, applied analysis and number theory (see [9], [10], [11], [20], [21], [22], [23], [28] and [31]).

The distribution formula for the modified q -Genocchi polynomials is given by (see [6])

$$\begin{aligned} \mathcal{G}_{n,q}(q^a [d]_q x) &:= \\ \frac{[d]_q^{n-1}}{[d]_q - q} \sum_{a=0}^{d-1} (-1)^a q^{a(n+1)} \mathcal{G}_{n,q^d} \left(x + \frac{[a]_q}{q^a [d]_q} \right), \quad (11) \end{aligned}$$

$$\text{for } d \equiv 1 \pmod{2}.$$

Araci *et al.* [8] derived several new identities for the (h, q) -Genocchi polynomials and gave symmetric identities of the (h, q) -Zeta type functions. Yuan He [14] gave symmetric identities for Carlitz's q -Bernoulli numbers (see also [12] and [13]). Kim also obtained symmetric identities for the q -Euler polynomials and derived the symmetric identities for the q -Euler Zeta function (see [15]). Simsek [25] gave the complete sum of products of (h, q) -extension of the Euler polynomials. Bagdasaryan investigated the elementary evaluation of the Zeta function and presented a real analytic approach to the values of the Riemann Zeta function (see, for details, [9] and [10]).

The symmetric identity of the Genocchi polynomials is given by Theorem 1 below (see [11]).

Theorem 1. Let a and b be odd integers. Then we have

$$\sum_{i=0}^m \binom{m}{i} a^{i-1} b^{m-i} G_i(bx) S_{m-i}(a) = \sum_{i=0}^m \binom{m}{i} b^{i-1} a^{m-i} G_i(ax) S_{m-i}(b), \quad (12)$$

where

$$S_m(a) := \sum_{j=0}^{a-1} (-1)^j j^m. \quad (13)$$

Motivated essentially by some of the aforementioned investigations, the fundamental aim of this paper is to generalize Theorem 1 by presenting an interesting and potentially useful extension of the symmetry identity (12) to hold true for the modified q -Genocchi polynomials arising from the above-mentioned q -Zeta type functions. Several other related results are also considered.

2 The q -Zeta Type Functions

In this section, we recall from (8) that

$$\tilde{\zeta}_q(s, x) = [2]_q \sum_{m=0}^{\infty} \frac{(-1)^m q^m}{(x + [m]_q)^s}. \quad (14)$$

In view of (10), we consider (14) in the following form:

$$q^{-asbx-sbj} \tilde{\zeta}_{q^a} \left(s, q^{-a} \left[bx + \frac{bj}{a} \right]_{q^{-a}} \right) = [2]_{q^a} \sum_{m=0}^{\infty} \frac{(-1)^m q^{ma}}{\left[m + bx + \frac{bj}{a} \right]_{q^a}^s}. \quad (15)$$

For non-negative integers k and i such that $m = bk + i$ with $0 \leq i \leq b-1$, if we suppose that $a \equiv 1 \pmod{2}$ and $b \equiv 1 \pmod{2}$, then we have

$$\begin{aligned} & q^{-asbx-sbj} \tilde{\zeta}_{q^a} \left(s, q^{-a} \left[bx + \frac{bj}{a} \right]_{q^{-a}} \right) \\ &= [a]_q^s [2]_{q^a} \sum_{m=0}^{\infty} \frac{(-1)^m q^{ma}}{\left[ma + abx + bj \right]_{q^a}^s} \\ &= [a]_q^s [2]_{q^a} \sum_{m=0}^{\infty} \sum_{i=0}^{b-1} \frac{(-1)^{i+mb} q^{(i+mb)a}}{\left[(i+mb)a + abx + bj \right]_{q^a}^s} \\ &= [a]_q^s [2]_{q^a} \sum_{i=0}^{b-1} (-1)^i q^{ia} \sum_{m=0}^{\infty} \frac{(-1)^m q^{mba}}{\left[ab(m+x) + ai + bj \right]_{q^a}^s}, \end{aligned} \quad (16)$$

which readily yields

$$\begin{aligned} & \sum_{j=0}^{a-1} (-1)^j q^{jb} q^{-asbx-sbj} \tilde{\zeta}_{q^a} \left(s, q^{-a} \left[bx + \frac{bj}{a} \right]_{q^{-a}} \right) \\ &= [a]_q^s [2]_{q^a} \sum_{j=0}^{a-1} (-1)^j q^{jb} \sum_{i=0}^{b-1} (-1)^i q^{ia} \times \\ & \quad \times \sum_{m=0}^{\infty} \frac{(-1)^m q^{mba}}{\left[ab(m+x) + ai + bj \right]_{q^a}^s}. \end{aligned} \quad (17)$$

Upon replacing a by b and j by i in (16), we get

$$\begin{aligned} & q^{-asbx-as} \tilde{\zeta}_{q^b} \left(s, q^{-b} \left[ax + \frac{ai}{b} \right]_{q^{-b}} \right) \\ &= [b]_q^s [2]_{q^b} \sum_{j=0}^{a-1} (-1)^j q^{jb} \times \\ & \quad \times \sum_{m=0}^{\infty} \frac{(-1)^m q^{mba}}{\left[ab(m+x) + ai + bj \right]_{q^a}^s}. \end{aligned} \quad (18)$$

Thus, by applying (17) in (18), we obtain the following theorem.

Theorem 2. For any odd integers a and b , we have

$$\begin{aligned} & \frac{[2]_{q^b}}{[a]_q^s} \sum_{i=0}^{a-1} (-1)^i q^{ib(1-s)} \tilde{\zeta}_{q^a} \left(s, q^{-a} \left[bx + \frac{bi}{a} \right]_{q^{-a}} \right) \\ &= \frac{[2]_{q^a}}{[b]_q^s} \sum_{i=0}^{b-1} (-1)^i q^{ia(1-s)} \times \\ & \quad \times \tilde{\zeta}_{q^b} \left(s, q^{-b} \left[ax + \frac{ai}{b} \right]_{q^{-b}} \right). \end{aligned} \quad (19)$$

Remark 1. Upon setting $b = 1$ in Theorem 2, we easily deduce that

$$\begin{aligned} & \tilde{\zeta}_q \left(s, q^{-1} [ax]_{q^{-1}} \right) = \\ & \frac{[2]_q}{[2]_{q^a} [a]_q^s} \sum_{i=0}^{a-1} (-1)^i q^{i(1-s)} \tilde{\zeta}_{q^a} \left(s, q^{-a} \left[x + \frac{i}{a} \right]_{q^{-a}} \right). \end{aligned} \quad (20)$$

Taking $a = 2$ in (20), we derive the following Corollary.

Corollary 1. For any odd integer a , we have

$$\begin{aligned} & \tilde{\zeta}_q \left(s, q^{-1} [2x]_{q^{-1}} \right) = \frac{[2]_q}{[2]_{q^2} [2]_q^s} \times \\ & \quad \left[\tilde{\zeta}_{q^2} \left(s, q^{-2} [x]_{q^{-2}} \right) - q^{b(1-s)} \tilde{\zeta}_{q^2} \left(s, q^{-2} \left[x + \frac{1}{2} \right]_{q^{-2}} \right) \right]. \end{aligned} \quad (21)$$

Remark 2. If we take $s = -n$ in Theorem 2, we get the following symmetric property of the modified q -Genocchi polynomials.

Theorem 3. For any odd integers a and b , we have

$$\begin{aligned}
 & [2]_{q^b} [a]_q^{n-1} \sum_{i=0}^{a-1} (-1)^i q^{ib(n+1)} \times \\
 & \quad \times \mathcal{G}_{n,q^a} \left(q^{-a} \left[bx + \frac{bi}{a} \right]_{q^{-a}} \right) = \\
 & = [2]_{q^a} [b]_q^{n-1} \sum_{i=0}^{b-1} (-1)^i q^{ia(n+1)} \times \\
 & \quad \times \mathcal{G}_{n,q^b} \left(q^{-b} \left[ax + \frac{ai}{b} \right]_{q^{-b}} \right). \quad (22)
 \end{aligned}$$

We now take $b = 1$ and replace x by $\frac{x}{a}$ in Theorem 3. We thus restate the distribution formula for the modified q -Genocchi polynomials as follows:

$$\begin{aligned}
 \mathcal{G}_{n,q} \left(-[-x]_q \right) &= \frac{[2]_q}{[2]_{q^a}} [a]_q^{n-1} \sum_{i=0}^{a-1} (-1)^i q^{i(n+1)} \times \\
 & \quad \times \mathcal{G}_{n,q^a} \left(q^{-a} \left[\frac{x+i}{a} \right]_{q^{-a}} \right), \quad (2 \nmid a). \quad (23)
 \end{aligned}$$

We next find from (3) that

$$\begin{aligned}
 \sum_{n=0}^{\infty} \mathcal{G}_{n,q}(x+y) \frac{t^n}{n!} &= [2]_q t \sum_{m=0}^{\infty} (-q)^m e^{(x+y+[m]_q)t} \\
 &= \left(\sum_{m=0}^{\infty} y^m \frac{t^m}{m!} \right) \left(\sum_{n=0}^{\infty} \mathcal{G}_{n,q}(x) \frac{t^n}{n!} \right),
 \end{aligned}$$

which, by applying the Cauchy product, yields

$$\begin{aligned}
 \sum_{n=0}^{\infty} \mathcal{G}_{n,q}(x+y) \frac{t^n}{n!} &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} \mathcal{G}_{k,q}(x) y^{n-k} \right) \frac{t^n}{n!}. \quad (24)
 \end{aligned}$$

Thus, by comparing the coefficients of $\frac{t^n}{n!}$ on both sides of this last equation (24), we get the following Corollary.

Corollary 2. For $n \in \mathbb{N}_0$, we obtain

$$\mathcal{G}_{n,q}(x+y) = \sum_{k=0}^n \binom{n}{k} \mathcal{G}_{k,q}(x) y^{n-k}. \quad (25)$$

By using Theorem 3 and (25), we can derive Theorem 4 below.

Theorem 4. For any odd integers a and b , we have

$$\begin{aligned}
 & [2]_{q^b} [a]_q^{n-1} \sum_{k=0}^n \binom{n}{k} [a]_{q^{-1}}^{k-n} [b]_{q^{-1}}^{n-k} \times \\
 & \quad \times \mathcal{G}_{k,q} \left(q^{-a} [bx]_{q^{-a}} \right) S_{n-k;q^{-b}}^{(-n-1)}(a) = \\
 & = [2]_{q^a} [b]_q^{n-1} \sum_{k=0}^n \binom{n}{k} [b]_{q^{-1}}^{k-n} [a]_{q^{-1}}^{n-k} \times \\
 & \quad \times \mathcal{G}_{k,q} \left(q^{-b} [ax]_{q^{-b}} \right) S_{n-k;q^{-a}}^{(-n-1)}(b). \quad (26)
 \end{aligned}$$

where

$$S_{m;q}^{(j)}(a) := \sum_{i=0}^{a-1} (-1)^i q^{ji} [i]_q^m. \quad (27)$$

Remark 3. Letting $q \rightarrow 1^-$ in Theorem 4, we can deduce the known symmetry identity (12).

3 Concluding Remarks and Observations

In this article, we have derived several symmetric properties of the q -Zeta type function $\tilde{\zeta}_q(s, x)$ defined by (8). As applications of these properties, we give new interesting symmetry identities for the modified q -Genocchi polynomials $\mathcal{G}_{n,q}(x)$ which are defined by (3). In the limit when $q \rightarrow 1^-$, this last result (Theorem 4) is shown to yield the known symmetry identity (12) for the Genocchi polynomials $G_n(x)$.

References

- [1] S. Araci, Novel identities involving Genocchi numbers and polynomials arising from applications from umbral calculus, *Applied Mathematics and Computation*, doi: 10.1016/j.amc.2014.01.013 (In press).
- [2] S. Araci, M. Acikgoz, E. Sen, Theorems on Genocchi polynomials of higher order arising from Genocchi basis, *Taiwanese Journal of Mathematics*, doi: 10.11650/tjm.18.2014.3006 (In press).
- [3] S. Araci, Novel identities for q -Genocchi numbers and polynomials, *J. Function Spaces Appl.*, **2012**, Article ID 214961, 1–13 (2012).
- [4] S. Araci, M. Acikgoz, A. Bagdasaryan and E. Şen, The Legendre polynomials associated with Bernoulli, Euler, Hermite and Bernstein polynomials, *Turkish J. Anal. Number Theory*, **1**, 1–3 (2013).
- [5] S. Araci, M. Acikgoz and E. Şen, On the extended Kim's p -adic q -deformed fermionic integrals in the p -adic integer ring, *J. Number Theory*, **133**, 3348–3361 (2013).
- [6] S. Araci, A. Bagdasaryan, E. Agyüz and M. Acikgoz, On the modified q -Genocchi numbers and polynomials and their applications, arXiv:1311.5992 [math.NT].
- [7] S. Araci, D. Erdal and J. J. Seo, A study on the fermionic p -adic q -integral representation on \mathbb{Z}_p associated with weighted q -Bernstein and q -Genocchi polynomials, *Abstr. Appl. Anal.*, **2011**, Article ID 649248, 1–10 (2011).
- [8] S. Araci, J. J. Seo and D. Erdal, New construction weighted (h, q) -Genocchi numbers and polynomials related to Zeta type function, *Discrete Dynamics Nature Soc.*, **2011**, Article ID 487490, 1–7 (2011).
- [9] A. Bagdasaryan, An elementary and real approach to values of the Riemann zeta function, *Phys. Atom. Nucl.*, **73**, 251–254 (2010).
- [10] A. Bagdasaryan, Elementary evaluation of the zeta and related functions: An approach from a new perspective, *AIP Conf. Proc.*, **1281**, 1094–1097 (2010).
- [11] E. Cetin, M. Acikgoz, I. N. Cangul and S. Araci, A note on the (h, q) -zeta-type function with weight α , *J. Inequal. Appl.*, **2013**, Article ID 100, 1–6 (2013).

- [12] J. Choi, P. J. Anderson and H. M. Srivastava, Some q -extensions of the Apostol-Bernoulli and the Apostol-Euler polynomials of order n , and the multiple Hurwitz zeta function, *Appl. Math. Comput.*, **199**, 723–737 (2008).
- [13] J. Choi, P. J. Anderson and H. M. Srivastava, Carlitz's q -Bernoulli and q -Euler numbers and polynomials and a class of q -Hurwitz zeta functions, *Appl. Math. Comput.*, **215**, 1185–1208 (2009).
- [14] Y. He, Symmetric identities for Calitz's q -Bernoulli numbers and polynomials, *Adv. Difference Equations*, **2013**, Article ID 246, 1–10 (2013).
- [15] D. S. Kim, T. Kim, S.-H. Lee and J. J. Seo, Symmetric identities of the q -Euler Polynomials, *Adv. Stud. Theoret. Phys.*, **7**, 1149–1155 (2013).
- [16] T. Kim, On the q -extension of Euler and Genocchi numbers, *J. Math. Anal. Appl.*, **326**, 1458–1465 (2007).
- [17] T. Kim, On the analogs of Euler numbers and polynomials associated with p -adic q -integral on \mathbb{Z}_p at $q = 1$, *J. Math. Anal. Appl.*, **331**, 779–792 (2007).
- [18] T. Kim, Symmetry p -adic invariant integral on \mathbb{Z}_p for Bernoulli and Euler polynomials, *J. Difference Equations Appl.*, **14**, 1267–1277 (2008).
- [19] T. Kim, Some identities on the q -Euler polynomials of higher order and q -Stirling numbers by the fermionic p -adic integral on \mathbb{Z}_p , *Russian J. Math. Phys.*, **16**, 484–491 (2009).
- [20] T. Kim, Note on the Euler q -zeta functions, *J. Number Theory*, **129**, 1798–1804 (2009).
- [21] T. Kim, S. H. Lee, H. H. Han and C. S. Ryoo, On the values of the weighted q -Zeta and L -functions, *Discrete Dynamics Nature Soc.*, **2011**, Article ID 476381, 1–7 (2011).
- [22] C. S. Ryoo and T. Kim, An analogue of the zeta function and its applications, *Appl. Math. Lett.*, **19**, 1068–1072 (2006).
- [23] L.-C. Jang, The q -analogue of twisted Lerch type Euler Zeta functions, *Bull. Korean Math. Soc.*, **47**, 1181–1188 (2010).
- [24] S.-H. Rim, J.-H. Jin, E.-J. Moon and S.-J. Lee, On multiple interpolation functions of the q -Genocchi polynomials, *J. Inequal. Appl.*, **2010**, Article ID 351419, 1–13 (2010).
- [25] Y. Simsek, Complete sum of products of (h, q) -extension of Euler polynomials and numbers, *J. Difference Equations Appl.*, **16**, 1331–1348 (2010).
- [26] H. M. Srivastava, Some generalizations and basic (or q -) extensions of the Bernoulli, Euler and Genocchi polynomials, *Appl. Math. Inform. Sci.*, **5**, 390–444 (2011).
- [27] H. M. Srivastava, A new family of the λ -generalized Hurwitz-Lerch Zeta functions with applications, *Appl. Math. Inform. Sci.*, **8**, 1–16 (2014).
- [28] H. M. Srivastava and J. Choi, *Series Associated with the Zeta and Related Functions*, Kluwer Academic Publishers, Dordrecht, Boston and London, (2001).
- [29] H. M. Srivastava and J. Choi, *Zeta and q -Zeta Functions and Associated Series and Integrals*, Elsevier Science Publishers, Amsterdam, London and New York, (2012).
- [30] H. M. Srivastava, B. Kurt and Y. Simsek, Some families of Genocchi type polynomials and their interpolation functions, *Integral Transforms Spec. Funct.*, **23**, 919–938 (2012); see also Corrigendum, *Integral Transforms Spec. Funct.*, **23**, 939–940 (2012).
- [31] H. M. Srivastava, M.-J. Luo and R. K. Raina, New results involving a class of generalized Hurwitz-Lerch zeta functions and their applications, *Turkish J. Anal. Number Theory*, **1**, 26–35 (2013).
- [32] H. M. Srivastava, R. K. Saxena, T. K. Pogány and R. Saxena, Integral and computational representations of the extended Hurwitz-Lerch Zeta function, *Integral Transforms Spec. Funct.*, **22**, 487–506 (2011).



Serkan Araci was born in Hatay, Turkey, on October 1, 1988. He received his B.S. and M.S. degrees in mathematics from the University of Gaziantep, Gaziantep, Turkey, in 2010 and 2013, respectively. Additionally, the title of his M.S. thesis is "Bernstein

polynomials and their reflections in analytic number theory" and, for this thesis, he received the Best Thesis Award of 2013 from the University of Gaziantep. He has published more than 50 papers in reputed international journals. His research interests include p -adic analysis, analytic theory of numbers, q -series and q -polynomials, and theory of umbral calculus. Araci is an editor and a referee for many international journals and serves as an Editor in Chief of both Turkish Journal of Analysis and Number Theory, Journal for Algebra and Number Theory Academia, and Journal of Analysis and Number Theory.



Armen Bagdasaryan is a senior research scientist of Russian Academy of Sciences in Moscow. He received his Ph.D. degree in applied mathematics from Russian Academy of Sciences and M.Sc. degree with honors in mathematics from Yerevan State University, Armenia. He

was thrice a recipient of a special research scholarship for outstanding young scientists who is under the age of thirty-five, awarded by Russian Academy of Sciences. He is a member of American Mathematical Society, Society for Industrial and Applied Mathematics, and Institute of Physics UK. He was a member of the IASTED Technical Committee on Modeling and Simulation, Canada. He serves on the editorial boards of several international journals in pure and applied mathematics, computer science. His research interests are in the areas of number theory, analysis, combinatorics, mathematical physics, topology, computer science and complex systems theory.



Cenap Özel received the PhD degree from University of Glasgow, Scotland, 1998. He is the recipient of Sedat Simavi Foundation National Science Laudable prize from Turkish Journalists Society 2005, and Young Outstanding Scientist prize from Turkish Science Academy 2007. His

interests are Algebraic Topology, especially complex cobordism of infinite dimensional manifolds, Lie theory, Quantum Mechanics and Mathematical Physics, Differential Geometry, and Number theory. He was a member of Abant Izzet Baysal University AIBU, Bolu Turkey since 1998 till 2014. Now he is a professor at University of Dokuz Eylul, Izmir Turkey.



H. M. Srivastava
For the author's biographical and other professional details, the interested reader should look into the following Web Site:

<http://www.math.uvic.ca/faculty/harimsri>