

2023

Extended Maxwell–Chern–Simons Lagrangian Density in Riemann–Liouville Fractional Derivatives

Amer D. Al-Oqali

Physics Department, Mutah University, Al-Karak, Jordan, aoqali@mutah.edu.jo

Follow this and additional works at: <https://digitalcommons.aaru.edu.jo/isl>

Recommended Citation

D. Al-Oqali, Amer (2023) "Extended Maxwell–Chern–Simons Lagrangian Density in Riemann–Liouville Fractional Derivatives," *Information Sciences Letters*: Vol. 12 : Iss. 2 , PP -.

Available at: <https://digitalcommons.aaru.edu.jo/isl/vol12/iss2/33>

This Article is brought to you for free and open access by Arab Journals Platform. It has been accepted for inclusion in Information Sciences Letters by an authorized editor. The journal is hosted on Digital Commons, an Elsevier platform. For more information, please contact rakan@aarj.edu.jo, marah@aarj.edu.jo, u.murad@aarj.edu.jo.

Extended Maxwell–Chern–Simons Lagrangian Density in Riemann–Liouville Fractional Derivatives

Amer D. Al-Oqali

Physics Department, Mutah University, Al-Karak, Jordan

Received: 2 Apr. 2022, Revised: 22 Jun. 2022, Accepted: 12 Jul. 2022

Published online: 1 Feb. 2023

Abstract: The Hamiltonian formulation for higher derivatives is reformulated using fractional derivatives. More precisely, the extended Maxwell–Chern–Simons Lagrangian density is reformulated using the Riemann–Liouville fractional derivative. The equations of motion resulting from the extended Maxwell–Chern–Simons Lagrangian density are obtained. Furthermore, the Hamiltonian of the system is constructed. When fractional derivatives are replaced by integer order derivatives, the classical results are obtained.

Keywords: Fractional derivatives; Extended Maxwell–Chern–Simons Lagrangian density; Lagrangian formulation; Hamiltonian formulation; Euler–Lagrange equations

1 Introduction

Fractional derivatives, which have many applications in physics, science, engineering, control, economics, mechanics, and other fields, can be viewed as a generalization of classical calculus [1, 2, 3, 4]. Actually, recently, physicists have applied a fractional derivatives approach in order to deal with problems that cannot be solved by classical methods. Therefore, fractional calculus appears as one of the most influential and commonly applicable methods for describing and explaining a set of complicated physical systems. [5, 6, 7]. Generally, higher derivative theories are an essential field in theoretical physics. Higher derivative theories were first proposed to remove infinities associated to point particles [8]. They can enhance ultraviolet features in quantum field theories [9]. Furthermore, they have been obtained from string theory [10] and non commutative theory [11] and have been utilized in electrodynamics [12], dark energy physics [13], and Lee-Wick models [14].

The extended Maxwell–Chern–Simons model is an attractive instance of higher derivative field theory [15] [16]. Generally, the Maxwell–Chern–Simons theory is a (2+1) dimensional field model, which defines charged fermions interactions with each other and with topologically massive propagating photons [17].

The fractional derivatives method was used by Jarab’ah and Nawafleh [18] to investigate nonconservative systems with second order Lagrangian. They obtained the fractional Hamilton’s equations for nonconservative systems. The generated equations were similar to fractional Euler–Lagrange equations.

Alawaida et al. [19] developed the Hamiltonian formulation of continuous field systems with third order using the fractional derivatives method. They generated the fractional Euler and fractional Hamilton equations for these systems from the fractional variational principle. In addition, Al-Oqali [20] reformulated Podolsky’s Lagrangian density in fractional form using the Riemann–Liouville fractional derivative. He gained the equations of motion using the fractional Euler–Lagrange equation. The Hamiltonian and the energy stress tensor are also generated in fractional form from the Lagrangian density.

The remainder of this paper is organized as follows: Section 2 presents some basic definitions of fractional derivatives. In Section 3, the Hamiltonian formulation of higher derivative field theories is reformulated using the Riemann–Liouville fractional derivative. In Section 4, the fractional form of the Euler–Lagrange equation of the extended Maxwell–Chern–Simons Lagrangian density is obtained. In section 5, the Hamiltonian density of the extended Maxwell–Chern–Simons model in fractional form is constructed. Finally, Section 6 is devoted to the paper’s conclusion.

* Corresponding author e-mail: aoqali@mutah.edu.jo

2 Definitions of fractional derivatives

In this section, we briefly present some basic definitions utilized in this work. The left and right Riemann–Liouville fractional derivative is defined as [21]:

The left Riemann–Liouville fractional derivative

$${}_a D_x^\alpha f(x) = \frac{1}{\Gamma(n-\alpha)} \left(\frac{d}{dx} \right)^n \int_a^x (x-\tau)^{n-\alpha-1} f(\tau) d\tau \quad (1)$$

The right Riemann–Liouville fractional derivative

$${}_x D_b^\alpha f(x) = \frac{1}{\Gamma(n-\alpha)} \left(\frac{-d}{dx} \right)^n \int_a^x (\tau-x)^{n-\alpha-1} f(\tau) d\tau. \quad (2)$$

where α represents the order of the derivative such that $n-1 \leq \alpha < n$ and Γ represents the gamma function. If α is an integer, these derivatives are defined in the usual sense, i.e.:

$${}_a D_x^\alpha f(x) = \left(\frac{d}{dx} \right)^n f(x) \quad (3)$$

$${}_a D_t^\alpha f(x) = \left(\frac{d}{dx} \right)^n f(t) \quad (4)$$

$\alpha=1,2,\dots$

3 The Hamiltonian formulation of higher derivative field theories with Riemann–Liouville fractional derivative

In this section, we will reformulate the generalized Euler–Lagrange equation, the generalized energy–momentum tensor, and the Hamiltonian density of higher derivative theories using fractional derivatives [22].

Let us start with a Lagrangian

$$\mathcal{L}(\phi, {}_a D_{x_\mu}^k \phi, {}_a D_{x_\mu}^k {}_a D_{x_\nu}^k \phi, \dots, {}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k \dots {}_a D_{x_{\mu_N}}^k \phi) \quad (5)$$

Which for simplicity we consider to be a function of a scalar field and to depend on a finite number of N derivatives. In this case, the action function corresponding to the above Lagrangian is given by

$$S = \int d^4x \mathcal{L}(\phi, {}_a D_{x_\mu}^k \phi, {}_a D_{x_\mu}^k {}_a D_{x_\nu}^k \phi, \dots, {}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k \dots {}_a D_{x_{\mu_N}}^k \phi) \quad (6)$$

The generalized Euler–Lagrange equation in fractional form is obtained by following the typical approach of extremizing the action.

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial \phi} - {}_a D_{x_\mu}^k \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_\mu}^k \phi)} \right) + {}_a D_{x_\mu}^k {}_a D_{x_\nu}^k \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_\mu}^k {}_a D_{x_\nu}^k \phi)} \right) - \dots \\ + (-1)^n {}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k \dots {}_a D_{x_{\mu_N}}^k \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k \dots {}_a D_{x_{\mu_N}}^k \phi)} \right) = 0 \quad (7) \end{aligned}$$

and the generalized energy–momentum tensor in the same way

$$\begin{aligned}
 T_v^\mu = & -\delta_v^\mu \mathcal{L} + \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_\mu}^k \phi)} \right) {}_a D_{x_v}^k \phi - \\
 & [{}_a D_{x_{\mu_1}}^k \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_\mu}^k {}_a D_{x_{\mu_1}}^k \phi)} \right) {}_a D_{x_v}^k \phi - \\
 & \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_\mu}^k {}_a D_{x_{\mu_1}}^k \phi)} \right) {}_a D_{x_{\mu_1}}^k {}_a D_{x_v}^k \phi] \\
 & + [{}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k {}_a D_{x_\mu}^k \phi)} \right) {}_a D_{x_v}^k \phi - \\
 & {}_a D_{x_{\mu_1}}^k \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k {}_a D_{x_\mu}^k \phi)} \right) {}_a D_{x_{\mu_2}}^k {}_a D_{x_v}^k \phi + \\
 & \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k {}_a D_{x_\mu}^k \phi)} \right) {}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k {}_a D_{x_v}^k \phi] + \\
 & \dots (-1)^{(N-1)} [{}_a D_{x_{\mu_1}}^k \dots {}_a D_{x_{\mu_{N-1}}}^k \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_{\mu_1}}^k \dots {}_a D_{x_{\mu_{N-1}}}^k {}_a D_{x_\mu}^k \phi)} \right) \\
 & {}_a D_{x_v}^k \phi - {}_a D_{x_{\mu_1}}^k \dots {}_a D_{x_{\mu_{N-2}}}^k \\
 & \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_{\mu_1}}^k \dots {}_a D_{x_{\mu_{N-1}}}^k {}_a D_{x_\mu}^k \phi)} \right) {}_a D_{x_{\mu_{N-1}}}^k {}_a D_{x_v}^k \phi + \\
 & \dots (-1)^{(N-1)} \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_{\mu_1}}^k \dots {}_a D_{x_{\mu_{N-1}}}^k {}_a D_{x_\mu}^k \phi)} \right) \\
 & {}_a D_{x_{\mu_1}}^k \dots {}_a D_{x_{\mu_{N-1}}}^k {}_a D_{x_v}^k \phi] \quad (8)
 \end{aligned}$$

The Hamiltonian density in fractional form, which corresponds to the component T_0^0 , is now given by

$$\begin{aligned} \mathcal{H} = & -\mathcal{L} + \frac{\partial \mathcal{L}}{\partial({}_a D_{x_0}^k \phi)} {}_a D_{x_0}^k \phi - \\ & [{}_a D_{x_{\mu_1}}^k \left(\frac{\partial \mathcal{L}}{\partial({}_a D_{x_0}^k {}_a D_{x_{\mu_1}}^k \phi)} \right) {}_a D_{x_0}^k \phi - \\ & \left(\frac{\partial \mathcal{L}}{\partial({}_a D_{x_0}^k {}_a D_{x_{\mu_1}}^k \phi)} \right) {}_a D_{x_{\mu_1}}^k {}_a D_{x_0}^k \phi] + \\ & [{}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k \left(\frac{\partial \mathcal{L}}{\partial({}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k {}_a D_{x_0}^k \phi)} \right) {}_a D_{x_0}^k \phi - \\ & {}_a D_{x_{\mu_1}}^k \left(\frac{\partial \mathcal{L}}{\partial({}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k {}_a D_{x_0}^k \phi)} \right) {}_a D_{x_{\mu_2}}^k {}_a D_{x_0}^k \phi + \\ & \left(\frac{\partial \mathcal{L}}{\partial({}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k {}_a D_{x_0}^k \phi)} \right) {}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k {}_a D_{x_0}^k \phi] + \\ & \dots (-1)^{N-1} [{}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k \dots {}_a D_{x_{\mu_{N-1}}}^k \\ & \left(\frac{\partial \mathcal{L}}{\partial({}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_2}}^k \dots {}_a D_{x_{\mu_{N-1}}}^k {}_a D_{x_0}^k \phi)} \right) {}_a D_{x_0}^k \phi - \\ & {}_a D_{x_{\mu_1}}^k \dots {}_a D_{x_{\mu_{N-2}}}^k \left(\frac{\partial \mathcal{L}}{\partial({}_a D_{x_{\mu_1}}^k {}_a D_{x_{\mu_{N-1}}}^k {}_a D_{x_0}^k \phi)} \right) \\ & {}_a D_{x_{\mu_{N-1}}}^k {}_a D_{x_0}^k \phi + (-1)^{N-1} \left(\frac{\partial \mathcal{L}}{\partial({}_a D_{x_{\mu_1}}^k \dots {}_a D_{x_{\mu_{N-1}}}^k {}_a D_{x_0}^k \phi)} \right) \\ & {}_a D_{x_{\mu_1}}^k \dots {}_a D_{x_{\mu_{N-1}}}^k {}_a D_{x_0}^k \phi] \quad (9) \end{aligned}$$

4 The fractional form of the Euler–Lagrange equation of the Extended Maxwell–Chern–Simons Lagrangian density

As an example of a Lagrangian with a higher derivative, let us consider the Lagrangian for the extended Maxwell–Chern–Simons model in 2+1 dimensions in the Lorentz gauge, which was proposed by Reyes [8]

$$\mathcal{L} = \frac{-1}{4} F_{\mu\nu} F^{\mu\nu} + \frac{g}{2} \varepsilon^{\alpha\beta\gamma} (\square A_\alpha) {}_a D_{x_\beta}^k A_\gamma - \frac{1}{2} ({}_a D_{x_\mu}^k A^\mu)^2 \quad (10)$$

where A_μ is the four-vector potential, $F_{\mu\nu} = {}_a D_{x_\mu}^k A^\nu - {}_a D_{x_\nu}^k A^\mu$ is a four dimension antisymmetric second rank tensor, $\eta_{\mu\nu} = \text{diag}(1, -1, 1)$ spacetime metric raises and lowers the indices, $\varepsilon^{\alpha\beta\gamma}$ the Levi–Civita tensor, fully antisymmetric, is given by $\varepsilon^{012} = 1$, and the constant $g > 0$ being a coupling coefficient of the Chern–Simons term.

To reformulate the extended Maxwell–Chern–Simons Lagrangian density in fractional form, we first define the relations

$${}_a D_{x_\mu}^k = ({}_a D_t^k, {}_a D_{x_i}^k), {}_a D_{x^\mu}^k = ({}_a D_t^k, -{}_a D_{x^i}^k) \quad (11)$$

$$A^\sigma = (\phi, \mathbf{A}), A_\sigma = (\phi, -\mathbf{A}) \quad (12)$$

In fractional form, the Euler–Lagrange equation for such a Lagrangian density is given by

$$\begin{aligned} \frac{\partial \mathcal{L}}{\partial A_\sigma} - {}_a D_{x_\lambda}^k \left(\frac{\partial \mathcal{L}}{\partial({}_a D_{x_\lambda}^k A_\sigma)} \right) + \\ {}_a D_{x_\mu}^k {}_a D_{x_\lambda}^k \left(\frac{\partial \mathcal{L}}{\partial({}_a D_{x_\mu}^k {}_a D_{x_\lambda}^k A_\sigma)} \right) = 0 \quad (13) \end{aligned}$$

The first term $\frac{\partial \mathcal{L}}{\partial A_\sigma} = 0$. The second term in Eq. (13) may be evaluated as follows:

$$\left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_\lambda}^k A_\sigma)} \right) = -F^{\lambda\sigma} + \frac{g}{2} \varepsilon^{\alpha\beta\gamma} (\square A_\alpha) \delta_\beta^\lambda \delta_\gamma^\sigma - \frac{1}{2} \left(\frac{\partial}{\partial ({}_a D_{x_\lambda}^k A_\sigma)} \right) [\eta^{\mu\mu'} {}_a D_{x_\mu}^k A_{\mu'} \eta^{\nu\nu'} {}_a D_{x_\nu}^k A_{\nu'}] \quad (14)$$

$$= -F^{\lambda\sigma} + \frac{g}{2} \varepsilon^{\alpha\lambda\sigma} (\square A_\alpha) - \frac{1}{2} \eta^{\mu\mu'} \eta^{\nu\nu'} [\delta_\mu^\lambda \delta_{\mu'a}^\sigma D_{x_\nu}^k A_{\nu'} + \delta_\nu^\lambda \delta_{\nu'a}^\sigma D_{x_\mu}^k A_{\mu'}] \quad (15)$$

which can be rewritten as

$$= -F^{\lambda\sigma} + \frac{g}{2} \varepsilon^{\alpha\lambda\sigma} (\square A_\alpha) - \frac{1}{2} [\eta^{\lambda\sigma} {}_a D_{x_\nu}^k A^\nu + \eta^{\lambda\sigma} {}_a D_{x_\mu}^k A^\mu] \quad (16)$$

$$= -F^{\lambda\sigma} + \frac{g}{2} \varepsilon^{\alpha\lambda\sigma} (\square A_\alpha) - \eta^{\lambda\sigma} {}_a D_{x_\mu}^k A^\mu \quad (17)$$

It follows that

$${}_a D_{x_\lambda}^k \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_\lambda}^k A_\sigma)} \right) = -{}_a D_{x_\lambda}^k F^{\lambda\sigma} + \frac{g}{2} \varepsilon^{\alpha\lambda\sigma} {}_a D_{x_\lambda}^k (\square A_\alpha) + \eta^{\lambda\sigma} {}_a D_{x_\lambda}^k {}_a D_{x_\mu}^k A^\mu \quad (18)$$

Similarly, the third term in Eq. (13) may be obtained as follows:

$$\left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_\mu}^k {}_a D_{x_\lambda}^k A_\sigma)} \right) = \left(\frac{\partial}{\partial ({}_a D_{x_\mu}^k {}_a D_{x_\lambda}^k A_\sigma)} \right) \left(\frac{g}{2} \varepsilon^{\alpha\beta\gamma} {}_a D_{x_\zeta}^k {}_a D_{x_\zeta}^k A_\alpha \right) {}_a D_{x_\beta}^k A_\gamma \quad (19)$$

which can be simplified to

$$= \frac{g}{2} \varepsilon^{\alpha\beta\gamma} \eta^{\zeta\nu} \delta_\mu^\zeta \delta_\lambda^\zeta \delta_\sigma^\alpha {}_a D_{x_\beta}^k A_\gamma \quad (20)$$

$$= \frac{g}{2} \varepsilon^{\sigma\beta\gamma} \eta^{\mu\lambda} {}_a D_{x_\beta}^k A_\gamma \quad (21)$$

then we have

$${}_a D_{x_\mu}^k {}_a D_{x_\lambda}^k \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_{x_\mu}^k {}_a D_{x_\lambda}^k A_\sigma)} \right) = \frac{g}{2} \varepsilon^{\sigma\beta\gamma} \eta^{\mu\lambda} {}_a D_{x_\mu}^k {}_a D_{x_\lambda}^k {}_a D_{x_\beta}^k A_\gamma \quad (22)$$

Substituting Eqs. (18 and 22) into Eq. (13), we get

$${}_a D_{x_\lambda}^k F^{\lambda\sigma} - \frac{g}{2} \varepsilon^{\alpha\lambda\sigma} {}_a D_{x_\lambda}^k (\square A_\alpha) + \eta^{\lambda\sigma} {}_a D_{x_\lambda}^k {}_a D_{x_\mu}^k A^\mu + \frac{g}{2} \varepsilon^{\sigma\beta\gamma} \eta^{\mu\lambda} {}_a D_{x_\mu}^k {}_a D_{x_\lambda}^k {}_a D_{x_\beta}^k A_\gamma = 0 \quad (23)$$

$$\begin{aligned}
 {}_a D_{x_\lambda}^k F^{\lambda\sigma} - \frac{g}{2} \varepsilon^{\alpha\lambda\sigma} (\square A_\alpha) {}_a D_{x_\lambda}^k A_\alpha + \\
 \frac{g}{2} \varepsilon^{\sigma\beta\gamma} {}_a D_{x_\lambda}^k {}_a D_{x_\lambda}^k {}_a D_{x_\beta}^k A_\gamma \\
 + {}_a D_{x_\sigma}^k {}_a D_{x_\mu}^k A^\mu = 0
 \end{aligned} \quad (24)$$

$$\begin{aligned}
 {}_a D_{x_\lambda}^k F^{\lambda\sigma} - \frac{g}{2} \square \left(\varepsilon^{\alpha\lambda\sigma} {}_a D_{x_\lambda}^k A_\alpha - \varepsilon^{\sigma\beta\gamma} {}_a D_{x_\beta}^k A_\gamma \right) + \\
 {}_a D_{x_\sigma}^k {}_a D_{x_\mu}^k A^\mu = 0
 \end{aligned} \quad (25)$$

Setting $\lambda = \gamma$ and $\alpha = \beta$ in the term $\varepsilon^{\alpha\lambda\sigma} {}_a D_{x_\lambda}^k A_\alpha$ in Eq. (25). This lead to following:

$$\begin{aligned}
 {}_a D_{x_\lambda}^k F^{\lambda\sigma} - \frac{g}{2} \square \left(\varepsilon^{\beta\gamma\sigma} {}_a D_{x_\gamma}^k A_\beta - \varepsilon^{\sigma\beta\gamma} {}_a D_{x_\beta}^k A_\gamma \right) + \\
 {}_a D_{x_\sigma}^k {}_a D_{x_\mu}^k A^\mu = 0
 \end{aligned} \quad (26)$$

$$\begin{aligned}
 {}_a D_{x_\lambda}^k F^{\lambda\sigma} - \frac{g}{2} \square \left(\varepsilon^{\beta\gamma\sigma} {}_a D_{x_\gamma}^k A_\beta - \varepsilon^{\sigma\beta\gamma} {}_a D_{x_\beta}^k A_\gamma \right) + \\
 {}_a D_{x_\sigma}^k {}_a D_{x_\mu}^k A^\mu = 0
 \end{aligned} \quad (27)$$

using the definition of the field strength tensor $F_{\beta\gamma} = {}_a D_{x_\beta}^k A_\gamma - {}_a D_{x_\gamma}^k A_\beta$, Eq. (27) can be written as:

$${}_a D_{x_\lambda}^k F^{\lambda\sigma} + \frac{g}{2} \varepsilon^{\sigma\beta\gamma} \square F_{\beta\gamma} + {}_a D_{x_\sigma}^k {}_a D_{x_\mu}^k A^\mu = 0 \quad (28)$$

The above equation represents the fractional form of the modified Maxwell equations. It's worth pointing out that for $k \rightarrow 1$, Eq. (28) simplifies to the usual Maxwell equation:

$$\partial_\lambda F^{\lambda\sigma} + \frac{g}{2} \varepsilon^{\sigma\beta\gamma} \square F_{\beta\gamma} + \partial^\sigma (\partial \cdot A) = 0 \quad (29)$$

Equation (28) can be rewritten as follows, see appendix A.

$$\left(\eta^{\sigma\alpha} + g \varepsilon^{\sigma\beta\alpha} \right) (\square A_\alpha) = 0 \quad (30)$$

5 The Hamiltonian formulation of the Extended Maxwell–Chern–Simons Model with the Riemann–Liouville fractional derivative

Consider $A_\mu(x)$ and ${}_a D_t^k A_\mu$ as two independent configuration field variables with their corresponding conjugate momenta defined by

$$P^\mu = \frac{\partial \mathcal{L}}{\partial ({}_a D_t^k A_\mu)} - {}_a D_t^k \Pi^\mu, \quad \Pi^\mu = \frac{\partial \mathcal{L}}{\partial ({}_a D_t^k {}_a D_t^k A_\mu)} \quad (31)$$

By using Eq. (17), we can write the conjugate momenta Π^μ as

$$\Pi^\mu = \frac{\partial \mathcal{L}}{\partial ({}_a D_t^k {}_a D_t^k A_\mu)} = \frac{g}{2} \varepsilon^{\mu\beta\gamma} {}_a D_{x_\beta}^k A_\gamma \quad (32)$$

which reads in components

$$\Pi^0 = \frac{\partial \mathcal{L}}{\partial ({}_a D_t^k {}_a D_t^k A_\mu)} = \frac{g}{2} \varepsilon^{0ij} {}_a D_{x_i}^k A_j \quad (33)$$

$$\Pi^i = \frac{g}{2} \varepsilon^{ij} \left({}_a D_t^k A_j - {}_a D_j^k A_0 \right) \quad (34)$$

and the conjugate momenta P^μ is given by

$$P^\mu = \frac{\partial \mathcal{L}}{\partial ({}_a D_t^k A_\mu)} - {}_a D_t^k \left(\frac{\partial \mathcal{L}}{\partial ({}_a D_t^k A_\mu)} \right) \tag{35}$$

By making use of Eq. (17), we obtain

$$\frac{\partial \mathcal{L}}{\partial ({}_a D_t^k A_\mu)} == -F^{0\mu} + \frac{g}{2} \varepsilon^{\alpha 0\mu} (\square A_\alpha) - \eta^{0\mu} ({}_a D_{x_\gamma}^k A^\gamma) \tag{36}$$

or

$$\frac{\partial \mathcal{L}}{\partial ({}_a D_t^k A_\mu)} == -F^{0\mu} - \frac{g}{2} \varepsilon^{\mu 0\alpha} (\square A_\alpha) - \eta^{0\mu} ({}_a D_{x_\gamma}^k A^\gamma) \tag{37}$$

Setting $\alpha = \gamma$ in the second term of Eq. (37), we get

$$\frac{\partial \mathcal{L}}{\partial ({}_a D_t^k A_\mu)} == -F^{0\mu} - \frac{g}{2} \varepsilon^{\mu 0\gamma} (\square A_\gamma) - \eta^{0\mu} ({}_a D_{x_\gamma}^k A^\gamma) \tag{38}$$

Substituting Eqs. (38 and 32) into Eq. (35), we get

$$P^\mu = -F^{0\mu} - \frac{g}{2} \varepsilon^{\mu 0\gamma} (\square A_\gamma) - \eta^{0\mu} ({}_a D_{x_\gamma}^k A^\gamma) - \frac{g}{2} \varepsilon^{\mu\beta\gamma} {}_a D_{x_\beta}^k {}_a D_{x_\gamma}^k A^\gamma \tag{39}$$

which read in components

$$P^0 = -F^{00} - \frac{g}{2} \varepsilon^{00\gamma} (\square A_\gamma) - \eta^{00} ({}_a D_{x_\gamma}^k A^\gamma) - \frac{g}{2} \varepsilon^{0\beta\gamma} {}_a D_{x_\beta}^k {}_a D_{x_\gamma}^k A^\gamma \tag{40}$$

$$P^0 = - ({}_a D_{x_\gamma}^k A^\gamma) - \frac{g}{2} \varepsilon^{ij} {}_a D_{x_i}^k {}_a D_{x_j}^k A_j \tag{41}$$

$$P^i = -F^{0i} - \frac{g}{2} \varepsilon^{i0\gamma} (\square A_\gamma) - \eta^{0i} ({}_a D_{x_\gamma}^k A^\gamma) - \frac{g}{2} \varepsilon^{i\beta\gamma} {}_a D_{x_\beta}^k {}_a D_{x_\gamma}^k A^\gamma \tag{42}$$

$$P^i = F_{0i} + \frac{g}{2} \varepsilon^{ij} (\square A_j) - \frac{g}{2} \varepsilon^{i\beta\gamma} {}_a D_{x_\beta}^k {}_a D_{x_\gamma}^k A^\gamma \tag{43}$$

Eq. (43) can be rewritten as

$$P^i = F_{0i} + \frac{g}{2} \varepsilon^{ij} (\square A_j) - \frac{g}{2} \varepsilon^{i0\gamma} {}_a D_{x_\beta}^k {}_a D_{x_\gamma}^k A^\gamma - \frac{g}{2} \varepsilon^{ij\gamma} {}_a D_{x_j}^k {}_a D_{x_\gamma}^k A^\gamma \tag{44}$$

$$P^i = F_{0i} + \frac{g}{2} \varepsilon^{ij} (\square A_j) + \frac{g}{2} \varepsilon^{0ij} {}_a D_{x_\beta}^k {}_a D_{x_\gamma}^k A^\gamma - \frac{g}{2} \varepsilon^{ij0} {}_a D_{x_j}^k {}_a D_{x_\gamma}^k A^\gamma \tag{45}$$

or

$$P^i = F_{0i} + \frac{g}{2} \varepsilon^{ij} (\square A_j) + \frac{g}{2} \varepsilon^{ij} {}_a D_t^k F_{jo} \quad (46)$$

The Hamiltonian density is given by

$$\mathcal{H} = P^\mu {}_a D_t^k A_\mu + \Pi^\mu {}_a D_t^k {}_a D_t^k A_\mu - \mathcal{L} \quad (47)$$

Taking into account Eqs. (10 and 32), then the Hamiltonian density in fractional form is

$$\begin{aligned} \mathcal{H} = P^0 {}_a D_t^k A_0 + P^i {}_a D_t^k A_i + \\ \frac{g}{2} \varepsilon^{\mu\beta\gamma} {}_a D_{x_\beta}^k A_\gamma {}_a D_t^k {}_a D_t^k A_\mu - \frac{1}{2} (F_{0j})^2 + \frac{1}{4} (F_{ij})^2 - \\ \frac{g}{2} \varepsilon^{\alpha\beta\gamma} (\square A_\alpha) {}_a D_{x_\beta}^k A_\gamma + \frac{1}{2} \left({}_a D_{x_\mu}^k A^\mu \right)^2 \end{aligned} \quad (48)$$

By using the relation $\square A = {}_a D_t^k {}_a D_t^k - \nabla^2$ Eq. (48), simplified to

$$\begin{aligned} \mathcal{H} = P^0 {}_a D_t^k A_0 + P^i {}_a D_t^k A_i - \frac{1}{2} (F_{0j})^2 + \frac{1}{4} (F_{ij})^2 + \\ \frac{g}{2} \varepsilon^{\alpha\beta\gamma} (\nabla^2 A_\alpha) {}_a D_{x_\beta}^k A_\gamma + \frac{1}{2} \left({}_a D_{x_\mu}^k A^\mu \right)^2 \end{aligned} \quad (49)$$

The second term in Eq. (49) can be written as

$$\Pi^i = \frac{g}{2} \varepsilon^{i\beta\gamma} {}_a D_{x_\beta}^k A_\gamma = \frac{g}{2} \varepsilon^{ij} \left({}_a D_{x_j}^k A_0 - {}_a D_t^k A_j \right) \quad (50)$$

or

$${}_a D_t^k A_j = -\frac{2}{g} \varepsilon^{ij} \Pi^i + {}_a D_{x_j}^k A_0 \quad (51)$$

$${}_a D_t^k A_i = \frac{2}{g} \varepsilon^{ij} \Pi^j + {}_a D_{x_i}^k A_0 \quad (52)$$

Multiply both sides of Eq. (52) by P^i , we get

$$P^i {}_a D_t^k A_i = \frac{2}{g} \varepsilon^{ij} P^i \Pi^j + P^i {}_a D_{x_i}^k A_0 \quad (53)$$

or

$$P^i {}_a D_t^k A_i = \frac{2}{g} \varepsilon^{ij} P_i \Pi_j - P_i {}_a D_{x_i}^k A_0 \quad (54)$$

The third term in Eq. (49) can be expressed as

$$\Pi^i = \frac{g}{2} \varepsilon^{ij} \left({}_a D_t^k A_j - {}_a D_{x_j}^k A_0 \right) = \frac{g}{2} \varepsilon^{ij} F_{0j} \quad (55)$$

It follows that

$$(F_{0i})^2 = 4 (\Pi_i)^2 \setminus (g^2) \quad (56)$$

The fourth term in Eq. (49) can be written in the following way:

$$\begin{aligned} \frac{g}{2} \varepsilon^{\alpha\beta\gamma} (\nabla^2 A_\alpha) {}_a D_{x_\beta}^k A_\gamma = \frac{g}{2} \varepsilon^{0\beta\gamma} (\nabla^2 A_0) {}_a D_{x_\beta}^k A_\gamma + \\ \frac{g}{2} \varepsilon^{i\beta\gamma} (\nabla^2 A_i) {}_a D_{x_\beta}^k A_\gamma \end{aligned} \quad (57)$$

$$\begin{aligned} = \frac{g}{2} \varepsilon^{ij} (\nabla^2 A_0) {}_a D_{x_i}^k A_j + \\ \frac{g}{2} \left(\varepsilon^{i0\gamma} {}_a D_{x_0}^k A_\gamma + \varepsilon^{i\beta 0} {}_a D_{x_\beta}^k A_0 \right) (\nabla^2 A_i) \end{aligned} \quad (58)$$

$$= \frac{g}{2} \varepsilon^{ij} (\nabla^2 A_0) {}_a D_{x_i}^k A_j + \frac{g}{2} \varepsilon^{ij} \left({}_a D_{x_j}^k A_0 - {}_a D_{x_0}^k A_j \right) \tag{59}$$

$$= \frac{g}{2} \varepsilon^{ij} (\nabla^2 A_0) {}_a D_{x_i}^k A_j - \frac{g}{2} \varepsilon^{ij} (F_{0j}) (\nabla^2 A_i) \tag{60}$$

Inserting Eq. (55) into Eq. (60), we get

$$\frac{g}{2} \varepsilon^{\alpha\beta\gamma} (\nabla^2 A_\alpha) {}_a D_{x_\beta}^k A_\gamma = \frac{g}{2} \varepsilon^{ij} (\nabla^2 A_0) {}_a D_{x_i}^k A_j - \Pi_i (\nabla^2 A_i) \tag{61}$$

Substituting Eqs. (54, 56, and 61) into Eq. (49), the Hamiltonian density can be written as

$$\begin{aligned} \mathcal{H} = P^0 {}_a D_t^k A_0 + \frac{2}{g} \varepsilon^{ij} P_i \Pi_j - P_i {}_a D_{x_i}^k A_j - \frac{2(\Pi_i)^2}{(g^2)} + \\ \frac{1}{4} (F_{ij})^2 + \frac{g}{2} \varepsilon^{ij} (\nabla^2 A_0) {}_a D_{x_i}^k A_j - \Pi_i (\nabla^2 A_i) + \\ \frac{1}{2} \left({}_a D_{x_\mu}^k A^\mu \right)^2 \end{aligned} \tag{62}$$

6 Conclusion

In this work, the Hamiltonian formulation for higher derivatives has been reformulated using fractional derivatives. For higher order derivatives, the extended Maxwell–Chern–Simons Lagrangian density is reformulated in fractional form using the Riemann–Liouville fractional derivative. The fractional form of the Euler–Lagrange equation and the Hamiltonian density of these systems are obtained. The classical results are obtained when fractional derivatives are replaced by integer order derivatives.

Appendix A: The Derivation of Equation (30):

We can write Eq. (28) as

$${}_a D_{x_\lambda}^k \left({}_a D_{x_\lambda}^k A^\sigma - {}_a D_{x_\sigma}^k A^\lambda \right) + {}_a D_{x^\sigma}^k {}_a D_{x_\mu}^k A^\mu + \frac{g}{2} \varepsilon^{\sigma\beta\gamma} \square \left({}_a D_{x_\beta}^k A_\gamma - {}_a D_{x_\gamma}^k A_\beta \right) = 0 \tag{A.1}$$

The above equation can be rearranged as:

$$\square A^\sigma - {}_a D_{x^\sigma}^k {}_a D_{x_\lambda}^k A^\lambda + {}_a D_{x^\sigma}^k {}_a D_{x_\mu}^k A^\mu + \frac{g}{2} \varepsilon^{\sigma\beta\gamma} \square \left({}_a D_{x_\beta}^k A_\gamma - {}_a D_{x_\gamma}^k A_\beta \right) = 0 \tag{A.2}$$

The second and third terms cancel after changing a dummy index from μ to λ . This leaves us with

$$\eta^{\sigma\alpha} \square A_\alpha + \frac{g}{2} \varepsilon^{\sigma\beta\gamma} \square \left({}_a D_{x_\beta}^k A_\gamma - {}_a D_{x_\gamma}^k A_\beta \right) = 0 \tag{A.3}$$

using $\varepsilon^{\sigma\beta\zeta} \varepsilon_{\sigma\beta\alpha} = \delta_\alpha^\zeta$, we rewrite Eq. (A.3) as:

$$\eta^{\sigma\alpha} \square A_\alpha + \frac{g}{2} \varepsilon^{\sigma\beta\gamma} \varepsilon^{\sigma\beta\alpha} \varepsilon_{\sigma\beta\alpha} \square \left({}_a D_{x_\beta}^k A_\gamma - {}_a D_{x_\gamma}^k A_\beta \right) = 0 \tag{A.4}$$

or

$$\eta^{\sigma\alpha} \square A_\alpha + \frac{g}{2} \left(\varepsilon^{\sigma\beta\alpha} {}_a D_{x_\beta}^k A_\alpha - \varepsilon^{\sigma\beta\alpha} {}_a D_{x_\alpha}^k A_\beta \right) = 0 \quad (\text{A.5})$$

let $\alpha \rightarrow \beta$ and $\beta \rightarrow \alpha$ in the term $\varepsilon^{\sigma\beta\alpha} {}_a D_{x_\alpha}^k A_\beta$ of the above equation, we get

$$\eta^{\sigma\alpha} \square A_\alpha + \frac{g}{2} \square \left(\varepsilon^{\sigma\beta\alpha} {}_a D_{x_\beta}^k A_\alpha + \varepsilon^{\sigma\beta\alpha} {}_a D_{x_\beta}^k A_\alpha \right) = 0 \quad (\text{A.6})$$

It follows that

$$\left(\eta^{\sigma\alpha} + g \varepsilon^{\sigma\beta\alpha} {}_a D_{x_\beta}^k \right) \square A_\alpha = 0 \quad (\text{A.7})$$

The author is grateful for the suggestions of anonymous referees, which have considerably improved the readability of the paper.

References

- [1] S. I. Muslih and D. Baleanu, Formulation of Hamiltonian Equations for Fractional Variational Problems, *Czechoslovak Journal of Physics*, **55**, 633-642 (2005).
- [2] E. K. Jaradat, Canonical Quantization for Fractional Schrödinger Lagrangian Density in Caputo Definition, *Jordan Journal of Physics*, **6**, 55-63 (2013).
- [3] Z. Yi, Fractional Differential Equations of Motion in Terms of Combined Riemann–Liouville derivatives, *Chines Physics B*, **21**, 084502- 084507 (2012).
- [4] A. D. Al-oqali, B. M. Al-khamiseh, E. K. Jaradat, and R. S. Hijjawi, The Linear Sigma Model Lagrangian Density: Fractional Formulation, *Canadian Journal of Pure and Applied Sciences*, **10**, 3803–3807 (2016).
- [5] T. Kang-Bo, L. Chang-Hong and D. Xiao-Jie, Electrodynamic Analysis of Dissipative Electromagnetic Materials Based on Fractional Derivative, *Chinese Physics Letters*, **24**, 847-850 (2007).
- [6] J. C. Gutiérrez-Vega, Fractionalization of Optical Beams: I. Planar Analysis, *Optics letters*, **32**, 1521–1523 (2007).
- [7] R. A. Magomedov, R. R. Meilanov, R. P. Meilanov, E. N. Akhmedov, V. D. Beybalaev and A. A. Aliverdiev, Generalization of Thermodynamics in of Fractional-Order Derivatives and Calculation of Heat-Transfer Properties of Noble Gases, *Journal of Thermal Analysis and Calorimetry*, **133**, 1189–1194 (2018).
- [8] C. M. Reyes, Testing symmetries in effective models of higher derivative field theories, *Physical Review D*, **80**, 105008-105020 (2009).
- [9] W. Thirring, Regularization as a Consequence of Higher Order Equations, *Physical Review*, **77**, 570-570 (1950)
- [10] D. A. Eliezer and R. P. Woodard, The Problem of Nonlocality in String Theory, *Nuclear Physics B*, **325**, 389-469 (1989).
- [11] C. S. Chu, J. Lukierski and W. J. Zakrzewski, Hermitian Analyticity, IR/UV Mixing and Unitarity of Noncommutative Field Theories, *Nuclear Physics B*, **632**, 219-239 (2002).
- [12] J. A. Wheeler and R. P. Feynman, Classical Electrodynamics in Terms of Direct Interparticle Action, *Reviews of Modern physics*, **21**, 425-432 (1949).
- [13] S. M. Carroll and M. Hoffman, Can the Dark Energy Equation-of-State Parameter w be Less Than -1?, *Physical Review D*, **68**, 0203509-0203519 (2003).
- [14] J. Karouby, T. Qiu, and R. Brandenberger, Instability of the Lee-Wick Bounce, *Physical Review D*, **84**, 043505- 043521 (2011).
- [15] G. Kaniadakis and A. M. Scarfone, A Maxwell–Chern–Simons Model for a Quantum System Obeying an Exclusion–Inclusion Principle, *Physica A*, **291**, 423-438 (2001).
- [16] J. Niederle, A. G. Nikitin and O. Kuriksha, Maxwell–Chern–Simons Models: Their Symmetries, Exact Solutions and Non-relativistic Limits, *Acta Polytechnica*, **50**, 96-101 (2010).
- [17] K. Haller and E. Lim-Lombridas, Maxwell–Chern–Simons Theory in Covariant and Coulomb Gauges, *Annals of Physics*, **246**, 1-48 (1996).
- [18] O. Jarab’ah and K. Nawafleh, Fractional Hamiltonian of Nonconservative Systems with Second Order Lagrangian, *American Journal of Physics and Applications*, **6**, 85-88 (2018).
- [19] Y. M. Alawaideh, R. S. Hijjawi and J. M. Khalifeh, Hamiltonian Formulation for Continuous Third Order Systems Using Fractionals Derivatives, *Jordan Journal of Physics*, **14**, 35-47 (2021).
- [20] A. D. Al-Oqali, Fractional Formulation of Podolsky Lagrangian Density, *International Journal of Advanced and Applied Sciences*, **9**, 136-141 (2022).
- [21] A. A. Diab, R. S. Hijjawi, J. H. Asad and J. M. Khalifeh, Hamiltonian Formulation of Classical Fields with Fractional Derivatives: Revisited, *Meccanica*, **48**, 323–330 (2013).
- [22] N. Moeller and B. Zwiebach, Dynamics with infinitely Many Time derivatives and Rolling Tachyons, *Journal of High Energy Physics*, **10**, 1-40 (2002).