

Solvability of Nonlinear Quadratic Functional Equations

H. H. G. Hashem* and A. M. A. El-Sayed

Faculty of Science, Alexandria University, Alexandria, Egypt

Received: 15 Feb. 2015, Revised: 17 May 2015, Accepted: 18 May 2015

Published online: 1 Sep. 2015

Abstract: We give an existence theorem for some quadratic functional equations which includes many key integral and functional equations that arise in nonlinear analysis and its applications. In particular, we extend the class of characteristic functions appearing in Chandrasekhar's classical integral equation from astrophysics and retain existence of its solutions. Also, some counter examples are considered.

Keywords: Quadratic functional integral equation; Positive monotonic solutions; Measure of noncompactness.

1 Introduction

The study of integral equations have gained much attention due to extensive applications of these equations in describing numerous events and problems of real world, and the theory of integral equations is rapidly developing with the help of several tools of functional analysis, topology and fixed point theory. For details, we refer to [2]-[11] and [17]-[27].

One of the kinds of integral equations is quadratic integral equations which have received increasing attention during recent years due to its applications in numerous diverse fields of science and engineering for example, the theory of radiative transfer, kinetic theory of gases, the theory of neutron transport and the traffic theory. Many authors have studied different kinds of nonlinear quadratic integral equations in different classes (see [2], [3], [7]-[14] and [17] - [27]). Especially, Chandrasekar's integral equation which has been a subject of much investigation since its appearance around fifty years ago [13].

Let $L_1 = L_1[0, T]$ be the class of Lebesgue integrable functions on $I = [0, T]$ with the standard norm.

Here, we are concerning with the nonlinear quadratic functional equation

$$x(t) = f(t, x(\phi_1(t))) + g(t, x(\phi_2(t))) \cdot \psi\left(t, \int_0^{\alpha(t)} u(t, s, x(\phi_3(s))) ds\right), \quad t \in I. \quad (1)$$

which includes as special cases numerous function, integral and functional integral equations encountered in

nonlinear analysis. For example, the quadratic integral equation of Chandrasekhar type

$$x(t) = 1 + \lambda x(t) \int_0^t \frac{t \phi(s)}{t+s} x(s) ds$$

and the quadratic integral equation of fractional order

$$x(t) = f(t) + g(t, x(t)) \int_0^t \frac{(t-s)^{\beta-1}}{\Gamma(\beta)} h(s, x(s)) ds, \quad \beta > 0.$$

2 Preliminaries

In this section, we introduce some notations and preliminary facts which are used in the paper.

Now let E be a Banach space with zero element θ and let X be a nonempty bounded subset of E . Moreover denote by $B_r = B(\theta, r)$ the closed ball in E centered at θ and with radius r . In the sequel we shall need some criteria for compactness in measure; the complete description of compactness in measure was given by Fréchet [4], but the following sufficient condition will be more convenient for our purposes (see [4]).

Theorem 1. Let X be a bounded subset of L_1 . Assume that there is a family of subsets $(\Omega_c)_{0 \leq c \leq b-a}$ of the interval (a, b) such that $\text{meas } \Omega_c = c$ for every $c \in [0, b-a]$, and for every $x \in X$, $x(t_1) \leq x(t_2)$, $(t_1 \in \Omega_c, t_2 \notin \Omega_c)$, then the set X is compact in measure.

* Corresponding author e-mail: hendhghashem@yahoo.com

The measure of weak noncompactness defined by De Blasi [1] and [15] is given by,

$$\beta(X) = \inf\{r > 0 : \text{there exists a weakly compact subset } Y \text{ of } E \text{ such that } X \subset Y + K_r\}$$

The function $\beta(X)$ possesses several useful properties which may be found in [15].

The convenient formula for the function $\beta(X)$ in L_1 was given by Appel and De Pascale (see [1])

$$\beta(X) = \lim_{\varepsilon \rightarrow 0} (\sup_{x \in X} (\sup_D [\int_D |x(t)| dt : D \subset [a, b], \text{meas } D \leq \varepsilon])), \quad (2)$$

where the symbol $\text{meas } D$ stands for Lebesgue measure of the set D .

Next, we shall also use the notion of the Hausdorff measure of noncompactness χ (see [4]) defined by

$$\chi(X) = \inf\{r > 0 : \text{there exists a finite subset } Y \text{ of } E \text{ such that } X \subset Y + K_r\}$$

In the case when the set X is compact in measure, the Hausdorff and De Blasi measures of noncompactness will be identical. Namely we have (see [1] and [15])

Theorem 2. Let X be an arbitrary nonempty bounded subset of L_1 . If X is compact in measure then $\beta(X) = \chi(X)$.

Finally, we will recall the fixed point theorem due to Darbo [6].

Theorem 3. Let Q be a nonempty, bounded, closed and convex subset of E and let $H : Q \rightarrow Q$ be a continuous transformation which is a contraction with respect to the Hausdorff measure of noncompactness χ , i.e., there exists a constant $\alpha \in [0, 1)$ such that $\chi(HX) \leq \alpha \chi(X)$ for any nonempty subset X of Q . Then H has at least one fixed point in the set Q .

3 Existence Theorem

Let the functional operator H be defined as

$$(Hx)(t) = \psi \left(t, \int_0^{\alpha(t)} u(t, s, x(s)) ds \right),$$

$$(Fx)(t) = f(t, x(t))$$

Then equation (1) may be written in operator form as:

$$(Ax)(t) = (Fx(\phi_1))(t) + (Gx(\phi_2))(t) \cdot (Hx(\phi_3))(t)$$

where $(Gx)(t) = g(t, x(t))$.

Consider the following assumptions:

- (i) $f, g : I \times R \rightarrow R$ are functions such that $f, g : I \times R_+ \rightarrow R_+$. Moreover, the functions f, g satisfy Carathéodory condition (i.e. are measurable in t for all $x \in R$ and continuous in x for all $t \in I$) and there exist two functions $m_1, m_2 \in L_1$ and constants $b_1 > 0, b_2 > 0$ such that

$$|f(t, x)| \leq m_1(t) + b_1 |x|, \quad |g(t, x)| \leq m_2(t) + b_2 |x|$$

$$\forall (t, x) \in I \times R.$$

Apart from this the functions f and g are nondecreasing in both variables.

- (ii) $u : I \times I \times R \rightarrow R$ is such that $u(t, s, x) \geq 0$ for $(t, s, x) \in I \times I \times R_+$ and $u(t, s, x)$ satisfies Carathéodory condition (i.e. it is measurable in (t, s) for all $x \in R$ and continuous in x for almost all $(t, s) \in I \times I$).
- (iii) There exist a positive constant b_3 , a function $m_3 \in L_1$ and a measurable (in both variables) function $k(t, s) = k : I \times I \rightarrow R_+$ such that

$$|u(t, s, x)| \leq k(t, s)(m_3(t) + b_3 |x|) \quad \forall t, s \in I \text{ and for } x \in R$$

and the integral operator K , generated by the function k and defined by

$$(Kx)(t) = \int_0^t k(t, s) x(s) ds, \quad t \in I. \quad (3)$$

maps continuously L_1 into L_∞ on I ;

- (iv) $t \rightarrow u(t, s, x)$ is a.e. nondecreasing on I for almost all fixed $s \in I$ and for each $x \in R_+$;
- (v) $\psi : I \times R \rightarrow R$ is a function such that $\psi : I \times R_+ \rightarrow R_+$. Moreover, the function ψ satisfies Carathéodory condition (i.e. is measurable in t for all $x \in R$ and continuous in x for all $t \in I$) and there exist bounded and measurable function $m(t)$ and a constant $b > 0$, such that

$$|\psi(t, x)| \leq m(t) + b |x| \quad \forall (t, x) \in I \times R.$$

Apart from this the function ψ is nondecreasing in both variables.

- (vi) $\alpha : I \rightarrow I$ is continuous.
- (vii) $\phi_i : I \rightarrow I, i = 1, 2, 3$ are increasing, absolutely continuous on and there exist positive constants $B_i, i = 1, 2, 3$ such that $\phi_i' \geq B_i$ a.e. on I ;
- (viii) Let

$$d > \sqrt{4 M b b_2 b_3 B_1^2 B_2 B_3 (||m_1|| + b M \cdot ||m_2|| ||m_3|| + N ||m_2||)},$$

$$|m(t)| \leq N, \quad M = ||K||_{L_\infty}$$

where we assume that

$$d = B_1 B_2 B_3 - b_1 B_2 B_3 - b b_2 M B_1 B_3 ||m_3|| - M b b_3 B_1 B_2 ||m_2|| + b b_2 N B_1 B_3.$$

Moreover, we assume that there exists a positive solution r of the quadratic equation

$$b b_2 b_3 B_1 M r^2 - d r + B_1 B_2 B_3 (||m_1|| + b M \cdot ||m_2|| ||m_3|| + N ||m_2||) = 0.$$

and define the set

$$B_r = \{x \in L_1 : \|x\| \leq r\}.$$

For the existence of at least one L_1 -positive solution of the quadratic functional equation (1) we have the following theorem.

Theorem 4. Let the assumptions (i)-(viii) be satisfied.

If $b_1 B_2 B_3 + M b b_2 B_1 B_3 \|m_3\| + r b b_2 b_3 M + b_2 B_1 B_3 N < B_1 B_2 B_3$, then the quadratic functional equation (1) has at least one solution $x \in L_1$ which is positive and a.e. nondecreasing on I .

Proof. Take an arbitrary $x \in L_1$, then, we get

$$\begin{aligned} |(Ax)(t)| &\leq |m_1(t)| + b_1 |x(\phi_1(t))| \\ &\quad + (m_2(t) + b_2 |x(\phi_2(t))|) \\ &\quad \left[m(t) + b \int_0^{\alpha(t)} k(t,s)(m_3(s) + b_3 |x(\phi_3(s))|) ds \right] \end{aligned}$$

and

$$\begin{aligned} \|(Ax)(t)\| &= \int_0^T |(Ax)(t)| dt \\ &\leq \int_0^T |m_1(t)| dt + b_1 \int_0^T |x(\phi_1(t))| dt \\ &\quad + \int_0^T (m_2(t) + b_2 |x(\phi_2(t))|) \left[m(t) + b \int_0^{\alpha(t)} k(t,s)(m_3(s) + b_3 |x(\phi_3(s))|) ds \right] dt \\ &\leq \|m_1\| + \frac{b_1}{B_1} \int_0^T |x(\phi_1(t))| \phi_1'(t) dt + \int_0^T m_2(t) m(t) dt + b \int_0^T m_2(t) \int_0^{\alpha(t)} k(t,s) m_3(s) ds dt \\ &\quad + b b_2 \int_0^T |x(\phi_2(t))| \int_0^{\alpha(t)} k(t,s) m_3(s) ds dt + b b_3 \int_0^T m_2(t) \int_0^{\alpha(t)} k(t,s) |x(\phi_3(s))| ds dt \\ &\quad + b b_2 b_3 \int_0^T |x(\phi_2(t))| \int_0^{\alpha(t)} k(t,s) |x(\phi_3(s))| ds dt + b_2 \int_0^T m(t) |x(\phi_2(t))| dt \\ &\leq \|m_1\| + \frac{b_1}{B_1} \int_0^T |x(\phi_1(t))| \phi_1'(t) dt + b \int_0^T m_2(t) \int_0^T k(t,s) m_3(s) ds dt \\ &\quad + \frac{b b_2}{B_2} \int_0^T |x(\phi_2(t))| \int_0^T k(t,s) m_3(s) ds dt + b b_3 \int_0^T m_2(t) \int_0^T k(t,s) |x(\phi_3(s))| ds dt + N \int_0^T m_2(t) dt \\ &\quad + b b_2 b_3 \int_0^T |x(\phi_2(t))| \int_0^T k(t,s) |x(\phi_3(s))| ds dt + b_2 N \int_0^T |x(\phi_2(t))| dt \\ &\leq \|m_1\| + \frac{b_1}{B_1} \int_0^{\phi_1(T)} |x(\phi_1(t))| \phi_1'(t) dt + b \int_0^T m_2(t) \int_0^T k(t,s) m_3(s) ds dt \\ &\quad + \frac{b b_2}{B_2} \int_0^{\phi_2(T)} |x(\phi_2(t))| \phi_2'(t) \int_0^T k(t,s) m_3(s) ds dt + \frac{b b_3}{B_3} \int_0^T |x(\phi_2(t))| \phi_2'(t) dt + N \|m_2\| \\ &\quad + \frac{b b_3}{B_3} \int_0^T m_2(t) \int_0^{\phi_3(T)} k(t,s) |x(\phi_3(s))| \phi_3'(s) ds dt \\ &\quad + \frac{b b_2 b_3}{B_2 B_3} \int_0^{\phi_2(T)} |x(\phi_2(t))| \phi_2'(t) \int_0^{\phi_3(T)} k(t,s) |x(\phi_3(s))| \phi_3'(s) ds dt \end{aligned}$$

$$\begin{aligned} &\leq \|m_1\| + \frac{b_1}{B_1} \int_0^T |x(\theta)| d\theta + b M \|m_2\| \|m_3\| + \frac{M b b_2 \|m_3\|}{B_2} \int_0^T |x(\theta)| d\theta + N \|m_2\|, \\ &\quad + \frac{M b b_3 \|m_3\|}{B_3} \int_0^T |x(\theta)| d\theta + \frac{M b b_2 b_3}{B_2 B_3} \int_0^T |x(\theta)| d\theta \cdot \int_0^T |x(\theta)| d\theta + \frac{b_2 N}{B_2} \int_0^{\phi_2(T)} |x(u)| du \\ &\leq \|m_1\| + \frac{b_1}{B_1} \|x\| + b M \|m_2\| \|m_3\| + \frac{M b b_2 \|m_3\|}{B_2} \|x\| + N \|m_2\| \\ &\quad + \frac{M b b_3 \|m_3\|}{B_3} \|x\| + \frac{M b b_2 b_3}{B_2 B_3} \|x\|^2 + \frac{b b_2 N}{B_2} \|x\|. \end{aligned}$$

which shows that the operator A maps the ball B_r into itself with

$$r = \frac{d - \sqrt{d^2 - 4 b M b_2 b_3 B_1^2 B_2 B_3 (\|m_1\| + b M \|m_2\| \|m_3\| + N \|m_2\|)}}{2 b M b_2 b_3 B_1} \quad (4)$$

Assumption (vii) implies

$$0 < d^2 - 4 b M b_2 b_3 B_1^2 B_2 B_3 (\|m_1\| + b M \|m_2\| \|m_3\| + N \|m_2\|) < d^2,$$

which implies that

$$0 < \sqrt{d^2 - 4 b M b_2 b_3 B_1^2 B_2 B_3 (\|m_1\| + b M \|m_2\| \|m_3\| + N \|m_2\|)} < d.$$

Then d is positive and hence r is a positive constant.

Let Q_r be a subset of $B_r \in L_1$ consisting of all functions which are a.e. nondecreasing on I .

Clearly, the set Q_r is nonempty, bounded, convex and closed [4]. Moreover this set is compact in measure [5].

Then we deduce that the operator A maps Q_r into itself.

Since the operator $(Ux)(t) = \int_0^{\alpha(t)} u(t,s,x(s)) ds$ is continuous, then the operator H is continuous and hence the product $G.H$ is continuous. Also, F is continuous. Thus the operator A is continuous on Q_r .

Let X be a nonempty subset of Q_r . Fix $\varepsilon > 0$ and take a measurable subset $D \subset I$ such that $\text{meas } D \leq \varepsilon$. Then, for any $x \in X$, using the same reasoning as in [4] and [5], we get

$$\begin{aligned} \|Ax\|_{L_1(D)} &= \int_D |(Ax)(t)| dt \\ &\leq \int_D |m_1(t)| dt + b_1 \int_D |x(\phi_1(t))| dt \\ &\quad + \int_D (m_2(t) + b_2 |x(\phi_2(t))|) \left[m(t) + b \int_0^{\alpha(t)} k(t,s)(m_3(s) + b_3 |x(\phi_3(s))|) ds \right] dt \\ &\leq \|m_1\|_{L_1(D)} + \frac{b_1}{B_1} \int_D |x(\phi_1(t))| \phi_1'(t) dt + b \int_D m_2(t) \int_0^T k(t,s) m_3(s) ds dt + N \int_D m_2(t) dt \\ &\quad + b b_2 \int_D |x(\phi_2(t))| \int_0^T k(t,s) m_3(s) ds dt + M b b_3 \int_D m_2(t) \int_0^T |x(\phi_3(s))| ds dt \\ &\quad + M b b_2 b_3 \int_D |x(\phi_2(t))| \int_0^T |x(\phi_3(s))| ds dt + \frac{b_2 N}{B_2} \int_D |x(\phi_2(t))| \phi_2'(t) dt \\ &\leq \|m_1\|_{L_1(D)} + \frac{b_1}{B_1} \int_D |x(\theta)| d\theta + b M \|m_2\|_{L_1(D)} \|m_3\|_{L_1} + \frac{M b b_2 \|m_3\|_{L_1}}{B_2} \int_D |x(\theta)| d\theta \\ &\quad + \frac{M b b_3 \|m_2\|_{L_1(D)}}{B_3} \int_0^T |x(\theta)| d\theta + \frac{M b b_2 b_3}{B_2 B_3} \int_D |x(\theta)| d\theta \cdot \int_0^T |x(\theta)| d\theta \\ &\quad + \frac{b_2 N}{B_2} \int_D |x(\theta)| d\theta + N \|m_2\|_{L_1(D)} \\ &\leq \|m_1\|_{L_1(D)} + \frac{b_1}{B_1} \|x\|_{L_1(D)} + b M \|m_2\|_{L_1(D)} \|m_3\|_{L_1} + \frac{M b b_2 \|m_3\|_{L_1}}{B_2} \|x\|_{L_1(D)} \\ &\quad + \frac{M b b_3 \|m_2\|_{L_1(D)}}{B_3} \|x\| + \frac{M b b_2 b_3}{B_2 B_3} \|x\|_{L_1(D)} \cdot \|x\| \\ &\quad + \frac{b_2 N}{B_2} \|x\|_{L_1(D)} + N \|m_2\|_{L_1(D)} \end{aligned}$$

$$\begin{aligned} &\leq \|m_1\|_{L_1(D)} + \frac{b_1}{B_1} \|x\|_{L_1(D)} + M b \|m_2\|_{L_1(D)} \|m_3\|_{L_1} + \frac{M b b_2 \|m_3\|_{L_1}}{B_2} \|x\|_{L_1(D)} \\ &+ \frac{r b M b_3 \|m_2\|_{L_1(D)}}{B_3} + \frac{r b M b_2 b_3}{B_2 B_3} \|x\|_{L_1(D)} \\ &+ \frac{b_2 N}{B_2} \|x\|_{L_1(D)} + N \|m_2\|_{L_1(D)} \end{aligned}$$

Since

$$\lim_{\varepsilon \rightarrow 0} \{ \sup \{ \int_D |m_i(t)| dt : D \subset I, \text{ meas } D < \varepsilon \} \} = 0, \quad i = 1, 2, 3.$$

We obtain

$$\beta(Ax(t)) \leq \left[\frac{b_1}{B_1} + \frac{M b b_2 \|m_3\|_{L_1}}{B_2} + \frac{M b b_2 b_3 r}{B_2 B_3} + \frac{b_2 N}{B_2} \right] \beta(x(t)).$$

This implies

$$\beta(AX) \leq \left[\frac{b_1}{B_1} + \frac{M b b_2 \|m_3\|_{L_1}}{B_2} + \frac{M b b_2 b_3 r}{B_2 B_3} + \frac{b_2 N}{B_2} \right] \beta(X), \quad (5)$$

where β is the De Blasi measure of weak noncompactness.

Keeping in mind Theorem 2 we can write (5) in the form

$$\chi(AX) \leq \left[\frac{b_1}{B_1} + \frac{M b b_2 \|m_3\|_{L_1}}{B_2} + \frac{M b b_2 b_3 r}{B_2 B_3} + \frac{b_2 N}{B_2} \right] \chi(X),$$

where χ is the Hausdorff measure of noncompactness.

Since $\frac{b_1}{B_1} + \frac{M b b_2 \|m_3\|_{L_1}}{B_2} + \frac{M b b_2 b_3 r}{B_2 B_3} + \frac{b_2 N}{B_2} < 1$, from Theorem 3 follows that A is contraction with respect to the measure of noncompactness χ . Thus A has at least one fixed point in Q_r which is a solution of the quadratic functional equation. ■

4 Special cases and applications

As particular cases of Theorem 4 we can obtain theorems on the existence of positive and a.e. nondecreasing solutions belonging to the space $L_1(I)$ of the following quadratic functional equations:

1.If $\psi(t, x) = \int_0^{\alpha(t)} u(t, s, x(\phi_3(s))) ds$, then we obtain the quadratic functional equation [26]

$$x(t) = f(t, x(\phi_1(t))) + g(t, x(\phi_2(t))) \int_0^{\alpha(t)} u(t, s, x(\phi_3(s))) ds, \quad t \in I.$$

2.If $\psi(t, x) = \int_0^t u(t, s, x(\phi_3(s))) ds$, then we obtain the quadratic functional equation

$$x(t) = f(t, x(\phi_1(t))) + g(t, x(\phi_2(t))) \int_0^t u(t, s, x(\phi_3(s))) ds, \quad t \in I.$$

that was studied in [14].

3.If $\psi(t, x) = \int_0^t u(t, s, x(\phi_3(s))) ds$, $g(t, x) = 1$ and $f(t, x) = a(t)$ then we obtain the functional integral equation of Urysohn type

$$x(t) = a(t) + \int_0^t u(t, s, x(\phi_3(s))) ds, \quad t \in I.$$

4.If $f(t, x) = a(t)$, $u(t, s, x) = h(t, x)$, then we obtain the quadratic integral equation

$$x(t) = a(t) + g(t, x(\phi_1(t))) \int_0^t h(s, x(\phi_2(s))) ds, \quad t \in I$$

that was proved in [22].

5.If $g(t, x) = 0$, then we obtain the functional equation

$$x(t) = f(t, x(\phi_1(t))), \quad t \in I$$

which is the same results proved by Banas [4].

6.If $f(t, x) = a(t)$, $u(t, s, x) = k(t, s) h(t, x)$, then we obtain the quadratic integral equation

$$x(t) = a(t) + g(t, x(t)) \int_0^t k(t, s) h(s, x(\phi(s))) ds, \quad t \in I$$

which is the same results proved in [21].

7.If $f(t, x) = a(t)$, $u(t, s, x) = k(t, s) h(t, x)$ and $g(t, x) = 1$ then we obtain the functional integral equation

$$x(t) = a(t) + \int_0^t k(t, s) h(s, x(\phi(s))) ds, \quad t \in I$$

which is the same results proved in [5].

8.If $f(t, x) = 0$ for any $t \in I$ and $x \in R_+$ we have

$$x(t) = g(t, x(\phi_2(t))) \int_0^t u(t, s, x(\phi_3(s))) ds, \quad t \in I.$$

9.If $\phi_i(t) = t$, $i = 1, 2, 3$, $\psi(t, x) = \int_0^t u(t, s, x(s)) ds$ for any $t \in I$ and $x \in R_+$ we have

$$x(t) = f(t, x(t)) + g(t, x(t)) \int_0^t u(t, s, x(s)) ds, \quad t \in I.$$

4.1 Chanrasekhar's integral equation

Example 1:

Let us consider the quadratic integral equation of Volterra type having the form

$$x(t) = a(t) + x(t) \int_0^t \frac{t}{t+s} u(t, s, x(s)) ds. \quad (6)$$

This equation represents the Volterra counterpart of the famous Chandrasekhar quadratic integral equation which has numerous application (cf. [2], [3], [7] and [13]). It arose originally in connection with scattering through a homogeneous semi-infinite plane atmosphere [13]. In astrophysical applications of the Chandrasekhar's equation the only restriction, that $\int_0^1 \phi(s) ds \leq 1/2$ is treated a necessary condition in [12].

In case $a(t) = 1$ and $u(t, s, x(s)) = \lambda \phi(s) x(s)$, λ is a positive constant and on I . Then Eqn.(6) has the form

$$x(t) = 1 + \lambda x(t) \int_0^t \frac{t \phi(s)}{t+s} x(s) ds.$$

In order to apply our results we have to impose an additional condition that the so-called "characteristic" function ϕ need not be continuous on I it sufficient only to be bounded and measurable on I .

In this case $r = \frac{1-\sqrt{1-4\lambda k}}{2\lambda k}$ and the assumption (vii) may be reduced to $4\lambda k \leq 1$ where $\sup_{s \in I} \phi(s) = k$.

Example 2:

Consider the following Chandrasekar's integral equation

$$x(t) = 1 + \frac{1}{10} x(t) \int_0^1 \frac{s}{t+s} x(s) ds, \quad t \in [0, 1] \quad (7)$$

where $\lambda = \frac{1}{2}$, $\phi(s) = \frac{s}{5}$ and $k = \frac{1}{10}$, then the condition $4\lambda k \leq 1$, is satisfied and $r = 4$.

Example 3:

Consider the following Chandrasekar's integral equation

$$x(t) = 1 + \frac{1}{15} x(t) \int_0^1 \frac{\exp(-s)t}{t+s} x(s) ds, \quad t \in \quad (8)$$

where $\lambda = \frac{1}{5}$, $\phi(s) = \frac{1}{3} \exp(-s)$ and $k \leq 1$, then the condition $4\lambda k \leq 1$ is satisfied.

4.2 Fractional integral equation

Example 1:

Now, consider the quadratic integral equation of fractional order

$$x(t) = f(t) + g(t, x(t)) \int_0^t \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} h(s, x(s)) ds, \quad \alpha > 0$$

which was studied in [11] (an existence theorem for continuous solutions was proved) and in [21] (an existence for integrable solutions was proved). In our case, $u(t, s, x(s)) = \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} h(s, x(s))$ and h satisfies

$|h(s, x(s))| \leq m(s) + \frac{b'}{(t-s)^{\alpha-1}} |x|$, $m \in L_1$. Then $|u(t, s, x(s))| \leq \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} m(s) + b|x|$, b is positive constant and $k(t, s) = \frac{(t-s)^{\alpha-1}}{\Gamma(\alpha)} m(s)$. The assumption $\int_0^t k(t, s) ds \leq M$ is reduced by $I^\beta m(t) \leq M'$, $\beta \leq \alpha$. For,

$$\int_0^t k(t, s) ds = I^\alpha m(t) = I^{\alpha-\beta} I^\beta m(t) \leq M' \int_0^t \frac{(t-s)^{\alpha-\beta-1}}{\Gamma(\alpha-\beta)} ds = \frac{M'}{\Gamma(\alpha-\beta+1)} = M$$


Acknowledgment:

The authors are thankful to the referee for the time taken to review this paper and for the remarks that helped improve the quality of this paper.

References

- [1] J. Appell, E. De Pascale, Su. alcuni parametri connessi con la misurabilità non compattezza di Hausdorff in spazi di funzioni misurabili, *Boll. Union Mat. Ital.*, Vol.6 no.3,(1984), 497-515.
- [2] I. K. Argyros, Quadratic equations and applications to Chandrasekhars and related equations, *Bull. Austral. Math. Soc.* 32(1985), 275-292.
- [3] I. K. Argyros, On a class of quadratic integral equations with perturbations, *Funct. Approx.* 20(1992), 51-63.
- [4] J. Banaś, On the superposition operator and integrable solutions of some functional equations, *Nonlin. Analysis T.M.A.* Vol. 12(1988), 777-784.
- [5] J. Banaś, Integrable solutions of Hammerstein and Urysohn integral equations, *J. Austral. Math. Soc. (Series A)* 46 (1989), 61-68.
- [6] J. Banaś, K. Goebel, Measure of noncompactness in Banach space, *Lecture Note in Pure and Appl. Math.*, Vol. 60. Dekker, New York, 1980.
- [7] J. Banaś, M. Lecko, W. G. El-Sayed, Existence Theorems of Some Quadratic Integral Equation, *J. Math. Anal. Appl.* 227 (1998), 276 - 279.
- [8] J. Banaś, A. Martinon, Monotonic solutions of a quadratic integral equation of Volterra type, *Comput. Math. Appl.* 47 (2004), 271 - 279.
- [9] J. Banaś, J. Rocha Martin and K. Sadarangani, On the solution of a quadratic integral equation of Hammerstein type, *Mathematical and Computer Modelling.* vol.43,(2006), 97-104.
- [10] J. Banaś, B. Rzepka, Monotonic solutions of a quadratic integral equations of fractional order *J. Math. Anal. Appl.* 332 (2007), 1370 -1378.
- [11] H. O. Bakodah, The Appearance of Noise Terms in Modified Adomian Decomposition Method for Quadratic Integral Equations, *American Journal of Computational Mathematics*, 2(2012),125-129.
- [12] J. Caballero, A. B. Mingarelli, K. Sadarangani, Existence Of Solutions Of An Integral Equation Of Chandrasekhar Type In The Theory Of Radiative, *Electronic Journal of Differential Equations*, Vol. 2006(2006), No. 57, pp. 111.
- [13] Chandrasekhar, S., Radiative Transfar, Dover, New York, 1960.
- [14] M. Cichon, M. A. Metwali, On quadratic integral equations in Orlicz spaces, *Journal of Mathematical Analysis and Applications*, Volume 387, Issue 1(2012), 419-432.
- [15] F. S. De Blasi, On a property of the unit sphere in Banach spaces, *Math. Soc. Sci. Math. R.S.Roum.*, vol.21, no.3-4, (1977) 259-262.
- [16] K. Deimling, Nonlinear Functional Analysis, *Springer - Verlag, Berlin*, 1985.
- [17] M. A. Darwish, On monotonic solutions of a singular quadratic integral equation with supremum, *Dynam. Syst. Appl.*, 17 (2008), 539-550.
- [18] A.M.A. El-Sayed, M.M. Saleh and E.A.A. Ziada, Numerical and Analytic Solution for Nonlinear Quadratic Integral Equations, *Math. Sci. Res. J.*, 12(8) (2008), 183-191.
- [19] A. M. A. El-Sayed, H. H. G. Hashem, Integrable and continuous solutions of nonlinear quadratic integral equation, *Electronic Journal of Qualitative Theory of Differential Equations* 25(2008), 1-10.

- [20] A. M. A. El-Sayed, H. H. G. Hashem, Monotonic positive solution of nonlinear quadratic Hammerstein and Urysohn functional integral equations, *Commentationes Mathematicae*. Vol.48, No.2(2008), 199-207.
- [21] A. M. A. El-Sayed, H. H. G. Hashem, Monotonic solutions of functional integral and differential equations of fractional order, *E. J. Qualitative Theory of Diff. Equ.*, No. 7. (2009), pp. 1-8.
- [22] El-Sayed, A.M.A and Hashem, H.H.G, Monotonic positive solution of a nonlinear quadratic functional integral equation, *Appl. Math. and Comput.*, 216 (2010) 2576-2580.
- [23] El-Sayed, A.M.A, Hashem, H.H.G and Ziada E. A. A. Picard and Adomian Methods for quadratic integral equation, *Comp. Appl. Math.*, Vol. 29, N. 3, (2010), 447-463.
- [24] A.M.A. El-Sayed, H. H. G. Hashem and Y. M. Y. Omar, Positive continuous solution of a quadratic integral equation of fractional orders, *Math. Sci. Lett.* Vol 2, No.1, 1-9 (2013).
- [25] H.H.G Hashem and M.S. Zaki, Carathéodory theorem for quadratic integral equations of Erdélyi-Kober type, *Journal of Fractional Calculus and its Applications*, Vol. 4 No. 5(2013)1-8.
- [26] H. H. G. Hashem and A. R. Al-Rwaily, Asymptotic Stability of Solutions to a Nonlinear Urysohn Quadratic Integral Equation, *Hindawi Publishing Corporation International Journal of Analysis Volume 2013, Article ID 259418*, 7 pages.
- [27] W. G. El-Sayed, B. Rzepka, Nondecreasing Solutions of a Quadratic Integral Equation of Urysohn Type, *Comput. Math. Appl.*, 51 (2006) 1065-1074.



differential and integral equations of fractional orders. She is referee and member of the editorial board of several international journals in the frame of pure and applied mathematics.

Hind Hashem

received the PhD degree in Mathematics at Alexandria University, Egypt. Here research interests are in the areas of pure mathematics (quadratic integral equations of different kinds, coupled systems of differential and integral equations, functional



several international journals in the frame of pure and applied mathematics. His main research interests are: dynamical systems, fractional calculus, modeling, computational methods, the fractional-order logistic equation, Equilibrium points, stability and numerical solutions of fractional-order predator-prey and rabies models, fractional and integral equations.

Ahmed El-Sayed

is Professor of pure mathematics at Alexandria University, faculty of Science, Mathematics and computer science department, Egypt. He is the Managing Editor of Journal of Fractional Calculus and Applications. He is referee and Editor of